

## Response of NPK and Vermicompost on Soil Properties, Growth and Yield Attributes of Cluster Bean (*Cyamopsis tetragonoloba* L.) Cv. Pusa Navbahar

### ABSTRACT

An experiment was conducted during *Kharif* (July to October) **on season** 2022-24 to study “Response of different level of NPK and vermicompost on soil properties, growth and yield attributes of cluster bean (*Cyamopsis tetragonoloba* L.) cv. Pusa Navbahar” at the research farm of Soil Science and Agricultural Chemistry, design laid out in randomized block design (RBD) with three replications. Cluster bean was taken for study with recommended doses of fertilizers (NPK @ 0, 50, and 100%  $\text{ha}^{-1}$ ) was applied along with Vermicompost (Vermicompost @ 0, 50 and 100%  $\text{ha}^{-1}$ ). A variety of cluster bean, Pusa Navbahar was taken for a research trial. Bulk density ( $\text{mg m}^{-3}$ ) and Particle density ( $\text{mg m}^{-3}$ ) was recorded maximum in T<sub>1</sub> (Absolute Control) and minimum in T<sub>9</sub> (NPK @ 100% + Vermicompost @ 100%). pH (1:2.5) w/v was recorded maximum in T<sub>1</sub> (Absolute Control) and minimum in T<sub>9</sub> (NPK @ 100% + Vermicompost @ 100%). Pore space (%), Water holding capacity (%), EC ( $\text{dS m}^{-1}$ ), Organic carbon (%), Nitrogen ( $\text{kg ha}^{-1}$ ), Phosphorus ( $\text{kg ha}^{-1}$ ) and Potassium ( $\text{kg ha}^{-1}$ ) was measured maximum in T<sub>9</sub> (NPK @ 100% + Vermicompost @ 100%) and minimum in T<sub>1</sub> (Absolute Control).

*Keywords: cluster bean, vermicompost, nitrogen, phosphorus, potassium*

### 1. INTRODUCTION

Soil is defined as the unconsolidated upper part of the earth's crust that serves as natural medium for the growth of higher plants [1]. The soil characteristics is categorized into three viz: physical, chemical and biological which interact to sustain the growth and development of crops. When the soil is properly managed, crop productivity is sustained. Soil physical and chemical properties play a central role in crop growth, development and yield by influencing the availability of air, nutrients and water to crops [2]. Cluster bean (*Cyamopsis tetragonoloba* L.) belongs to the family Fabaceae and subfamily Faboideae. It is commonly known as guar, an annual legume crop, are widely grown for its guar gum that contain ample amount of soluble dietary fibres. Cluster bean is a cash crop due to its application in agriculture, paper, juice, textile, purification of waste water, mining, petroleum, explosives, pharmaceuticals and food industries. Globally, India ranks first and produces about 80% of the world's cluster beans. Major cluster bean producing states in India are Rajasthan, Haryana, Gujarat, Uttar Pradesh, Punjab, and Madhya Pradesh. This legume is a valuable plant in a crop rotation cycle, as it reduces weeds and enrich the soil by partnering with nitrogen fixing bacteria in the soil. Being a legume crop, it has the capacity to fix atmospheric nitrogen by its effective root nodules. Hence, guar is considered as an excellent soil building crop, like other legumes, with respect to available nitrogen, which improve yield of succeeding crops [3]. Cluster beans or guar, among leguminous crops is comparatively more drought tolerant hardy crop, grown under rainfed condition in arid and semi-regions of India during *kharif* season. The crop can tolerate moderate salinity and alkalinity conditions. Its deep penetrating roots enable the plant to utilize available moisture more efficiently and thus offer better scope for rainfed cropping. Due to high degree of drought and salinity tolerance, guar could be a valuable alternative crop for the exploitation of the semi-arid environment, where high temperature, poor erratic rainfall and elevated soil salt content restrict the cultivation of other crops [4]. Nitrogen is considered as one of the significant macronutrients among all the mineral elements for all living tissues of the plant from metabolism to resource allocation, growth and development. It increases the protein content in grains, fruits and seeds of plants. Deficient plants may have stunted growth due to decreasing photosynthesis, leaf area and longevity of green leaves. Phosphorus plays a vital role in photosynthesis,

respiration, energy storage, cell elongation and improves the quality of crops. Potassium is an important essential nutrient after nitrogen and phosphorus and plays a vital role in plant cell sap, support enzymatic activity, photosynthesis and transportation of sugar, synthesis of protein and starch but doesn't bound with carbon or oxygen. It develops tolerance to drought condition and enhances plant ability to resist attacks of pests and diseases. Vermicompost is a peat-like organic fertilizer with high nutritional contents, aeration, porosity, and water-holding capacity, prepared by the joint action of earthworms and microbes. In addition to organic waste management, vermicompost is recognized as an effective plant growth promoter [5].

## 2. MATERIAL AND METHODS



The experiment was conducted during kharif season in 2023 at the central research farm of department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute (NAI), Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U.P., situated on the south of Prayagraj on the right side of the river Yamuna on the South of Rewa Road at a distance of about 6 km away from Prayagraj city. It is situated at 25°57'69" N latitude, 81°59'74" E longitude and at the altitude of 98 meter above the sea level. The area of Prayagraj district comes under subtropical belt with a sub-tropical and semi-arid climate in the South east of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46°C – 48°C and seldom falls as low as 4°C – 5°C. The relative humidity ranged between 20 to 94 percent. The average rainfall in this area is around 1100 mm annually and rains mostly during July- September. The soil samples were randomly collected from three different sites in the experiment plot prior to tillage operation from a depth of 0-15 cm. The size of the soil sample was reduced by coning and quartering the composites soil sample and was air dried, passed through a 2 mm sieve for preparing the sample for physical and chemical analysis [6] (bulk density, particle density, water holding capacity and pore space: Muthuvel et al. [7]; pH: Jackson [8]; EC: Wilcox [9]; organic carbon: Walkley and Black [10]; available nitrogen: Subbaih and Asija [11], phosphorus: Olsen et al. [12]; potassium: Toth and Prince, [13]).

### 2.1 Treatment Combination

**Table 1: Treatment combinations of cluster bean**

| Treatment      | Treatment Combination            | Symbol                        |
|----------------|----------------------------------|-------------------------------|
| T <sub>1</sub> | Control                          | L <sub>0</sub> V <sub>0</sub> |
| T <sub>2</sub> | NPK @ 0% + Vermicompost @ 50%    | L <sub>0</sub> V <sub>1</sub> |
| T <sub>3</sub> | NPK @ 0% + Vermicompost @ 100%   | L <sub>0</sub> V <sub>2</sub> |
| T <sub>4</sub> | NPK @ 50% + Vermicompost @ 0%    | L <sub>1</sub> V <sub>0</sub> |
| T <sub>5</sub> | NPK @ 50% + Vermicompost @ 50%   | L <sub>1</sub> V <sub>1</sub> |
| T <sub>6</sub> | NPK @ 50%+ Vermicompost @ 100%   | L <sub>1</sub> V <sub>2</sub> |
| T <sub>7</sub> | NPK @ 100% + Vermicompost @ 0%   | L <sub>2</sub> V <sub>0</sub> |
| T <sub>8</sub> | NPK @ 100% + Vermicompost @ 50%  | L <sub>2</sub> V <sub>1</sub> |
| T <sub>9</sub> | NPK @ 100% + Vermicompost @ 100% | L <sub>2</sub> V <sub>2</sub> |

Note: RDF: - 20:40:40 NPK, Vermicompost 4 t ha<sup>-1</sup>

## 3. RESULTS AND DISCUSSION


### 3.1 Response of N, P, K and Vermicompost on physical properties of soil after harvest of cluster bean

The data of soil samples showed maximum bulk density (1.35 Mg m<sup>-3</sup> and 1.36 Mg m<sup>-3</sup>) and particle density (2.55 Mg m<sup>-3</sup> and 2.55 Mg m<sup>-3</sup>) at 0-15 cm and 15-30 cm depth, respectively that were recorded in the treatment T<sub>1</sub> (Absolute control) and minimum bulk density (1.27 Mg m<sup>-3</sup> and 1.29 Mg m<sup>-3</sup>) and particle density (2.51 Mg m<sup>-3</sup> and 2.52 Mg m<sup>-3</sup>) at 0-15 cm and 15-30 cm depth, respectively that were recorded in the treatment T<sub>9</sub> (NPK @ 100% + Vermicompost @ 100%). The soil samples data also showed maximum percentage pore space (49.60 and 48.81 %) and water holding capacity (46.93 and


47.05 %) at 0-15 cm and 15-30 cm depth, respectively that were measured in the treatment T<sub>9</sub> (NPK @ 100% + Vermicompost @ 100%) and minimum percentage pore space (47.06 % and 46.67 %) and water holding capacity (42.79 % and 43.05 %) was recorded in treatment T<sub>1</sub> (Absolute control) at 0-15 cm and 15-30 cm depth, respectively (Table 2 and fig. 2). Similar findings were also observed [14], [15] and [16].

**Table 2. Effect of Different Levels of N, P, K and Vermicompost on physical properties of post-harvest soil sample of cluster bean**

| Treatments     | Bulk Density (Mg m <sup>-3</sup> ) |         | Particle Density (Mg m <sup>-3</sup> ) |         | Pore space (%) |         | Water holding capacity (%) |         |
|----------------|------------------------------------|---------|--|---------|----------------|---------|----------------------------|---------|
|                | 0-15cm                             | 15-30cm | 0-15cm                                 | 15-30cm | 0-15cm         | 15-30cm | 0-15cm                     | 15-30cm |
| T <sub>1</sub> | 1.35                               | 1.36    | 2.55                                   | 2.55    | 47.06          | 46.67   | 42.79                      | 43.05   |
| T <sub>2</sub> | 1.32                               | 1.34    | 2.54                                   | 2.54    | 48.03          | 47.24   | 43.96                      | 44.75   |
| T <sub>3</sub> | 1.29                               | 1.31    | 2.53                                   | 2.54    | 49.01          | 48.42   | 45.16                      | 46.07   |
| T <sub>4</sub> | 1.34                               | 1.35    | 2.54                                   | 2.55    | 47.24          | 46.85   | 42.15                      | 42.96   |
| T <sub>5</sub> | 1.31                               | 1.33    | 2.53                                   | 2.53    | 48.22          | 47.43   | 43.16                      | 44.17   |
| T <sub>6</sub> | 1.28                               | 1.30    | 2.52                                   | 2.53    | 49.21          | 48.62   | 45.18                      | 45.70   |
| T <sub>7</sub> | 1.33                               | 1.34    | 2.53                                   | 2.53    | 47.43          | 47.04   | 43.00                      | 43.75   |
| T <sub>8</sub> | 1.30                               | 1.32    | 2.52                                   | 2.52    | 48.41          | 47.62   | 44.14                      | 44.95   |
| T <sub>9</sub> | 1.27                               | 1.29    | 2.51                                   | 2.52    | 49.60          | 48.81   | 46.93                      | 47.05   |
| F- test        | NS                                 | NS      | NS                                     | NS      | S              | S       | S                          | S       |
| S.Em. (±)      | -                                  | -       | -                                      | -       | 0.46           | 0.45    | 0.55                       | 0.67    |
| C.D (P=0.05)   | -                                  | -       | -                                      | -       | 1.41           | 1.38    | 1.67                       | 2.03    |



**Fig. 1. Effect of Different Levels of N, P, K and Vermicompost on Bulk density and Particle density of the post-harvest soil sample**



**Fig. 2. Effect of Different Levels of N, P, K and Vermicompost on Pore space and Water holding capacity of the post-harvest soil sample**

### 3.2 Response of N, P, K and Vermicompost on chemical properties of soil after harvest of cluster bean

Table 3. and fig. 3 showed that highest pH (6.88 and 6.89) but lowest EC  $0.43$  and  $0.44 \text{ dS m}^{-1}$  and organic carbon (0.34 and 0.33 %) was recorded in treatment T<sub>1</sub> (Absolute control) at 0-15 cm depth and at 15- 30 cm depth, respectively and T<sub>9</sub> (NPK @ 100% + Vermicompost @ 100%) resulted the lowest pH (6.80 and 6.82) but highest EC ( $0.46 \text{ dS m}^{-1}$  and  $0.47 \text{ dS m}^{-1}$ ), organic carbon (0.38% and 0.37%) at 0-15 cm depth and at 15- 30 cm depth, respectively. Table 4. and fig. 4 showed that lowest available nitrogen ( $265.51$  and  $263.29 \text{ kg ha}^{-1}$ ), available phosphorus ( $14.87$  and  $14.27 \text{ kg ha}^{-1}$ ) and available potassium ( $178.36$  and  $172.66 \text{ kg ha}^{-1}$ ) was measured in treatment T<sub>1</sub> (Absolute control) at 0-15 cm depth and at 15- 30 cm depth, respectively and T<sub>9</sub> (NPK @ 100% + Vermicompost @ 100%) recorded the highest available nitrogen ( $288.05$  and  $285.22 \text{ kg ha}^{-1}$ ), available phosphorus ( $25.28$  and  $24.28 \text{ kg ha}^{-1}$ ) and available potassium ( $201.36$  and  $193.73 \text{ kg ha}^{-1}$ ) at 0-15 cm depth and at 15- 30 cm depth, respectively. Similar findings were also observed [13], [14] and [15].

**Table 3. Effect of Different Levels of N, P, K and Vermicompost on pH, EC and Organic carbon of the post-harvest soil sample**

| Treatments     | pH     |         | Electrical conductivity ( $\text{dS m}^{-3}$ ) |         | Organic carbon (%) |         |
|----------------|--------|---------|--|---------|--------------------|---------|
|                | 0-15cm | 15-30cm | 0-15cm   | 15-30cm | 0-15cm             | 15-30cm |
| T <sub>1</sub> | 6.88   | 6.89    | 0.43   | 0.44    | 0.34               | 0.33    |
| T <sub>2</sub> | 6.86   | 6.87    | 0.44   | 0.45    | 0.35               | 0.34    |
| T <sub>3</sub> | 6.82   | 6.84    | 0.45   | 0.46    | 0.36               | 0.35    |
| T <sub>4</sub> | 6.87   | 6.88    | 0.43   | 0.44    | 0.35               | 0.34    |
| T <sub>5</sub> | 6.85   | 6.86    | 0.44   | 0.45    | 0.36               | 0.35    |
| T <sub>6</sub> | 6.81   | 6.83    | 0.45   | 0.46    | 0.37               | 0.36    |
| T <sub>7</sub> | 6.86   | 6.87    | 0.44   | 0.45    | 0.36               | 0.35    |
| T <sub>8</sub> | 6.84   | 6.85    | 0.45   | 0.46    | 0.37               | 0.36    |
| T <sub>9</sub> | 6.80   | 6.82    | 0.46   | 0.47    | 0.38               | 0.37    |

|                                 |           |           |           |           |          |          |
|---------------------------------|-----------|-----------|-----------|-----------|----------|----------|
| <b>F- test</b>                  | <b>NS</b> | <b>NS</b> | <b>NS</b> | <b>NS</b> | <b>S</b> | <b>S</b> |
| <b>S.Em. (<math>\pm</math>)</b> | -         | -         | -         | -         | 0.01     | 0.01     |
| <b>C.D (P=0.05)</b>             | -         | -         | -         | -         | 0.02     | 0.02     |





Fig. 3. Effect of Different Levels of N, P, K and Vermicompost on pH, EC and Organic carbon of the post-harvest soil sample

Table 4. Effect of Different Levels of N, P, K and Vermicompost on Available Nitrogen, Available Phosphorus and Available Potassium of the post-harvest soil sample

| Treatments                      | Available Nitrogen (kg ha <sup>-1</sup> ) |          | Available Phosphorus (kg ha <sup>-1</sup> ) |          | Available Potassium (kg ha <sup>-1</sup> ) |          |
|---------------------------------|---|----------|---|----------|--|----------|
|                                 | 0-15cm                                    | 15-30cm  | 0-15cm                                      | 15-30cm  | 0-15cm                                     | 15-30cm  |
| <b>T<sub>1</sub></b>            | 265.51                                    | 263.29   | 14.87                                       | 14.27    | 178.36                                     | 172.66   |
| <b>T<sub>2</sub></b>            | 268.14                                    | 266.02   | 15.36                                       | 15.26    | 180.46                                     | 173.58   |
| <b>T<sub>3</sub></b>            | 271.09                                    | 268.06   | 16.89                                       | 16.38    | 184.68                                     | 174.90   |
| <b>T<sub>4</sub></b>            | 274.37                                    | 271.54   | 15.27                                       | 15.02    | 186.79                                     | 179.56   |
| <b>T<sub>5</sub></b>            | 276.49                                    | 274.47   | 18.29                                       | 17.01    | 187.46                                     | 181.67   |
| <b>T<sub>6</sub></b>            | 279.25                                    | 276.41   | 23.27                                       | 22.45    | 189.89                                     | 183.59   |
| <b>T<sub>7</sub></b>            | 282.49                                    | 280.24   | 17.28                                       | 16.25    | 196.97                                     | 185.23   |
| <b>T<sub>8</sub></b>            | 285.34                                    | 283.32   | 20.49                                       | 19.26    | 198.98                                     | 192.43   |
| <b>T<sub>9</sub></b>            | 288.05                                    | 285.22   | 25.28                                       | 24.28    | 201.36                                     | 193.73   |
| <b>F- test</b>                  | <b>S</b>                                  | <b>S</b> | <b>S</b>                                    | <b>S</b> | <b>S</b>                                   | <b>S</b> |
| <b>S.Em. (<math>\pm</math>)</b> | 3.37                                      | 4.03     | 0.26  | 0.28     | 2.98                                       | 2.62     |
| <b>C.D (P=0.05)</b>             | 10.20                                     | 12.10    | 0.79  | 0.84     | 8.96                                       | 7.90     |



**Fig. 4. Effect of Different Levels of N, P, K and Vermicompost on Available Nitrogen, Available Phosphorus and Available Potassium of the post-harvest soil sample**

#### 4. CONCLUSION

Based on the results, the application of NPK and Vermicompost was found to improve the soil's health in references to cluster bean. Application of T<sub>9</sub> (@ 100% NPK + @ 100% Vermicompost) was found optimal for improving Soil Properties like Pore space, Water holding capacity, Organic Carbon and Available Nitrogen, Phosphorus, Potassium. The results showed that the soil's bulk density, particle density, pH, and electrical conductivity were found non-significant after the crop was harvested, while the soil's organic carbon, nitrogen, phosphorus, nitrogen, porosity, and water holding capacity were found significant after the crop was harvested with the various applications of the N, P, K and vermicompost.

#### REFERENCES

1. Ahmed, N., Masood, S., Ahmad, S. and Bashir, S. Soil Management for Better Crop Production and Sustainable Agriculture. *Agronomic Crops* 2019; 47-71.
2. Cardone, L., Castronuovo, D., Perniola, M., Scrano, L., Cicco, N. and Candido, V. The Influence of Soil Physical and Chemical Properties on Saffron (*Crocus sativus* L.) Growth, Yield and Quality. *Agronomy* 2020; 10(8): 1154.

3. El-Lethy, S. R., Talaat, I. M., Shahira, A. T. and El-Moursi, A. Impacts of some biostimulants on guar (*Cyamopsis tetragonoloba* L.) plants. *Bulletin of the National Research Centre*. 2019; 43: 169.
4. Choudhary, B., Bamboriya, S. J., Thomas, T., Sarita, and Bijarnia, L. S. Effect of NPK and Neem Cake on Physical and Chemical Properties of Soil in Cluster bean (*Cyamopsis tetragonoloba* L.) Cv. Pusa nav-bahar. *International Journal Current Microbiological Application Science*. 2020; 9(4): 1777-1783.
5. Rehman, S. U., Castro, F. D., Aprile, A., Benedetti, M., Fanizzi, F. P. Vermicompost: Enhancing Plant Growth and Combating Abiotic and Biotic Stress. *Agronomy* 2023; 13(4): 1134.
6. Joshi, A., Thomas, T., David, A. A., Kumar, S. and Dangi, M. Response of vermicompost and neem cake on Soil health and yield attributes of cluster bean (*Cyamopsis tetragonoloba* L.) Var. Durgapura Safed. *International Journal of Chemical Studies*. 2020; 8(4): 3349-3353.
7. Muthuval, P. S, Udaysoorian, C., Natesan, R., Ramaswami, P. P. *Introduction to Soil analysis*, Tamil Nadu Agricultural University, Coimbatore, 641002; 1992.
8. Jackson, M. L. Soil chemical analysis, Second edition Indian Reprint, prentice hall of India, New Delhi. 1958; pp-498.
9. Wilcox L. V. Electrical conductivity. *Am. Water work Assoc. J.* 1950; 42:775 -776.
10. Walkley, A. and Black, I. A. Critical examination of rapid method for determining organic carbon in soils, effect of variance in digestion conditions and of inorganic soil constituents. *Soil science* 1947; pp. 632-251.
11. Subbiah, B. V., Asija, L. A. (1956) Rapid procedure for estimation of available nitrogen in soils. *Current Science*. 1956, 25:259.
12. Olsen, S. R., Cole, C. V., Watanhe, F. S. and Dean, L. A. Estimation of available Phosphorus in soils by extraction with sodium bicarbonate. U. S. Dept. Agri. Circ. 1954; 939.
13. Toth, S. J., Prince, A. L. Estimation of Cation Exchange Capacity and exchangeable Ca, K and Na Content of Soil by Flame Photometer technique. *Soil Science*. 1949; 6(7): 439- 445.
14. Kumari, U., Hasan, A., Thomas, T., David, A. A., Khatana, R. N. S., and Kumar, U. Impact of vermicompost and zinc implementation on soil physico-chemical properties after harvest of mung bean. *Asian Journal of Microbiology Biotechnology and Environmental Sciences*. 2022; 24(3): 519-522.
15. Dangi, M., Thomas, T., David, A. A., Kumar, S. and Joshi, A. Response of different levels of Zinc and Vermicompost on Soil health and yield attributes of Cluster bean (*Cyamopsis tetragonoloba* L.) viz. Durgapura Jay. *International Journal of Chemical Studies*. 2020; 8(4): 3559-3563.
16. Yadav M. K., Tarence, T. and Rao P. S. Response of NPK and Neem Cake on Soil Properties, Growth and Yield of Cluster Bean (*Cyamopsis tetragonoloba* L.) Var. Laxmi-50. *International Journal of Current Microbiology and Applied Sciences*. 2019; 8(5): 1752-1758.