

“Effect of different plant growth regulators on seed germination, seedling growth and establishment of papaya (*Carica papaya* L.) c.v. Taiwan-7”

ABSTRACT:-

An experiment was conducted at the Department of Horticulture, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (U.P)-211007, India during year 2018-19 to investigate the “Effect of different plant growth regulators on seed germination, seedling growth and establishment of papaya (*Carica papaya* L.) c.v. Taiwan-7, under Prayagraj agro climatic conditions”. The experiment was laid out in Randomized Design (RBD). There were Ten treatments including control replicated three times in which growing media such as pond soil, vermicompost, and Plant growth regulators mixed in various percentages. The different levels of plant growth regulators Such as GA₃, NAA and BA with different composition in different combination showed positive response on physical quality attributes of Papaya plant. The maximum values of physical quality characters of papaya plant like Survival percentage (90.68), number of germination days (10.01), germination percentage (90.01), number of leaves (10.46), plant height (12.76 cm), leaf area (38.4 cm²) and chlorophyll content (52.84), Number of primary roots (10.46), number of secondary roots(27.68) and stem diameter (3.85 cm) were recorded in T3 (GA₃ with 150 ppm) and followed by the treatment T6 (NAA with 150 ppm) while, minimum values of these characters maximum value of physical parameter was recorded in control treatment (T₀). From the present investigation it is concluded that treatment (T3) Performed best in terms of growth parameters followed by (T6) and (T9). The maximum B:C ratio was also observed in the same treatment i.e. (T3) with **2.02** .

Key words: Plant growth regulators, Vermicompost, Taiwan-7, GA₃, NAA-

1. INTRODUCTION:-

Papaya (*Carica papaya* L.): A tropical fruit crop with rapid growth, continuous fruiting cycle, and evergreen nature. It is an exceptionally wholesome, refreshing, and delicious tropical fruit that belongs to the Caricaceae family. Native to tropical America, papaya is also known by

Formatted: Subscript

Commented [h1]: Rewrite it as per format.
Remove title
Focus on results part

Commented [h2]: Correct it

the names papaw or pawpaw. It stands as an ideal fruit choice for cultivation in various settings, including kitchen gardens, residential backyards, fields, and even orchards as a filler plant. Its proximity to urban areas makes it a convenient choice for cultivation, and its nutritional richness further contributes to its popularity.

Nutritional Value and Richness: Papaya stands out as a nutritional powerhouse, offering a plethora of health benefits. It contains approximately 2500 International Units (IU) of vitamin A and around 85 milligrams of vitamin C per 100 grams of its succulent pulp. These vitamins are essential for maintaining healthy skin, boosting the immune system, and supporting overall well-being. In addition, papaya is notably rich in calcium and a range of other essential minerals that contribute to bone health, nerve function, and more. **Medicinal Significance:** Beyond its delectable taste and nutritional bounty, papaya holds high medicinal value. The extraction of papain from the dried latex of immature papaya fruits serves a variety of applications. Papain, a proteolytic enzyme, is used to tenderize meat, manufacture chewing gum, create cosmetics, degum natural silk, and enhance the shrink resistance of wool. The diverse utility of papain highlights its significance in industries beyond the culinary realm. **Anticancer Potential:** Recent research has uncovered another facet of papaya's potential health benefits. Papaya juice has demonstrated an *in vitro* antiproliferative effect on liver cancer cells. This effect is believed to be attributed to compounds like lycopene present in papaya. The promising findings suggest that papaya may have a role to play in promoting health and potentially combating certain diseases.

Soil Requirements and Climatic Considerations: **Optimal Soil Conditions:** Papaya thrives in well-drained and fertile soils, which provide the ideal foundation for its growth and development. Well-drained soils prevent waterlogging, which can be detrimental to the root system. Fertile soils offer the necessary nutrients for papaya plants to flourish.

The soil structure should allow for efficient root penetration, promoting healthy root growth and nutrient absorption. **Climatic Preferences and Protection:** Papaya plants have specific climatic preferences that contribute to their successful cultivation. They thrive in sheltered environments that offer protection against adverse wind conditions. While papaya requires abundant sunlight for growth, exposure to strong winds can damage the plants and hinder their development. By choosing a sheltered location with ample sunlight, cultivators can provide the optimal climatic conditions for papaya to thrive. **Seedling Raising Time and Challenges:** Best

Raising Time: Optimal seedling raising time for papaya is June to September, with an alternative window of February to May in North India due to frost.

Naphthalene Acetic Acid (NAA) is a synthetic plant growth regulator with significant effects on plant growth and development: Root Development: NAA promotes the formation and growth of roots. It stimulates cell elongation and division in root tissues, leading to the development of a robust root system. Lateral Root Formation: NAA is involved in the initiation and development of lateral roots, which branch from the primary root. This branching increases the root surface area, enhancing nutrient and water absorption. Fruit Growth: NAA can influence fruit development by preventing premature fruit drop and enhancing fruit growth. Its application can lead to increased fruit size and yield in specific crops

Gibberellic Acid (GA) is a plant growth regulator with diverse roles in influencing plant development: Stem Elongation: GA stimulates cell division and elongation in the internodes, leading to increased stem length and overall plant height. Seed Germination: GA breaks seed dormancy and promotes germination by activating enzymes that degrade stored nutrients, facilitating embryo growth. Flowering Induction: GA interacts with other hormones and environmental cues to influence flowering. It's widely used in horticulture to trigger flowering in certain plants.

The Taiwan-7 papaya variety, scientifically known as *Carica papaya* L., is characterized by its distinct features: Fruit Colour: The standout feature of this variety is its vibrant red-orange flesh, setting it apart from traditional yellow or orange varieties. Flavour and Taste: "Taiwan-7" papaya is known for its sweet and tropical flavour, making it a favourite for consumption. Fruit Size: The fruits of this variety are medium to large in size, making them suitable for both individual consumption and commercial use. Yield Potential: This variety boasts high yield potential, making it an attractive option for farmers seeking productive cultivars.

Commented [h3]: Merge short sentences

2. MATERIALS AND METHODS:-

2.1 Geographical location of the experimental site

The experimental site is located at a latitude of 25.41° North and longitude of 81.84 ° East, with an altitude of 98 meters above the mean sea level (MSL). [ADD MORE DETAILS](#)

Formatted: Font color: Red

2.2 Climatic conditions of the experimental area

The area of Prayagraj comes under humid sub-tropical climate, which experiences warm humid monsoon, hot dry summer and cold dry winter. The annual mean temperature is 26.1°C while monthly mean temperatures are 18-29°C. The daily average maximum temperature is about 22°C and the minimum temperature is 9°C. The average annual rainfall received is 1042.2 mm. At this location, the temperature reaches upto 46°C-48°C and the minimum temperature recorded is 4°C-5°C. The relative humidity ranges in this location ranges between 20-94%.

2.3 Experimental details

Formatted: Body Text, Right: 0.08", Space After: 12 pt, Add space between paragraphs of the same style, No bullets or numbering, Tab stops: Not at 0.1"

Table 1 Treatment Details

TREATMENT-(N0.)	Treatments
cT0	Control (Unsprayed)
T ₁	GA3@50ppm
T ₂	GA3@100ppm
T ₃	GA3@150ppm
T ₄	NAA@50ppm
T ₅	NAA@100ppm
T ₆	NAA@150ppm
T ₇	BA@50ppm
T ₈	BA@100ppm
T ₉	BA@150ppm

3. RESULTS AND DISCUSSION

a) Number of days to germination

The minimum for Number of days required for initiation of germination was recorded in the treatment T3 (GA3 @ 150 ppm) 10.01 followed by 10.68 with T6 (NAA @ 150 ppm) and maximum days for initiation of Germination 12.68 was recorded in T0 control.

b) Number of leaves per plant

At 45 DAT the maximum Number of leaves per plant was recorded in the treatment T4 (GA3 @ 150 ppm) 10.46 followed by 10.40 with T6 (NAA @ 150 ppm) and minimum Number of leaves per plant 8.36 was recorded in T0 control.

c) Height of seedling

At 45 days the maximum Height of seedling was recorded in the treatment T4 (GA3 @ 150 ppm) 12.76 cm followed by 12.00 cm with T6 (NAA @ 150 ppm) and minimum Height of seedling 9.96 cm was recorded in T0 control.

d) Stem Diameter

The maximum Stem Diameter was recorded in the treatment T4 (GA3@ 150 ppm) 3.85 mm followed by 3.67 mm with T6 (NAA @ 150 ppm) and minimum Stem Diameter 3.37 mm was recorded in T0 control

e) Fresh weight of shoot (g)

At 45 DAT the maximum Fresh weight of shoot (g) was recorded in the treatment T4 (GA3@ 150 ppm) 5.05 g followed by 4.60 g with T6 (NAA @ 150 ppm) and minimum Fresh weight of shoot (g) 3.98 was recorded in T0 control.

f) Fresh weight of root (g)

Fresh weight of root (g) of four tagged seedlings was measured by metric scale at 45 days and average value was Calculated. The maximum Fresh weight of root (g) was recorded in the treatment T4 (GA3 @ 150 ppm) 1.26 g followed by 1.09 g with T6 (NAA @ 150 ppm) and minimum days for Fresh weight of shoot (g) 0.60 was recorded in T0 control.

g) Leaf area

The maximum Leaf Area was recorded in the treatment T4 (GA3 @ 150 ppm) 38.40 cm² / plant followed by 38.21 cm² / plant with T8 (NAA @ 150 ppm) and minimum Leaf Area 33.69 cm² / plant was recorded in T0 control.

h) Survival percentage of papaya seedling

The maximum Survival percentage of papaya seedling was recorded in the treatment T4 (GA3 @ 150 ppm) 90.86 % followed by 87.98% with T6 (NAA @ 150 ppm) and minimum Survival percentage of papaya seedling 77.79% was recorded in T0 control.

i) Germination Percentage

The germination percentage was worked out after the final germination, i.e., after stoppage of germination. It was calculated by dividing to total number of seeds sown with the number of seeds germination and germination percent was calculated

The maximum Germination Percentage was recorded in the treatment T3 (GA3 @ 150 ppm) 90.01 % and minimum Germination Percentage 60.01% was recorded in T0 control.

j) Chlorophyll content

Chlorophyll content of four tagged seedlings was measured with the help of spectrophotometric ally. An average value was calculated. The maximum Chlorophyll Content was recorded in the treatment T4 (GA3 @ 150 ppm) 52.84 µm/g followed by 51.82 µm/g with T8 (NAA @ 150 ppm) and minimum Chlorophyll Content 45.17 µm/g was recorded in T0 control.

Table-2 Effect of PGRs on germination percentage, days to germination, plant height (cm) and survival percentage in papaya

Treatment (No.)	Treatment	Germination Percentage	Number of Days of Germination	Survival Percentage (%)	Plant Height (cm)
T ₀	Control	60.01	12.68	77.79	9.96
T ₁	GA3@50ppm	68.34	11.34	84.34	10.23
T ₂	GA3@100ppm	73.34	11.01	87.01	10.46
T ₃	GA3@150ppm	90.01	10.01	90.68	12.76
T ₄	NAA@50ppm	66.68	12.34	82.34	10.03
T ₅	NAA@100ppm	70.01	11.01	85.72	10.3
T ₆	NAA@150ppm	86.68	10.68	87.98	12
T ₇	BA@50ppm	61.13	12.68	81.34	10
T ₈	BA@100ppm	68.34	11.34	85.01	10.06
T ₉	BA@150PPM	75.01	10.68	87.68	11.16
F test		S	S	S	S
SE(m)		3.11	0.29	1.18	0.31
C.D. at 5%		1.56	0.14	0.59	0.15
C.V		13.67	8	4.4	9.08

Table 3 Effect of PGRs on Stem diameter (cm), Leaf area and number of leaves in papaya.

Treatment (No.)	Treatment	Number of leaves	Stem Diameter (mm)	Chlorophyll (SPAD Value)	Leaf Area(cm ²)
T ₀	Control	8.36	3.37	45.17	33.69
T ₁	GA3@50ppm	9.33	3.46	49.24	36.28
T ₂	GA3@100ppm	10.26	3.66	49.77	37.61
T ₃	GA3@150ppm	10.46	3.85	52.84	38.4
T ₄	NAA@50ppm	9.06	3.41	48.67	34.87
T ₅	NAA@100ppm	9.5	3.54	49.25	37.38
T ₆	NAA@150ppm	10.4	3.67	51.82	38.21
T ₇	BA@50ppm	8.76	3.41	47.8	34.25
T ₈	BA@100ppm	8.83	3.51	49.46	36.35
T ₉	BA@150PPM	10.3	3.67	50.54	37.95
F test		S	S	S	S
SE(m)		0.25	0.05	0.67	0.54
C.D. at 5%		0.12	0.02	0.33	0.27
C.V		8.2	3.53	4.27	4.69

Table 4 Effect of PGRs on root and shoot value of different term in papaya

Treatment	Treatment	Fresh weight of shoot (g)	Dry weight of shoot (g)	Fresh weight of roots (g)	Dry weight of roots (g)	Root to shoot ratio
T ₀	Control	3.98	0.6	2.12	0.2	0.54
T ₁	GA3@50ppm	4.16	0.88	2.24	0.22	0.55
T ₂	GA3@100ppm	4.33	1.05	2.62	0.28	0.61
T ₃	GA3@150ppm	5.05	1.26	2.9	0.3	0.58
T ₄	NAA@50ppm	4.26	0.71	2.23	0.21	0.53
T ₅	NAA@100ppm	4.31	0.74	2.18	0.22	0.52
T ₆	NAA@150ppm	4.6	1.09	2.21	0.24	0.49
T ₇	BA@50ppm	4.03	0.62	2.14	0.2	0.54
T ₈	BA@100ppm	3.85	0.65	2.1	0.2	0.55
T ₉	BA@150PPM	4.12	0.63	2.12	0.2	0.52
F test		S	S	S	S	S
SE(m)		0.11	0.07	0.08	0.01	0.01
C.D. at 5%		0.05	0.04	0.04	0.01	0.01
C.V		8.12	28.84	12.18	16.55	6.26

Table 4 Economics of different bagging treatment of guava fruit.

Treatment	Cost of cultivation	Gross Return	Net Return	B:C Ratio
T ₀	13100	35000	21900	1.68
T ₁	13130	37000	23870	1.82
T ₂	13150	38500	25350	1.93
T ₃	13220	40000	26780	2.02
T ₄	13120	37000	23880	1.82
T ₅	13130	37500	24370	1.86
T ₆	13140	39000	25860	1.97
T ₇	13120	37000	23880	1.83
T ₈	13140	38000	24860	1.9
T ₉	13160	38800	25640	1.95

Conclusion:-

Growing media and growth regulators are being commonly used now-a-days to get a high germination percentage in a short period. From this paper we can conclude that by using GA3 with a combination of different types of growing media resulted in effective seed germination and good seedling vigor. The resulted high survival % of seedlings in papaya seeds attributed to GA3 which in turn catalyzes the starch conversion into simple carbohydrates by participating in the activity of alpha amylase which liberates chemical energy for the activation of the embryo. On the other hand, growing media helps the germination by providing adequate cation-exchange for holding of nutrients and also having good water holding capacity and sufficiently porous structure that permits adequate moisture and exchange of gases between the growing media and the embryo which is important for fast and invariant germination of seeds.

References:-

Ashish, D, & Patel, D. & Hemant, K. & Panigrahi, M. & Tani, M. (2022). Studies on the influence of plant growth regulators and bio-inoculants on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Red Lady under net tunnel.

Bhardwaj, R.L. (2013). Effect of growing media on seed germination and seedling growth of

papaya cv. Red Lady. Indian J. Agric. Res. 47(2): 163-168.

Choudhary RC, Kanwar J, Agarwal H, Prakash O.(2020) Effect of GA3 and growing media on seed germination of papaya (*Carica papaya* L.) cv. Pusa Nanha. IJCS. 2020;8(5):1423-1425.

Chauhan, R.& Jadhav, S.K. & Quraishi, A. (2014). An Efficient Seed Germination and Seedling Establishment Protocol for Hybrid *Carica papaya* Linn. With Application of Plant Growth Regulator. Biotechnology. 13. 139-142. 10.3923/biotech.2014.139.142.

Dash, B.P.(2019) Effect of different growing media on seed germination and subsequent seedling growth of papaya (*Carica papaya* Linn) cv. Pusa Nanha under Prayagraj agro climatic conditions. Journal of Pharmacognosy and Phytochemistry. 2019;8(5):141-143.

Desai A, Trivedi A, Panchal B, Desai V. (2017) Improvement of papaya seed germination by different growth regulator and growing media under net house condition. Int. J Curr. Microbio. App-Sci, 2017, 6(9).

Hazarika, B., Gogoi, S., Das, R. & Bordoloi, A. (2023). Studies on seed germination, growth and vigour of papaya seedlings cv. 'Sapna' as influenced by growth regulators and propagation media. The Pharma Innovation. 12. 2387-2391.

Kumararvind, R.; Patel, K.M. and Upadhyay, N.V. (2015). Effect of different growing media and containers on germination and establishment of seedlings of papaya (*Carica papaya* L.) cv. Madhubindu. Tre. Biosci. 8(1): 227-230.

Kumawat, R.; Maji, S.; Govind and Meena, D.C. (2014). Studies on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Coorg Honey Dew as influenced by media and chemicals. Journal of Crop and Weed. 10(2): 281-286.

Mali, D.S.; Nikumbhe, P.H.; Kharde, R.P. and Sonavane, P.N. (2015). Effect of pre-sowing seed treatment with plant growth regulators on germination and growth of papaya (Carica papaya L.). Bioinfolet. 12 (3 B): 668-674.

Malik N, Sengar RS, Yadav MK, Singh SK, Singh G, Kumar M. (2019) Effect of Different Plant Growth Regulators on In-vitro Callus Induction in *Carica papaya* (cv. Pusa Nanha). Int. J Curr. Microbiol. App. Sci. 2019;8(7):1217-1225.

Maneesha, SR. (2019) Effect of Calcium Nitrate and Potassium Nitrate Priming on Seed Germination and Seedling Vigour of Papaya (*Carica papaya* L.). Journal of Horticultural Sciences. 2019;14(2):149-154.

Paikra P, Paikra MS, Anil Netam. (2021) Effect of gibberellic acid and growing media on seed germination, growth and vigour of papaya (*Carica papaya* L.) seedling. The Pharma Innovation Journal. 2021;10(12):2343-2347.

Purba JN, Nyoman Srilaba, Putu Sri Wahyuni, Putu Bayu Wahyuda.(2021) Response of papaya (*Carica papaya* L.) seeds growth to the gibberellin (GA3) and cow bio urine. Journal Agronomy Tanaman Tropika, 2021, 3(2).

Ramteke, V.; Paithankar, D.H.; Ningot, E.P. and Kurrey, V.K. (2015). Effect of GA3 and propagation media on germination, growth and vigour of papaya cv. Coorg Honey Dew. The Bioscan. 10(3): 1011-1016.

Ramteke, V.; Paithankar, D.H.; Kamatyanatti, M.; Baghel, M.M.; Chauhan, J. and Khichi, P. (2015). Seed germination and seedling growth of papaya as influenced by GA3 and potting media. J. of Pro. Agri. 6: 129-133.