

Effect of different plant growth regulators on seed germination, seedling growth and establishment of papaya (*Carica papaya* L.) c.v. Taiwan-7

ABSTRACT:-

An experiment was conducted at the Department of Horticulture, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (U.P)-211007, India during year 2023-24. The experiment was laid out in Randomized Design (RBD). There were Ten treatments including control replicated three times in which growing media such as pond soil, vermicompost, and Plant growth regulators mixed in various percentages. The different levels of plant growth regulators Such as GA3, NAA and BA with different compositions in different combinations showed positive responses on the physical quality attributes of the papaya plant. The maximum values of physical quality characteristics of papaya plant like Survival percentage (90.68), number of germination days (10.01), germination percentage (90.01), number of leaves (10.46), plant height (12.76 cm), leaf area (38.4 cm²) and chlorophyll content (52.84), Number of primary roots (10.46), number of secondary roots (27.68) and stem diameter (3.85 cm) were recorded in T3 (GA3 with 150 ppm) and followed by the treatment T6 (NAA with 150 ppm) while, minimum values of these characters maximum value of physical parameter was recorded in control treatment (T0). From the present investigation it is concluded that treatment (T3) Performed best in terms of growth parameters followed by (T6) and (T9). The maximum B:C ratio (2.02) was also observed in the same treatment i.e. (T3)

Keywords: Plant growth regulators, Vermicompost, Taiwan-7, GA3, NAA.

1. INTRODUCTION:-

Papaya (*Carica papaya* L.): A tropical fruit crop with rapid growth, continuous fruiting cycle, and evergreen nature. It is an exceptionally wholesome, refreshing, and delicious tropical fruit that belongs to the Caricaceae family. Native to tropical America, papaya is also known by the names papaw or pawpaw. It stands as an ideal fruit choice for cultivation in various settings, including kitchen gardens, residential backyards, fields, and even orchards as a filler plant. Its

proximity to urban areas makes it a convenient choice for cultivation, and its nutritional richness further contributes to its popularity. **(Choudhary *et al.*, 2020)**

Nutritional Value and Richness: Papaya stands out as a nutritional powerhouse, offering a plethora of health benefits. It contains approximately 2500 International Units (IU) of vitamin A and around 85 milligrams of vitamin C per 100 grams of its succulent pulp. **(Ram, 2007)**. These vitamins are essential for maintaining healthy skin, boosting the immune system, and supporting overall well-being. In addition, papaya is notably rich in calcium and a range of other essential minerals that contribute to bone health, nerve function, and more. **Medicinal Significance:** Beyond its delectable taste and nutritional bounty, papaya holds high medicinal value. The extraction of papain from the dried latex of immature papaya fruits serves a variety of applications. Papain, a proteolytic enzyme, is used to tenderize meat, manufacture chewing gum, create cosmetics, degum natural silk, and enhance the shrink resistance of wool. The diverse utility of papain highlights its significance in industries beyond the culinary realm. **Anticancer Potential:** Recent research has uncovered another facet of papaya's potential health benefits. Papaya juice has demonstrated an *in vitro* antiproliferative effect on liver cancer cells [15-17]. This effect is believed to be attributed to compounds like lycopene present in papaya. The promising findings suggest that papaya may have a role to play in promoting health and potentially combating certain diseases. **(Chauhan *et al.*, 2014)**

Soil Requirements and Climatic Considerations: The optimal Soil Conditions: Papaya thrives in well-drained and fertile soils, which provide the ideal foundation for its growth and development. Well-drained soils prevent waterlogging, which can be detrimental to the root system. Fertile soils offer the necessary nutrients for papaya plants to flourish. **(Bhardwaj 2013)**

The soil structure should allow for efficient root penetration, promoting healthy root growth and nutrient absorption. **Climatic Preferences and Protection:** Papaya plants have specific climatic preferences that contribute to their successful cultivation. They thrive in sheltered environments that offer protection against adverse wind conditions. While papaya requires abundant sunlight for growth, exposure to strong winds can damage the plants and hinder their development. By choosing a sheltered location with ample sunlight, cultivators can provide the optimal climatic conditions for papaya to thrive. **Seedling Raising Time and Challenges:** Best

Raising Time: Optimal seedling raising time for papaya is June to September, with an alternative window of February to May in North India due to frost(Ashishet *al.*, 2022).

Naphthalene Acetic Acid (NAA) is a synthetic plant growth regulator with significant effects on plant growth and development: Root Development: NAA promotes the formation and growth of roots. It stimulates cell elongation and division in root tissues, leading to the development of a robust root system. Lateral Root Formation: NAA is involved in the initiation and development of lateral roots, which branch from the primary root. This branching increases the root surface area, enhancing nutrient and water absorption. Fruit Growth: NAA can influence fruit development by preventing premature fruit drop and enhancing fruit growth. Its application can lead to increased fruit size and yield in specific crops

Gibberellic Acid (GA) is a plant growth regulator with diverse roles in influencing plant development: Stem Elongation: GA stimulates cell division and elongation in the internodes, leading to increased stem length and overall plant height. Seed Germination: GA breaks seed dormancy and promotes germination by activating enzymes that degrade stored nutrients, facilitating embryo growth. Flowering Induction: GA interacts with other hormones and environmental cues to influence flowering. It's widely used in horticulture to trigger flowering in certain plants.Vorugantiet *al.*, 2022.

The Taiwan-7 papaya variety, scientifically known as *Carica papaya* L., is characterized by its distinct features: Fruit Color: The standout feature of this variety is its vibrant red-orange flesh, setting it apart from traditional yellow or orange varieties. Flavor and Taste: "Taiwan-7" papaya is known for its sweet and tropical flavor, making it a favorite for consumption. Fruit Size: The fruits of this variety are medium to large, making them suitable for both individual consumption and commercial use. Yield Potential: This variety boasts high yield potential, making it an attractive option for farmers seeking productive cultivars.

2. MATERIALS AND METHODS:-

2.1 Geographical location of the experimental site

The experimental site is located at a latitude of 25.41° North and longitude of 81.84° East, with an altitude of 98 meters above the mean sea level (MSL) of Prayagraj.

2.2 Climatic conditions of the experimental area

The area of Prayagraj comes under a humid sub-tropical climate, which experiences warm humid monsoon, hot dry summer and cold dry winter. The annual mean temperature is 26.1°C while monthly mean temperatures are 18-29°C. The daily average maximum temperature is about 22°C and the minimum temperature is 9°C. The average annual rainfall received is 1042.2 mm. At this location, the temperature reaches up to 46°C-48°C and the minimum temperature recorded is 4°C-5°C. The relative humidity ranges in this location between 20-94%.

Table 1 Treatment Details

| TREATMENT | Treatments |
|----------------|---------------------|
| T ₀ | Control (Unsprayed) |
| T ₁ | GA3@50ppm |
| T ₂ | GA3@100ppm |
| T ₃ | GA3@150ppm |
| T ₄ | NAA@50ppm |
| T ₅ | NAA@100ppm |
| T ₆ | NAA@150ppm |
| T ₇ | BA@50ppm |
| T ₈ | BA@100ppm |
| T ₉ | BA@150ppm |

3. RESULTS AND DISCUSSION

a) Number of days to germination

The minimum for Number of days required for initiation of germination was recorded in the treatment T3 (GA3 @ 150 ppm) at 10.01 followed by 10.68 with T6 (NAA @ 150 ppm) and the maximum days for initiation of Germination at 12.68 was recorded in the T0 control.

“Maximum number of days to germination was recorded when seeds soaked in GA3 might be due to the fact that GA3 involved in the activation of cytological enzymes which stimulates α -amylase enzyme that converts insoluble starch into soluble sugars and it also initiates the radical growth by removing some metabolic blocks” (Nagaret *al.* 2016).

b) Number of leaves per plant

At 45 DAT the maximum Number of leaves per plant was recorded in the treatment T4 (GA3 @ 150 ppm) at 10.46 followed by 10.40 with T6 (NAA @ 150 ppm) and the minimum Number of leaves per plant at 8.36 was recorded in the T0 control.

“Number of leaves/plant was noticed in seed treated with GA3 - 200 ppm. This may be due to GA3 are used for weakening of the seed coat so that the emergence of radical and plumule is positively influenced for root and shoot initiation. Beside this, GA3 also helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings” Mewada and Bakshi (2022)

c) Height of seedling

At 45 days the maximum height of the seedling was recorded in the treatment T4 (GA3 @ 150 ppm) at 12.76 cm followed by 12.00 cm with T6 (NAA @ 150 ppm) and minimum height of seedling at 9.96 cm was recorded in the T0 control.

“Height of seedling was noticed in seed treated with GA3 - 200 ppm. This may be due to GA3 are used for weakening of the seed coat so that the emergence of radical and plumule is positively influenced for root and shoot initiation. Beside this, GA3 also helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings” (Meenaet *al.*, 2017).

d) Stem Diameter

The maximum stem Diameter was recorded in the treatment T3 (GA3@ 150 ppm) at 3.85 mm followed by 3.67 mm with T6 (NAA @ 150 ppm) and the minimum stem Diameter at 3.37 mm was recorded in T0 control.

“Stem diameter was noticed in seed treated with GA3 - 200 ppm. This may be due to GA3 are used for weakening of the seed coat so that the emergence of radical and plumule is positively influenced for root and shoot initiation. Beside this, GA3 also helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings” (Deyaswari *et al.*, 2018).

e) Fresh weight of shoot (g)

At 45 DAT the maximum Fresh weight of shoot (g) was recorded in the treatment T4 (GA3@ 150 ppm) at 5.05 g followed by 4.60 g with T6 (NAA @ 150 ppm) and the minimum Fresh weight of shoot (g) at 3.98 was recorded in T0 control.

“Fresh weight of shoot (g) was noticed in seed treated with GA3 - 200 ppm. This may be due to GA3 are used for weakening of the seed coat so that the emergence of radical and plumule is positively influenced for root and shoot initiation. Beside this, GA3 also helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings” (Das, 2019).

f) Fresh weight of root (g)

Fresh weight of root (g) of four tagged seedlings was measured by metric scale at 45 days and average value was Calculated. The maximum Fresh weight of root (g) was recorded in the treatment T4 (GA3 @ 150 ppm) at 1.26 g followed by 1.09 g with T6 (NAA @ 150 ppm) and minimum days for Fresh weight of shoot (g) at 0.60 was recorded in T0 control.

“Fresh weight of root (g) was noticed in seed treated with GA3 - 200 ppm. This may be due to GA3 are used for weakening of the seed coat so that the emergence of radical and plumule is positively influenced for root and shoot initiation. Beside this, GA3 also helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings” (Maneesha, 2019).

g) Leaf area

The maximum Leaf Area was recorded in the treatment T4 (GA3 @ 150 ppm) at 38.40 cm² / plant followed by 38.21 cm² / plant with T8 (NAA @ 150 ppm) and the minimum Leaf Area at 33.69 cm² / plant was recorded in T0 control.

“Leaf area was noticed in seed treated with GA3 - 200 ppm. This may be due to GA3 are used for weakening of the seed coat so that the emergence of radical and plumule is positively influenced for root and shoot initiation. Beside this, GA3 also helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings” (Maliket *et al.*, 2019).

h) Survival percentage of papaya seedling

The maximum Survival percentage of papaya seedling was recorded in the treatment T4 (GA3 @ 150 ppm) at 90.86 % followed by 87.98% with T6 (NAA @ 150 ppm) and the minimum Survival percentage of papaya seedling at 77.79% was recorded in T0 control.

“Maximum survival was noticed in seed treated with GA3 - 200 ppm. This may be due to GA3 are used for weakening of the seed coat so that the emergence of radical and plumule is positively influenced for root and shoot initiation. Beside this, GA3 also helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings” (Ramteke, 2015).

i) Germination Percentage

The germination percentage was worked out after the final germination, i.e., after stoppage of germination. It was calculated by dividing to total number of seeds sown with the number of seeds germination and germination percent was calculated

The maximum Germination Percentage was recorded in the treatment T3 (GA3 @ 150 ppm) at 90.01 % and the minimum Germination Percentage at 60.01% was recorded in T0 control.

“Maximum germination percentage was recorded when seeds soaked in GA3 might be due to the fact that GA3 involved in the activation of cytological enzymes which stimulates α -amylase enzyme that converts insoluble starch into soluble sugars and it also initiates the radical growth by removing some metabolic blocks” (Babu *et al.* 2010).

j) Chlorophyll content

Chlorophyll content of four tagged seedlings was measured with the help of spectrophotometric ally. An average value was calculated. The maximum Chlorophyll Content

was recorded in the treatment T4 (GA3 @ 150 ppm) at 52.84 $\mu\text{m/g}$ followed by 51.82 $\mu\text{m/g}$ with T8 (NAA @ 150 ppm) and the minimum Chlorophyll Content at 45.17 $\mu\text{m/g}$ was recorded in T0 control.

k) Shoot:ratio

A perusal of data given in Table 4 revealed that use of different plant growth regulators brought out perceptible variation in shoot:root ratio of papaya seedlings at 45 days after sowing. Maximum and significantly higher shoot: root ratio (0.61) was observed in treatment T₂ and it was statistically at par with treatment T₄, T₅, T₈ and T₉ during the course of experimentation. However, minimum shoot:root ratio (0.54) was found in treatment T₆.

“Fresh weight and dry weight of roots recorded significantly increase with treatment GA3 - 200 ppm. This seems to be the effect of mobilization of water and nutrients transported at higher rate which might have promoted more production of photosynthetic product and translocated them to various plant parts which might have resulted in better growth of the seedlings and hence more fresh and dry weight” as suggested by **Desai et al., (2017)**.

Economics of papaya

It is evident from the data (Table 5) that plant growth regulators GA 3@ 150 ppm had maximum and significantly higher net returns in papaya seedlings (26780 per 1000 seedlings) over rest of the treatments except T₈, T₅, T₆ and T₂ which were found at par. Further, the minimum net return (21900 per 1000 seedlings) was recorded under the treatment T₀.

Table-2 Effect of PGRs on germination percentage, days to germination, plant height (cm) and survival percentage in papaya

| Treatment (No.) | Treatment | Germination Percentage | Number of Days of Germination | Survival Percentage (%) | Plant Height (cm) |
|-----------------|------------|------------------------|-------------------------------|-------------------------|-------------------|
| T ₀ | Control | 60.01 | 12.68 | 77.79 | 9.96 |
| T ₁ | GA3@50ppm | 68.34 | 11.34 | 84.34 | 10.23 |
| T ₂ | GA3@100ppm | 73.34 | 11.01 | 87.01 | 10.46 |
| T ₃ | GA3@150ppm | 90.01 | 10.01 | 90.68 | 12.76 |
| T ₄ | NAA@50ppm | 66.68 | 12.34 | 82.34 | 10.03 |
| T ₅ | NAA@100ppm | 70.01 | 11.01 | 85.72 | 10.3 |
| T ₆ | NAA@150ppm | 86.68 | 10.68 | 87.98 | 12 |

| | | | | | |
|----------------|-----------|-------|-------|-------|-------|
| T ₇ | BA@50ppm | 61.13 | 12.68 | 81.34 | 10 |
| T ₈ | BA@100ppm | 68.34 | 11.34 | 85.01 | 10.06 |
| T ₉ | BA@150PPM | 75.01 | 10.70 | 87.68 | 11.16 |
| F test | | S | S | S | S |
| SE(m) | | 3.11 | 0.29 | 1.18 | 0.31 |
| C.D. at 5% | | 1.56 | 0.14 | 0.59 | 0.15 |
| C.V | | 13.67 | 8 | 4.4 | 9.08 |

Table 3 Effect of PGRs on Stemdiameter (mm), Leaf area and number of leaves in papaya.

| Treatment (No.) | Treatment | Number of leaves | Stem Diameter (mm) | Chlorophyll (SPAD Value) | Leaf Area(cm ²) |
|-----------------|------------|------------------|--------------------|--------------------------|-----------------------------|
| T ₀ | Control | 8.36 | 3.37 | 45.17 | 33.69 |
| T ₁ | GA3@50ppm | 9.33 | 3.46 | 49.24 | 36.28 |
| T ₂ | GA3@100ppm | 10.26 | 3.66 | 49.77 | 37.61 |
| T ₃ | GA3@150ppm | 10.46 | 3.85 | 52.84 | 38.4 |
| T ₄ | NAA@50ppm | 9.06 | 3.41 | 48.67 | 34.87 |
| T ₅ | NAA@100ppm | 9.5 | 3.54 | 49.25 | 37.38 |
| T ₆ | NAA@150ppm | 10.4 | 3.67 | 51.82 | 38.21 |
| T ₇ | BA@50ppm | 8.76 | 3.41 | 47.8 | 34.25 |
| T ₈ | BA@100ppm | 8.83 | 3.51 | 49.46 | 36.35 |
| T ₉ | BA@150PPM | 10.3 | 3.65 | 50.54 | 37.95 |
| F test | | S | S | S | S |
| SE(m) | | 0.25 | 0.05 | 0.67 | 0.54 |
| C.D. at 5% | | 0.12 | 0.02 | 0.33 | 0.27 |
| C.V | | 8.2 | 3.53 | 4.27 | 4.69 |

Table 4 Effect of PGRs on root and shoot value of different term in papaya

| Treatment | Treatment | Fresh weight of shoot (g) | Fresh weight of roots (g) | Root to shoot ratio |
|----------------|------------|---------------------------|---------------------------|---------------------|
| T ₀ | Control | 3.98 | 2.12 | 0.54 |
| T ₁ | GA3@50ppm | 4.16 | 2.24 | 0.55 |
| T ₂ | GA3@100ppm | 4.33 | 2.62 | 0.61 |
| T ₃ | GA3@150ppm | 5.05 | 2.9 | 0.58 |
| T ₄ | NAA@50ppm | 4.26 | 2.23 | 0.53 |
| T ₅ | NAA@100ppm | 4.31 | 2.18 | 0.52 |

| | | | | |
|----------------|------------|------|-------|------|
| T ₆ | NAA@150ppm | 4.6 | 2.21 | 0.49 |
| T ₇ | BA@50ppm | 4.03 | 2.14 | 0.54 |
| T ₈ | BA@100ppm | 3.85 | 2.1 | 0.55 |
| T ₉ | BA@150PPM | 4.12 | 2.12 | 0.52 |
| F test | | S | S | S |
| SE(m) | | 0.11 | 0.08 | 0.01 |
| C.D. at 5% | | 0.05 | 0.04 | 0.01 |
| C.V | | 8.12 | 12.18 | 6.26 |

Table 5 Economics of different treatment.

| Treatment | Cost of cultivation | Gross Return | Net Return | B:C Ratio |
|----------------|---------------------|--------------|------------|-----------|
| T ₀ | 13100 | 35000 | 21900 | 1.68 |
| T ₁ | 13130 | 37000 | 23870 | 1.82 |
| T ₂ | 13150 | 38500 | 25350 | 1.93 |
| T ₃ | 13220 | 40000 | 26780 | 2.02 |
| T ₄ | 13120 | 37000 | 23880 | 1.82 |
| T ₅ | 13130 | 37500 | 24370 | 1.86 |
| T ₆ | 13140 | 39000 | 25860 | 1.97 |
| T ₇ | 13120 | 37000 | 23880 | 1.83 |
| T ₈ | 13140 | 38000 | 24860 | 1.9 |
| T ₉ | 13160 | 38800 | 25640 | 1.95 |

Conclusion:-

Growing media and growth regulators are being commonly used nowadays to get a high germination percentage in a short period. From this paper we can conclude that using GA3 with a combination of different types of growing media resulting in effective seed germination and good seedling vigor. The resulted high survival % of seedlings in papaya seeds is attributed to GA3 which in turn catalyzes the starch conversion into simple carbohydrates by participating in the activity of alpha-amylase which liberates chemical energy for the activation of the embryo. On the other hand, growing media helps the germination by providing adequate cation-exchange for holding nutrients and also having good water holding capacity and sufficiently porous structure that permits adequate moisture and exchange of gases between the growing media and

the embryo which is important for fast and invariant germination of seeds.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

References:-

1. **Ashish, D, & Patel, D. & Hemant, K. & Panigrahi, M. & Tani, M. (2022).** Studies on the influence of plant growth regulators and bio-inoculants on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Red Lady under net tunnel.
2. **Babu, D.; Patel, K.; Singh, R.K.; Yadav, A.; De, D.S.; L.C. and Deka, B.C. (2010).** *Seed germination, seedling growth and vigour of papaya under north east Indian condition. Acta Hort. 851: 299- 306.*
3. **Bhardwaj, R.L. (2013).** *Effect of growing media on seed germination and seedling growth of papaya cv. Red Lady. Indian J. Agric. Res. 47(2): 163-168.*
4. **Chauhan, R.& Jadhav, S.K. & Quraishi, A. (2014).** An Efficient Seed Germination and Seedling Establishment Protocol for Hybrid *Carica papaya* Linn. With Application of Plant Growth Regulator. *Biotechnology. 13. 139-142. 10.3923/biotech.2014.139.142.*
 - a. **Choudhary RC, Kanwar J, Agarwal H, Prakash O.(2020)** Effect of GA3 and growing media on seed germination of papaya (*Carica papaya* L.) cv. Pusa Nanha. *IJCS. 2020;8(5):1423-1425.*
5. **Das, B.P.(2019)** Effect of different growing media on seed germination and subsequent seedling growth of papaya (*Carica papaya* Linn) cv. Pusa Nanha under Prayagraj agro climatic conditions. *Journal of Pharmacognosy and Phytochemistry. 2019;8(5):141-143*

6. **Dayeswari D, Auxilia J, Malarkodi K, Vijayakumar RM.(2018)** Effect of Chemicals and Bio-Inoculants on Seedling Growth and Vigour of TNAU Papaya CO. 8 (*Carica papaya* L.). Int. J Curr. Microbiol. App. Sci, 2018;7(3):3007-3014.
7. **Desai A, Trivedi A, Panchal B, Desai V. (2017)** Improvement of papaya seed germination by different growth regulator and growing media under net house condition. Int. J Curr. Microbio. App-Sci, 2017, 6(9).
8. **Malik N, Sengar RS, Yadav MK, Singh SK, Singh G, Kumar M. (2019)** Effect of Different Plant Growth Regulators on In-vitro Callus Induction in *Carica papaya* (cv. Pusa Nanha). Int. J Curr. Microbiol. App. Sci. 2019;8(7):1217-1225.
9. **Maneesha, SR. (2019)** Effect of Calcium Nitrate and Potassium Nitrate Priming on Seed Germination and Seedling Vigour of Papaya (*Carica papaya* L.). Journal of Horticultural Sciences. 2019;14(2):149-154.
10. **Meena AK, Garhwal OP, Mahawar AK, Singh SP.(2017)** Effect of different growing media on seedling growth parameters and economics of papaya (*Carica papaya* L) cv. Pusa delicious. Int J Curr Microbiol App Sci. 2017;6(6):2964-2972.
11. **Mewada, R. & Bakshi, M. (2022).** Effect of different chemicals on germination and seedling growth in papaya cvs. red lady and red sun. 11. 1400-1403.
12. **Nagar SK, Vihol N, Husain S, Nagar PK.(2016)** Effect of different growing media on growth of seedlings of p seedlings of papaya (*Carica papaya* L.) cv. Madhu bindu. Madhu bindu under net house conditions, 2016.
13. **Ram, M. 2007.** Papaya.Indian council of Agricultural Research, New Delhi.pp.142-143.
14. **Voruganti., S.R., Singh, J. and Jayarajan, S. (2022)** Effect of growth regulator and growing media on seed germination of papaya: A review The Pharma Innovation Journal 2022; 11(4): 1121-1124.
15. Yakubu, Mohammed Saani, Bonaventure Kissinger Maalekuu, and Paul Kweku Tandoh.

2019. "Effect of Plant Growth Regulators on Chemical Quality Characteristics of Stored Seeds of Two Cotton Varieties". Asian Journal of Advances in Agricultural Research 8 (3):1-9. <https://doi.org/10.9734/AJAAR/2018/46011>.

16. Pun , Amar Bahadur, Arjun Kumar Shrestha, Kalyani Mishra Tripathi, and Dilli Ram Baral. 2023. "Effects of Plant Growth Regulators Including Micro-Nutrients on Fruit Set and Yield of Mandarin in Dhankuta District of Nepal". Asian Journal of Agricultural and Horticultural Research 10 (3):65-73. <https://doi.org/10.9734/ajahr/2023/v10i3232>.

17. Gaspar T, Kevers C, Penel C, Greppin H, Reid DM, Thorpe TA. Plant hormones and plant growth regulators in plant tissue culture. In vitro Cellular & Developmental Biology-Plant. 1996 Oct;32:272-89.

18. Choudhary RC, Kanwar J, Chouhan GS, Sing P, Tanwar DR. Effect of ga3 and growing media on seedling growth of papaya (Carica papaya L.) cv. pusa Nanha. International Journal of Chemical Studies. 2018;6(6):1008-12.