

Evaluation of Physico-Chemical and Organoleptic Properties of Value added Lotus Stem Pickle

ABSTRACT

The present study was carried in Post-harvest Laboratory of Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh, India. The current experiment was carried out to prepare the lotus stem pickle value added with carrot and green chilli using different vinegar like apple vinegar, jamun vinegar, sugarcane vinegar with the objective to assess the physico-chemical and organoleptic properties of the pickle. The experiment was conducted in Completely Randomized Design (CRD), with nine treatments and three replications. Based on the statistical analysis, it has been observed that treatment T8 (Lotus stem + Green chili + Carrot + Sugarcane vinegar) was found best in terms of physico-chemical properties viz. pH, acidity (%), total soluble solids (°Brix), vitamin C (mg/100g), total sugar (%). Based on the statistical analysis, it has been observed that treatment T6 (Lotus stem + Green chili + Carrot + Apple vinegar) was found best in terms organoleptic properties viz. colour, taste, flavour, texture and overall acceptability.

Keywords: Lotus stem, Organoleptic properties, Pickle, Physico-chemical properties, Vinegar

Commented [1]: Your opening should be a brief introduction of your topic

Commented [2]: was observed

Commented [3]: Was observed

INTRODUCTION

Nelumbo nucifera, (2n = 16) commonly known as lotus or sacred lotus is an aquatic perennial plant belonging to family Nelumbonaceae. It is a large aquatic herb with stout creeping yellowish white colored rhizomes and is of perennial in nature (Chopra et al., 1958). Lotus

Commented [4]: This reference is so old, please update

stem or Kamal kakdi, is (most of the times wrongly called as lotus root) very popular vegetable in India and several other Asian countries including China and Japan. Lotus stem is loaded with full of nutrition, it contains abundant amount of protein, amino acids, dietary fibre, starch and Vitamin C, Vitamin B1 and Vitamin B2 and has some tremendous health benefits. Lotus stem has extraordinary medicinal value. It has high iron, calcium, dietary fibre and effective constituents like starch, protein, asparagines, pyrocatechol, gallic-catechin, neochlorogenic acid, leucocyanidin, peroxides, vitamins B and C. It is used for curing fever, diarrhea, hemorrhages, dysentery, high BP, excessive menstruation. Even after such good functional qualities it is found to be under consumed vegetable. Incorporation of this in traditional ingredients is best way to introduce them in daily diet of people. It is widely favored by Asian because of its hard and crispy texture and distinctive aroma and taste. It is often used to make dishes such as salads, pickled vegetables, stir-fried foods and confections. Because lotus root contains a high concentration of polyphenolic compounds, it possesses good antioxidant activity (Liu *et al.*, 2010). Lotus stem is beneficial in relieving constipation, promote digestive health, It has cholesterol lowering property, low glycemic index, control high blood pressure, it has antioxidant property, support nervous system, anti-viral and anti-bacterial property, promote bone health and help in weight loss (Ohkoshi *et al.*, 2007).

Commented [5]: italicize

Commented [6]: rephrase

Commented [7]: This should be linked to....curing fever, diarrhea, hemorrhages

The rhizomes are 60-140 cm long 0.5 to 2.5 cm in diameter. The rhizome bears nodes and each of it produces a leaf (Sheikh, 2014). The colour of the rhizome varies from yellowish white to yellowish brown in colour, smooth longitudinally striated with brown patches, nodes and internodes are present (Pal *et al.*, 2015). The lotus rhizome and its extracts have shown diuretic, psycho-pharmacological, anti-diabetic, anti-obesity, hypoglycemic, antipyretic and antioxidant activities (Mukherjee *et al.*, 1996).

Commented [8]: The sentence can be merged

Carrot (*Daucus carota* L.) is one of the popular root vegetables grown throughout the world and is the most important source of dietary carotenoids in Western countries including the United States of America (Block, 1994; Hashimoto and Nagayama, 2004). It is one of the important root vegetables rich in bioactive compounds like carotenoids and dietary fibers with appreciable levels of several other functional components having significant health-promoting properties. In recent years, the consumption of carrot and its products have increased steadily due to their recognition as an important source of natural antioxidants besides, anticancer activity of β -carotene being a precursor of vitamin A (Speizer *et al.*, 1999; Sharma *et al.*, 2012). It contains many important vitamins and minerals and rich in antioxidants beta-carotene

Commented [9]: Can this be updated?

which have protective properties against certain forms of cancer and cardiovascular diseases (Basu *et al.*, 2001).

Green chilli (*Capsicum frutescens* L.) is a good source of plant derived chemical compounds that are known to have disease preventing and health promoting properties. Fresh chillies are an excellent source of vitamin A, tocopherol and ascorbic acid as well as neutral and acidic phenolic compounds which are important antioxidants (Howard *et al.*, 2000; Ahmed and Shivhare, 2001; Ajaykumar *et al.*, 2012). They are also good sources of provitamin A, carotenoid, viz., β -carotene, α -carotene, β -cryptoxanthin and oxygenated carotenoids or xanthophylls which can vary in composition and concentration due to differences genetics and degree of ripening (Markus *et al.*, 1999).

Pickling is the oldest and useful method which is used for the preservation of food by anaerobic fermentation or immersion in the vinegar and resulting food is called pickle. The word pickle is derived from Dutch language “pekel” which mean brine. Pickle is fermented food which contains various vegetables (beans, carrot, mango, green tomatoes, pepper, cabbage etc.) and spices while additives are added for the taste improvement. Sweet, vinegar, salt and oil pickles are very common commercial manufacture varieties. Salt pickle contains fresh and pure materials and salt; it is preserved with 12-15%. Basically, lactic acid bacteria are used for the conversion of sugar into acid (Nurul and Asmah, 2012; Wikipedia, 2017).

Pickling process is used to preserve the vegetables for long time. Vinegar and vegetable oil is used as pickling medium in Asia (Park *et al.*, 2014). Pickles contain phytochemicals and minerals which come from ingredients (Vegetables and spices), therefor pickles are useful against different diseases such as cancer, inflammation, brain dysfunction, atherosclerosis. The bioactive compounds are phenolic acids mostly flavonoids and tannins which scavenge the free radicals which are harmful for health and cause different diseases (Pietta, 2000; Huang *et al.*, 2012).

MATERIALS AND METHODS

The experiment was conducted in the Post-harvest Technology Laboratory, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during the year 2022-2024. The experiment was laid out

Commented [10]: delete

Commented [11]: Check spelling

Commented [12]: ??

Commented [13]: spelling

in Completely Randomized Design (CRD) having 09 treatments with three replications. The treatments were T₀ (Control), T₁ (Lotus stem + Apple vinegar), T₂ (Lotus stem + Jamun vinegar), T₃ (Lotus stem + Sugarcane vinegar), T₄ (Lotus stem + Vinegar + Sugar), T₅ (Lotus stem + Green Chilli + Carrot), T₆ (Lotus stem + Green Chilli + Carrot + Apple vinegar), T₇ (Lotus stem + Green Chilli + Carrot + Jamun vinegar), T₈ (Lotus stem + Green Chilli + Carrot + Sugarcane vinegar).

Preparation and Storage of Lotus Stem Pickle

The lotus stem was washed thoroughly and peeled. Then, it was cut into slices and boiled until they became soft. The boiled pieces were left to sun dry to remove moisture. After drying, the slices were placed into a container, and oil and spices were added. The mixture was stirred well with a spoon to ensure the spices were thoroughly mixed with the slices. Vinegar and sodium benzoate were then added to the mixture. The pickle was filled into sterilized plastic bottles and stored at ambient room temperature.

Commented [14]: The sentence can be emerged

Commented [15]: removed

Evaluation of Physico-chemical Properties of Lotus Stem Pickle

The value added lotus stem pickles were evaluated for various physico-chemical properties like pH, acidity, TSS, moisture, vitamin C, total sugar. The pH content was analyzed by digital pH meter. The acidity content was analyzed by titration method. The TSS content was analyzed by hand refractometer. The moisture content was analyzed by oven dry method. The vitamin C content was analyzed by 2, 6-dichlorophenol-inndophenol visual titration method. The total sugar content was analyzed by lane and eynon method. The recorded data of all samples for different parameters were tabulated and statistically analyzed to find out the most suitable treatment combination in terms of physico-chemical properties.

Commented [16]: You can mрге

Evaluation of Organoleptic Properties of Lotus Stem Pickle

The value added lotus stem pickles were evaluated by a panel of five judges to determine colour, taste, flavour, texture and overall acceptability. Each sample was evaluated and given a score by the panelists on the scale of 1-9 using hedonic rating method. The mean scores of all samples of all the five members were tabulated and statistically analyzed to find out which treatment combination is most acceptable in terms of sensory acceptability.

RESULTS AND DISCUSSION

Physico-chemical Properties of Lotus Stem Pickle

The nutritional value of lotus stem pickle was evaluated by analyzing its physico-chemical properties like pH, acidity, TSS, moisture, vitamin C and total sugar. The data recorded on physico-chemical properties of lotus stem pickle have been presented in Table 1.

Effect of different treatments on pH of Lotus Stem Pickle

Statistical analysis revealed that pH content differed significantly across all treatments. The mean values of pH content ranged from 4.30 to 5.02%. The minimum pH content 4.30 was recorded in T₈ (Lotus stem + Green chili + Carrot + Sugarcane vinegar), while the maximum pH content 5.02 was recorded in T₁ (Lotus stem + Apple vinegar). Similar result were reported by Mondal *et al.* (2013) in jackfruit pickle; Pal *et al.* (2018) in wood apple pickle; Kokani and Mohape (2021) in carrot pickle; Sultana *et al.* (2021) in carrot, green chilli and brinjal pickle.

Commented [17]: You can show values got by Mondal....etc

Effect of different treatments on Acidity of Lotus Stem Pickle

Statistical analysis revealed that acidity content differed significantly across all treatments. The mean values of acidity content ranged from 0.78 to 1.37%. The minimum acidity content 0.78% was recorded in T₂ (Lotus stem + Jamun vinegar), while the maximum acidity content 1.37% was recorded in T₈ (Lotus stem + Green chili + Carrot + Sugarcane vinegar). Similar results were reported by Mir *et al.* (2020) in carrot pickle; Kokani and Mohape (2021) in carrot pickle; Sultana *et al.* (2021) in carrot, green chilli and brinjal pickle; Rymbai and Chaurasiya (2022) in carrot, pea and ginger pickle.

Effect of different treatments on Total Soluble Solids of Lotus Stem Pickle

Statistical analysis revealed that TSS content differed significantly across all treatments. The mean values of TSS content ranged from 17.14 to 26.17°Brix. The minimum TSS content 17.14°Brix was recorded in T₂ (Lotus stem + Jamun vinegar), while the maximum TSS content 26.17°Brix was recorded in T₄ (Lotus stem + Vinegar + Sugar). Similar results were reported by Pal *et al.* (2018) in wood apple pickle; Verma *et al.* (2023) in wood apple pickle.

Effect of different treatments on Moisture of Lotus Stem Pickle

Statistical analysis revealed that moisture content differed significantly across all treatments. The mean values of moisture content ranged from 45.47 to 49.03%. The minimum moisture content 45.47% was recorded in T₀ (Control), while the maximum moisture content 49.03% was recorded in T₈ (Lotus stem + Green chili + Carrot + Sugarcane vinegar). Similar results

were reported by Mondal *et al.* (2013) in jackfruit pickle; Ali *et al.* (2013) in brinjal pickle; Rymbai and Chaurasiya (2022) in carrot, pea and ginger pickle; Meena *et al.* (2023) in mango pickle.

Commented [18]: You can also show the results of this author

Effect of different treatments on Vitamin C of Lotus Stem Pickle

Statistical analysis revealed that vitamin C content differed significantly across all treatments. The mean values of vitamin C content ranged from 9.01 to 14.46mg/100g. The minimum vitamin C content 9.01mg/100g was recorded in T₀ (Control), while the maximum vitamin C content 14.46mg/100g was recorded in T₈ (Lotus stem + Green chili + Carrot + Sugarcane vinegar). Similar results were reported by Rymbai and Chaurasiya (2022) in carrot, pea and ginger pickle; Meena *et al.* (2023) in mango pickle.

Commented [19]: Please repeat same as above here

Effect of different treatments on Total sugar of Lotus Stem Pickle

Statistical analysis revealed that total sugar content differed significantly across all treatments. The mean values of total sugar content ranged from 5.10 to 7.48%. The minimum total sugar content 5.10% was recorded in T₀ (Control), while the maximum total sugar content 7.48% was recorded in T₄ (Lotus stem + Vinegar + Sugar). Similar results were reported by Thakur *et al.* (2017) in mango pickle; Saroj and Singh (2018) in mango pickle; Meena *et al.* (2023) in mango pickle.

Organoleptic Properties of Lotus Stem Pickle

The sensory acceptability of lotus stem pickle was evaluated by analyzing its organoleptic properties like colour, taste, flavour, texture and overall acceptability. The data recorded on organoleptic properties of lotus stem pickle have been presented in Table 2.

Effect of different treatments on Colour of Lotus Stem Pickle

Statistical analysis revealed that organoleptic score for colour differed significantly across all treatments. The mean values of organoleptic score for colour ranged from 4.7 to 8.3. The minimum organoleptic score 4.7 for colour was recorded in T₀ (Control), while the maximum organoleptic score 8.3 for colour was recorded in T₆ (Lotus stem + Green chili + Carrot + Apple vinegar).

Effect of different treatments on Taste of Lotus Stem Pickle

Statistical analysis revealed that organoleptic score for taste differed significantly across all treatments. The mean values of organoleptic score for taste ranged from 4.7 to 8.7. The minimum organoleptic score 4.7 for taste was recorded in T₀ (Control), while the maximum

organoleptic score 8.7 for taste was recorded in T₆ (Lotus stem + Green chili + Carrot + Apple vinegar).

Effect of different treatments on Flavour of Lotus Stem Pickle

Statistical analysis revealed that organoleptic score for flavour differed significantly across all treatments. The mean values of organoleptic score for flavour ranged from 4.3 to 8.7. The minimum organoleptic score 4.3 for flavour was recorded in T₀ (Control), while the maximum organoleptic score 8.7 for flavour was recorded in T₆ (Lotus stem + Green chili + Carrot + Apple vinegar).

Effect of different treatments on Texture of Lotus Stem Pickle

Statistical analysis revealed that organoleptic score for texture differed significantly across all treatments. The mean values of organoleptic score for texture ranged from 4.3 to 8.3. The minimum organoleptic score 4.3 for texture was recorded in T₀ (Control), while the maximum organoleptic score 8.3 for texture was recorded in T₆ (Lotus stem + Green chili + Carrot + Apple vinegar).

Effect of different treatments on Overall acceptability of Lotus Stem Pickle

Statistical analysis revealed that organoleptic score for overall acceptability differed significantly across all treatments. The mean values of organoleptic score for overall acceptability ranged from 5.3 to 8.7. The minimum organoleptic score for overall acceptability 5.3 was recorded in T₀ (Control), while the maximum organoleptic score for overall acceptability 8.7 was recorded in T₆ (Lotus stem + Green chili + Carrot + Apple vinegar).

Table 1: Effect of different treatments on Physico-chemical properties of Lotus Stem Pickle

Treatment	pH	Acidity (%)	TSS (°Brix)	Moisture (%)	Vitamin C (mg/100g)	Total Sugar (%)
T ₀	4.58	1.02	18.15	45.47	9.01	5.10
T ₁	5.02	0.80	17.66	45.95	9.72	5.45
T ₂	4.87	0.78	17.14	46.47	10.99	5.37
T ₃	4.96	0.87	18.27	46.53	10.09	5.22
T ₄	4.64	0.90	26.17	46.36	11.80	7.48
T ₅	4.38	1.28	19.89	47.65	12.91	6.27
T ₆	4.41	1.26	20.16	48.17	14.21	6.46
T ₇	4.52	1.30	20.34	48.27	13.67	6.22
T ₈	4.30	1.37	21.27	49.03	14.46	6.31
F-test	S	S	S	S	S	S
SE(d)	0.045	0.044	0.171	0.281	0.445	0.206
CV	1.202	5.082	1.053	0.725	4.585	4.214
CD at 5%	0.096	0.093	0.359	0.590	0.934	0.433

Table 2: Effect of different treatments on Organoleptic properties of Lotus Stem Pickle

Treatment	Colour	Taste	Flavour	Texture	Overall acceptability
T ₀	4.7	4.7	4.3	4.3	5.3
T ₁	7.0	7.3	7.3	6.7	6.7
T ₂	6.7	7.0	7.0	7.0	7.3
T ₃	6.7	7.0	7.0	6.7	6.7
T ₄	7.0	7.3	7.3	7.3	7.0
T ₅	7.7	7.7	7.7	7.0	7.3
T ₆	8.3	8.7	8.7	8.3	8.7
T ₇	8.0	8.3	8.3	7.7	8.3
T ₈	8.0	8.3	8.0	8.0	8.3
F-test	S	S	S	S	S
S.Ed	0.588	0.497	0.471	0.471	0.521
CV	10.126	8.257	7.193	8.248	8.748
CD at 5%	1.235	1.044	0.990	0.990	1.095

CONCLUSION

Based on the statistical analysis, it has been observed that treatment T8 (Lotus stem + Green chili

+ Carrot + Sugarcane vinegar) was found best in terms of physico-chemical properties viz. pH (4.30), acidity (1.37%), total soluble solids (21.27°Brix), vitamin C (14.46mg/100g), total sugar (6.31%). Based on the statistical analysis, it has been observed that treatment T6 (Lotus stem + Green chili + Carrot + Apple vinegar) was found best in terms organoleptic properties viz. colour (8.3), taste (8.7), flavour (8.7), texture (8.3) and overall acceptability (8.7).

This research sheds light on the untapped potential of lotus stem, demonstrating its versatility and value beyond traditional uses. By promoting the production and consumption of lotus stem pickle, we contribute to minimizing agricultural losses, fostering awareness about this nutritious vegetable, and enriching culinary experiences.

REFERENCES

- Ahmed, J. & Shivhare, U. S. (2001).** Effect of pretreatment on drying characteristics and color of dehydrated green chillies. *Journal of Food Science and Technology* 38, 504-506.
- Ajaykumar, M. T., Sandeep, L. J. & Madhukar, G. B. (2012).** Effect of Pretreatments on Quality Attributes of Dried Green Chilli Powder. *ISCA Research Journal of Engineering Sciences*, 1(1), 71-74.
- Ali, A., Mahomud, S., Ahmed, M., & Hasan, S. M. K. (2013).** Period of acceptability of brinjal pickle stored in glass bottles at room temperature (20-27°C). *Agricultural Science Research Journal*, 3(9), 267-272.
- Basu, H. N., Del Vecchio, A. J., Flider, F. & Orthofer, F. T. (2001).** Nutritional and potential disease prevention properties of carotenoids. *Journal of the American Oil Chemists' Society*, 78, 665–675.
- Block, G. (1994).** Nutrient source of pro-vitamin A carotenoids in American diet. *American Journal of Epidemiology*, 139, 290–293.
- Chopra, R. N., Chopra, I. C. & Handa, K. L. (1958).** Indigenous drugs of India. U.N. Dhur and sons Private Limited, Calcutta, Second Edition; 1958, 679.
- Hashimoto, T. & Nagayama, T. (2004).** Chemical compositions of ready-to eat fresh carrot. *Journal of the Food Hygienic Society of Japan*, 39, 324–328.
- Howard, L. R., Talcot, S. T., Brenes, C. H. & Villalon, B. (2000).** Changes in phytochemical

and antioxidant activity of selected pepper cultivars (*Capsicum* species) as influenced by maturity. *Journal of Agricultural and Food Chemistry* 48(5), 1713-1720.

Huang, S., Huang, M. & Feng, B. (2012). Antioxidant activity of extracts produced from pickled and dried mustard (*Brassica juncea* Coss. Var. *foliosa* Bailey). *International journal of food properties*, 15(2), 374-384.

Kokani, R. C. & Mohape, M. N. (2021). Standardization and storage study of carrot (*Daucus carota*) pickle. *International Journal of Food Science and Nutrition*, 6(2), 103-106.

Liu, J., Zhang, M. & Wang, S. (2010). Processing characteristics and flavour of full lotus root powder beverage. *Journal of the Science of Food and Agriculture*, 90(14), 2482-2489.

Markus, F., Daood, H. G., Kapitany, J. & Biacs, P. A. (1999). Change in the carotenoid and antioxidant content of spice red pepper paprika) as a function of ripening and some technological factor. *Journal of Agricultural and Food Chemistry*, 47, 100-107.

Meena, R., Patil, S., Yadav, N., Sagore, B., Prachi, P., Meena, B. & Jain, S. (2023). Studies on Preparation of Mango Pickle from Different Genotypes of Akola Maharashtra Region. *International Journal of Plant & Soil Science*, 35(23), 474-483.

Mir, S., Gull, A., Akhtar, G., Jeelani, R., Ganaie, T. A., Bakshi, R. A. & Masoodi, F. A. (2020). Effect of microwave heat and sodium benzoate on the quality of carrot pickles during storage at ambient temperature. *Annals: Food Science & Technology*, 21(3), 407-413.

Mondal, C., Remme, R. N., Mamun, A. A., Sultana, S., Ali, M. H. & Mannan, M. A. (2013). Product development from jackfruit (*Artocarpus heterophyllus*) and analysis of nutritional quality of the processed products. *Journal of Agriculture and Veterinary Science*, 4(1), 76-84.

Mukherjee, P. K., Balasubramanian, R., Saha, K., Saha, B. P. & Pal, M. (1996). A review on *Nelumbo nucifera* gaertn. *Ancient Science of life*, 15(4), 268-276.

Nurul, S. R. & Asmah, R. (2012). Evaluation of antioxidant properties in fresh and pickled papaya. *International Food research journal*, 9, 1117-1124.

Ohkoshi, E., Miyazaki, H., Shindo, K., Watanabe, H., Yoshida, A. & Yajima, H. (2007).

Constituents From The Leaves of *Nelumbo nucifera* Stimulate Lipolysis in The White Adipose Tissue of Mice. *Planta Medica*, 73, 1255–1259.

Pal, I. & Dey, P. (2015). A review on lotus (*Nelumbo nucifera*) seed. *International Journal of Science and Research*, 4(7), 1659-1665.

Pal, S. L., Singh, D. & Attri, P. K. (2018). Evaluation of physico-chemical properties of different types of pickles of wood apple (*Limonia acidissima*). *Journal of Pharmacognosy and Phytochemistry*, 7(1S), 1184-1187.

Park, B. R., Park, J. J., Hwang, I. G., Han, H. M., Shin, M., Shin, D. S. & Yoo, S. M. (2014). Quality and antioxidant activity characteristics during storage of tea leaf pickles with different vinegar contents. *Korean journal of food and cookery science*, 30(4), 402-411.

Pietta, P. G. (2000). Flavonoids as antioxidants. *Journal of natural products*, 63, 1035-1042.

Rymbai, I. M. R. & Chaurasiya, A. K. (2022). Quality Assessment of Mixed Pickle (Carrot, Pea and Ginger) Fermented by Lactic Acid Bacteria. *Asian Journal of Agricultural and Horticultural Research*, 9(4), 129-135.

Saroj, & Singh, K. K. (2018). Standardization of recipe and method for mango pickle. *Int. J. Chem. Stud*, 6(2), 2033-2037.

Sharma, K. D., Karki, S., Thakur, N. S. & Attri, S. (2012). Chemical composition, functional properties and processing of carrot-a review. *Journal of Food Science and Technology*, 49(1), 22–32.

Sheikh, S. A. (2014). Ethno-medicinal uses and pharmacological activities of lotus (*Nelumbo nucifera*). *J. Med. Plants Stud*, 2(6), 42-46.

Speizer, F. E., Colditz, G. A., Hunter, D. J., Rosner, B. & Hennekens, C. (1999). Prospective study of smoking, antioxidant intake and lung cancer in middle aged women. *Cancer Causes Control*, 10, 475–482.

Sultana, S., Iqbal, A. & Islam, M. N. (2014). Preservation of carrot, green chilli and brinjal by fermentation and pickling. *Int Food Res J*, 21(6), 2405-2412.

Thakur, N. S., Kumar, D., Thakur, A., Chauhan, M. & Kumar, P. (2017). Studies on development and storage quality of seedling mango (*Mangifera indica* L.) pickle of Himachal Pradesh. *Indian Journal of Ecology*, 44(6), 690-696.

Verma, A., Singh, D. & Bala, K. L. (2023). Standardization of Wood Apple (*Limonia acidissima*) Pickle. *International Journal of Plant & Soil Science*, 35(18), 1305-1311.

Wikipedia (2017). Pickling. Retrieved from <https://en.wikipedia.org/wiki/Pickling>.

UNDER PEER REVIEW