

Original Research Article

Influence of foliar spray of plant growth regulator and nutrients on fruit growth and yield attributing characteristics of Guava (*Psidium guajava* L.) cv. Apple Colour.

ABSTRACT:

The study was conducted on a 10-year-old guava plant grown in a (3*3) high density plantation at the Fruit Research Station Imaliya, Department of Horticulture, J.N.K.V.V., Jabalpur (M.P.) during mrig-bahar 2022-23. The field experiment was designed using FRBD (Factorial randomised block design) and included 20 treatment combinations of PGR (salicylic acid) and various nutrients with three replications. The findings showed that among all the fruit development parameters, i.e., fruit length (6.89 cm), fruit width (7.62 cm), average fruit weight (238.50 g), fruit volume (215.50 ml), and specific gravity (1.11g/cc), were achieved greatest in the treatments combination of (Salicylic acid 300 ppm + KNO₃ 0.5%) and all the yield attributing characteristics i.e., number of fruits per plant (97.83 fruit), yield / plant (22.40 kg) and yield / hectare (248.86 qt). The highest value was found in the treatment combination (Salicylic acid 200 ppm + Borax 0.5%) followed by (Salicylic acid 200 ppm + ZnSO₄ 0.5%) as compare (control).

Key words: foliar spray, morphological characteristics, yield, PGR, nutrients, FRBD

1. INTRODUCTION

Guava (*Psidium guajava* L.) is a significant tropical and subtropical fruit crop from the Myrtaceae family. It is native to tropical America, ranging from Mexico to Peru, and has steadily gained commercial relevance as a fruit crop in various nations. It was introduced to India in the early 17th century and has since become a commercial crop throughout the country. In India, guava is mainly grown in Uttar Pradesh, Madhya Pradesh, Bihar, Gujarat, Karnataka, Andhra Pradesh, and Maharashtra, with a total cultivated area of 3.08 lakh hectares, an annual production of 4582 thousand MT (NHB 2020-21), and productivity 23.7 metric tonnes per ha, whereas in Madhya Pradesh, the area, production, and productivity of guava is 41.69 thousand ha, 776.75 MT and 19.58 MT/ha, respectively (Annon, 2021).

Guava claims superiority over several other fruits because of its commercial and nutritional value. It is a rich and cheap source of vitamin C (2 to 5 times more than fresh orange juice, 260mg/100g) and pectin (a polysaccharide substance). The ripe fruit contain 12.3-26.3% dry matter, 77.9-86.9% moisture, 0.511% ash, 0.10-0.70% crude fat, 0.82- 1.45% crude protein and 2.0-7.2% crude fibre (Mitra and Bose, 2001).

In recent years Guava orchards in India showing nutrient deficiency and could be responsible for lesser yield and quality compared to international market. Nutrient play an important role in production and its deficiency leads in lowering the production, productivity, and quality of fruits. Among the trace elements, zinc and boron play significant role in flowering and fruiting process. Boron and zinc increase the fruit set reduce fruit drop and improve fruit quality in various fruit crops (El. Sherif *et al.*, 1997). Calcium compounds extend the shelf-life of fruits by maintaining firmness, minimizing rate of respiration, protein breakdown, disintegration of tissues and disease incidence (Bangerth *et al.*, 1972). Potassium also stimulates the synthesis of chlorophyll and increased photosynthetic activity resulting in increased stored food material in the tissue leading in increase the size of fruits (Jat and Kacha, 2014).

The plant growth regulators (PGR) act as messengers and needed in small quantities at low concentration. They enhance the rapid changes in physiological activity and improve crop productivity and quality. The use of plant growth regulators has resulted in some outstanding achievements in several fruit crops with respect to growth, yield, and quality (Suman, Pency, Meghawal, & Sahu, 2017). Among these plant hormones, salicylic acid (SA) shown potential effectiveness in maintaining yield and morphological growth of fruits (Rahmani *et al.*, 2017). Fruit yield per plant is also attributed to fruit retention percentage.

Thus, fruit yield per plant seems to be the combined effect on single fruit weight and number of fruits per plant (Harsha *et al.*, 2023). Zinc sulphate is also helpful to increase fruit size and yield (Chauhan *et al* (2019).

The basic concept of nutrient and plant growth regulators is the adjustment of plant nutrient supply to an optimum level for sustaining the desired crop productivity. The foliar application of nutrients and plant growth regulator plays a vital role in improving the growth, yield, quality of fruits.

2.0 MATERIALS AND METHODS

The experiment was carried out in the Fruit Research Station Imaliya, Department of Horticulture, J.N.K.V.V., Jabalpur (M.P.) on 10-year-old guava plant under (3*3) high density plantation during mrig-bahar of 2022-23. Trees were maintained under uniform cultural schedule. The experimental was laid out in FRBD (Factorial randomized block design) comprising 20 treatment combinations and were replicated thrice. There were two factors, first is plant growth regulator (Salicylic acid) containing 4 level and second is nutrient which contains 5 levels. The plants were sprayed with different concentrations of plant growth regulator (Salicylic acid 100, 200 and 300 ppm), nutrients (KNO₃ 0.5%, ZnSO₄ 0.5%, Ca(NO₃)₂ 2 % and Borax 0.5%) and control. Treatments were given thrice i.e., first, before bud initiation, second, at fruit setting stage and third after pre harvest stage. The following treatment combinations has been used, are presented in Table1.

Table 1: Various treatment combinations

S.No.	Notation	Treatment combination
1	S ₀ M ₀	Control
2	S ₀ M ₁	KNO ₃ 0.5 %
3	S ₀ M ₂	ZnSO ₄ 0.5%
4	S ₀ M ₃	Ca(NO ₃) ₂ 2%
5	S ₀ M ₄	Borax 0.5%
6	S ₁ M ₀	Salicylic acid 100 ppm
7	S ₁ M ₁	Salicylic acid 100 ppm + KNO ₃ 0.5 %
8	S ₁ M ₂	Salicylic acid 100 ppm + ZnSO ₄ 0.5%
9	S ₁ M ₃	Salicylic acid 100 ppm + Ca (NO ₃) ₂ 2%
10	S ₁ M ₄	Salicylic acid 100 ppm + Borax 0.5%
11	S ₂ M ₀	Salicylic acid 200ppm
12	S ₂ M ₁	Salicylic acid 200 ppm + KNO ₃ 0.5 %
13	S ₂ M ₂	Salicylic acid 200 ppm + ZnSO ₄ 0.5%
14	S ₂ M ₃	Salicylic acid 200 ppm + Ca(NO ₃) ₂ 2%
15	S ₂ M ₄	Salicylic acid 200 ppm + Borax 0.5%
16	S ₃ M ₀	Salicylic acid 300 ppm
17	S ₃ M ₁	Salicylic acid 300 ppm + KNO ₃ 0.5 %
18	S ₃ M ₂	Salicylic acid 300 ppm + ZnSO ₄ 0.5%
19	S ₃ M ₃	Salicylic acid 300 ppm + Ca(NO ₃) ₂ 2%
20	S ₃ M ₄	Salicylic acid 300 ppm + Borax 0.5%

2.0
Data

Collection

The study evaluated the impact of various treatments on fruit characteristics and yield parameters. Fruit length was measured in centimeters from stem to calyx end using vernier calipers, while fruit width was recorded at the center of the fruit. The average fruit weight was determined by weighing five randomly selected fruits from each tree using an electric balance and calculating the mean. Fruit volume was measured via the water displacement method using a measuring cylinder, and specific gravity was computed by dividing the average fruit weight by the average fruit volume. The number of fruits per plant was determined by counting all mature fruits harvested from each tree. Yield per plant was recorded using an electric weighing machine, summing the weights of all harvested fruits. Yield per hectare was estimated by

multiplying the average yield per plant by the total number of plants per hectare, calculated to be 1111 plants based on a planting density of 3 m x 3 m.

3.0 RESULTS AND DISCUSSION

The interaction effect of various PGR concentrations and nutrient treatments on guava characteristics, as delineated in Table 2 and figure 1, revealed significant improvements as compare to control. The application of Salicylic acid at 300 ppm + KNO₃ at 0.5% (S3 M1) resulted in the highest observed values for fruit length (6.89 cm), fruit width (7.62 cm), fruit weight (238.50 g), fruit volume (215.50 ml), and specific gravity (1.11 g/cc). Conversely, the control (S0 M0) recorded the lowest values for these parameters, with fruit length at 5.28 cm, fruit width at 6.14 cm, fruit weight at 177.90 g, fruit volume at 170.36 ml, and specific gravity at (1.04 g/cc).

The interaction effect of PGRs and nutrients, as detailed in Table 3 and figure 2, demonstrated significantly enhanced outcomes for the number of fruits per tree, yield per plant (kg), and yield per hectare (qt). The combination of Salicylic acid at 200 ppm + Borax at 0.5% (S2 M4) resulted in the highest number of fruits per tree (97.83), the highest yield per plant (22.40 kg), and the highest yield per hectare (248.86 qt). While, the control treatment (S0 M0) recorded the lowest values, with (73.84) fruits per tree, a yield per plant of (13.14 kg), and a yield per hectare of (145.99 qt).

The reason behind the increment of both morphological as well as yield attributing characteristics might be because of positive influence of salicylic acid on growth characteristics of guava are in agreement with those reported by Khodary et al. (2004), Szepesi et al. (2005), and Stevens et al. (2006) on tomato, Gunes et al. (2005) on maize, El-Tayeb et al. (2005) on barley, Amin et al. (2007) on onion, and Yildirim et al. (2008) on cucumber. The positive effect of salicylic acid could explain that SA plays an important role in the regulation of several vital processes and growth in plants (Raskin, 1992). As a natural phenolic compound in many plants, salicylic acid is an important component in the plant signal transduction pathway. It was reported that salicylic acid application promotes cell division and cell enlargement (Hayat et al., 2005) (Javaheri et al 2012).

Among all the nutrient treatments, the increment of both morphological as well as yield attributing characteristics may be linked to the role of KNO₃ in the present study helped to optimizing the plant growth attributes by expediting the transportation of photosynthates from leaves to maturing fruits. The increase in fruit weight may be due to potassium availability, which enhanced the stream of sucrose to the apoplast resulting in increased sugar transportation to the sink tissues and hence promoted the fruit growth (Taiz and Zeiger, 2004). A similar finding was also reported by Gill et al.

Pottasium application significantly induced fruit size this could be ascribed to activation of enzymes by K and its involvement in adenosine triphosphate (ATP) production which is important in regulating the rate of photosynthesis which enable the plants to have more food to be stored in the fruits (Havlin, et al., 2005). ATP is also used as the energy source for many plant activities (Van Brunt and Sultenfuss, 1998) including cell divisions. Then cell division largely determines the final number of cells in a fruit and thereafter the final fruit size (Lemaire-Chamley et al., 2005)

Movchan and Soboroikova (1972) reported that boron boosts nitrogen absorption, therefore aiding photosynthesis process, which, in turn, leading to a greater carbohydrate accumulation, thus contributing to increased fruit size and weight. The aforementioned findings have been repeatedly confirmed by the findings of Kumar et al. (2018), Gaund et al. (2022), Tiwari et al. (2014), M. Thiruppathi (2020), Vani et al. (2020), Shreekant (2017), Poojan et al. (2020), Kumar et al. (2022), Pratap et al. (2022), Lenka et al. (2019), Goyal et al. (2019), Pippal et al. (2019), Kumar et al. (2015), and Saini et al. (2021).

Table.2 Interaction effect of foliar spray of PGR and nutrients on fruit length (cm), fruit width (cm), average fruit weight (g), fruit volume (ml) and specific gravity (g/cc) of guava (*Psidium guajava L.*) cv. Apple Colour.

S.No.	Notation	Fruit length (cm)	Fruit width (cm)	Average fruit weight (g)	Fruit volume (ml)	Specific gravity g/cc
-------	----------	-------------------	------------------	--------------------------	-------------------	-----------------------

1	S ₀ M ₀	5.28	6.14	177.90	170.36	1.04
2	S ₀ M ₁	6.12	6.72	212.10	195.60	1.08
3	S ₀ M ₂	6.05	6.72	205.70	190.25	1.08
4	S ₀ M ₃	5.52	6.35	184.10	173.68	1.06
5	S ₀ M ₄	5.81	6.60	195.60	181.20	1.07
6	S ₁ M ₀	5.36	6.25	180.50	172.42	1.04
7	S ₁ M ₁	6.47	7.16	221.30	201.80	1.09
8	S ₁ M ₂	6.31	6.88	218.40	199.52	1.09
9	S ₁ M ₃	5.68	6.40	187.50	174.95	1.07
10	S ₁ M ₄	5.88	6.64	198.90	186.24	1.08
11	S ₂ M ₀	5.45	6.32	181.20	173.10	1.05
12	S ₂ M ₁	6.78	7.46	231.60	208.50	1.11
13	S ₂ M ₂	6.55	7.30	225.80	205.60	1.10
14	S ₂ M ₃	5.75	6.48	190.40	177.46	1.07
15	S ₂ M ₄	6.25	6.78	214.50	198.56	1.08
16	S ₃ M ₀	5.63	6.38	185.40	174.23	1.06
17	S ₃ M ₁	6.89	7.62	238.50	215.50	1.11
18	S ₃ M ₂	6.81	7.55	235.50	211.95	1.11
19	S ₃ M ₃	5.76	6.52	194.20	181.10	1.07
20	S ₃ M ₄	6.69	7.35	226.40	206.50	1.10
SEm±		0.068	0.081	2.666	2.190	0.001
CD at 5%		0.195	0.231	7.634	6.270	0.002

Table.3 Interaction effect of foliar spray of PGR and nutrients on Number of fruits per plant), Yield per plant (kg), and Yield per hectare (qt) of guava (*Psidium guajava L.*) cv. Apple Colour.

S.No.	Notation	Number of fruits per plant	Yield per plant (kg)	Yield per hectare (q)
1	S ₀ M ₀	73.84	13.14	145.99
2	S ₀ M ₁	78.25	15.62	173.54
3	S ₀ M ₂	84.74	17.11	190.09
4	S ₀ M ₃	76.12	14.62	162.43
5	S ₀ M ₄	87.11	17.45	193.87
6	S ₁ M ₀	74.58	14.00	155.54
7	S ₁ M ₁	80.55	16.00	177.76
8	S ₁ M ₂	91.24	18.77	208.53
9	S ₁ M ₃	77.20	15.20	168.87
10	S ₁ M ₄	92.28	19.64	218.20
11	S ₂ M ₀	75.35	14.24	158.21
12	S ₂ M ₁	88.74	18.00	199.98
13	S ₂ M ₂	96.71	21.24	235.98
14	S ₂ M ₃	83.24	16.74	185.98
15	S ₂ M ₄	97.83	22.40	248.86
16	S ₃ M ₀	76.95	15.02	166.87
17	S ₃ M ₁	90.38	18.21	202.31
18	S ₃ M ₂	92.80	20.12	223.53
19	S ₃ M ₃	82.64	16.22	180.20
20	S ₃ M ₄	94.47	20.65	229.42
SEm±		1.193	0.393	4.371
CD at 5%		3.416	1.126	12.513




Fig. 1. Interaction effect of foliar spray of PGR and nutrients on fruit length (cm), fruit width (cm), average fruit weight (g), fruit volume (ml) and specific gravity (g/cc) of guava (*Psidium guajava* L.) cv. Apple Colour.




Fig. 2. Interaction effect of foliar spray of PGR and nutrients on Number of fruits per plant, Yield per plant (kg), and Yield per hectare (q) of guava (*Psidium guajava* L.) cv. Apple Colour.

CONCLUSION

Foliar feeding of PGR and was an effective way for improvement of morphological as well as yield attributing characteristics of guava. The treatment combination S₃ M₁ (Salicylic acid 300 ppm + KNO₃ 0.5 %) followed by S₃ M₂ (Salicylic acid 300 ppm + ZnSO₄ 0.5%) was found to be effective in maximising the morphological parameters significantly. In case of yield parameters treatment combination S₂ M₄ (Salicylic acid 200 ppm + Borax 0.5%) followed by S₂ M₂ (Salicylic acid 200 ppm + ZnSO₄ 0.5%) was significantly effective as compare to control.

REFERENCES

- Amin, A. A., M. Rashad, and H.M.H. El-Abagy.. Physiological effect of indole-3- butyric acid and salicylic acid on growth, yield and chemical concentration of onion plants. *J. Appl. Sci. R.* 2007;3(11):1554-1563
- Anonymous. National Horticulture Board database 2021. Ministry of agriculture government of India
- Bangerth, F., Dilley, D. R., Dewey, D. H.. Effect of postharvest calcium treatments on internal breakdown and respiration of apple fruits. *J. Amer. Sot. Hort. Sci.* 1972;97: 679-682
- Chauhan, A. S., Kumar, K., Saini, P. K., Singh, V., & Singh, J. P.. Effect of NAA and zinc sulphate on fruiting, yield of litchi (*Litchi chinensis* Sonn.) cv. Calcuttia. 2019.
- El-Sherif AA, Saeed WT, Nauman UF . Effect of foliar application of potassium and zinc on behaviour of montakhab E.L. Kanater guava true. *Bull. Hort. Res. Ins. Gizd, Jat, G., & LAXMIDAS, K. H.* (2014). Response of guava to foliar application of urea and zinc on fruit set, yield and quality. *Journal of agrisearch.*1997-19981;(2).
- El-Tayeb, M.A.. Response of barley grains to the interactive effect of salinity and salicylic acid. *Plant Growth Regulation* 2005;45:215-224
- Gill, P.P.S., M.Y. Ganaie, W.S. Dhillon, and N.P. Singh.. Effect of foliar sprays of potassium on fruit size and quality of 'Patharnakh'pear. *Indian J. Hortic.* 2012;69:512–516
- Gunes, A., A. Ind, M. Alpaslan, M. Cicek, E. Guneri, and E. Guzelordu.. Effects of exogenously applied salicylic acid on the induction of multiple stress tolerance and mineral nutrition in maize. *Arch. Agro.Soil Sci.* 2005;51:678-695.
- Harsha, P. K. G., Raipuriya, S., Parihar, C., Dangi, J. P., Malik, V., Kantwa, R. C., & Sonaniya, P. Effect of foliar feeding of plant growth regulators and nutrients on morphological and yield attributing characteristics of guava (*Psidium guajava* L.) cv. Gwalior-27. *Metabolism.* 2023; 35, 36.
- Havlin, J.L., Beaton, J.D., Tisdale, S.L. and Nelson, W.L.. *Soil Fertility and Fertilizers: an introduction to nutrient management* (7thed.). Pearson Educational, Inc, NJ, USA. 2005
- Hayat Q, Hayat S, Irfan M. Ahmad A.. Effect of exogenous salicylic acid under changing environment: A review. *Environmental and Experimental Botany.* 2010;68: 14-25.
- Javaheri, M., Mashayekhi, K., Dadkhah, A., & Tavallaee, F. Z.. Effects of salicylic acid on yield and quality characters of tomato fruit (*Lycopersicum esculentum* Mill.). *Int. J. Agric. Crop Sci.* 2012;4(16), 1184-1187.
- Khodary, S.E.A. Effect of salicylic acid on the growth, photosynthesis and carbohydrate metabolism in salt-stressed maize plants. *J. Agric. Biol.* 2004; 6:5-8.
- Lemaire-Chamley, M., Petit, J., Garcia, V., Just, D., Baldet, P., Germain, V., Fegard, M., Mouassite, M., Cheniclet, C. and Rothan, C.. Changes in transcriptional profiles are associated with early fruit tissue specialization in tomato. *Plant Physiol.*, 2005;139(2): 750-769.
- Mitra, S.K. and Bose, T.K.. *Fruits: Tropical & Sub tropical VI.* Astral Publication Naya udyog, Calcutta, pp. 2001; 610-611..

- Prasad, B., Das, S., Chatterjee, D. and Singh, U.P.. Effect of foliar application of urea, zinc and boron on yield of guava. *Journal of Applied Biology*, 2005;15 (1): 44-47.
- Rahmani, N., Ahlawat, T., Kumar, S., & Mohammadi, N.. Improving productivity in Mango (*Mangifera indica* L.) cv. Kesar through foliar sprays of silicon and salicylic acid. *International Journal of Chemical Studies* 2017;5(6), 1440-1443.
- Raskin, I. Salicylate, a new plant hormone. *Plant physiol.* 1992;99:799-803.
- Stevens, J., T. Senaranta, and K. Sivasithamparam.. Salicylic acid induces salinity tolerance in tomato (*Lycopersicon esculentum* cv. Roma): Associated changes in gas exchanges, water relations and membrane stabilization. *Plant Growth Regul.* 2006;49:77-83.
- Suman, M., Pency, D.S., Meghawal, D.R., & Sahu, O.P..Effect of plant growth regulators on fruit crops. *Journal of Pharmacognosy and Phytochemistry.* 2017;6, 331–337.
- Szepesi, A., J. Csiszar, B. Bajkan, K. Gemes, and I. Tari.. Role of salicylic acid pretreatment on the accumulation of tomato plants to salts and osmotic stress. *Acta. Biol. Szeg.* 2005;49:123-125
- Taiz, Z., and E. Zeiger.. *Fisiologia Vegetal.* Armed, Porto Alegre. 2004;719
- Van Brunt, J.M. and Sultenfuss, J.H.. Functions of Potassium in plants. *Better crops.* 1998;82(3): 4-5.
- Yildirim, E., E. Turan, and M. Guvenc.. Effect of foliar salicylic acid application on growth, chlorophyll and mineral content of cucumber grown under salt stress. *J. Plant Nutr.* 2008;31:593-612.