

## The Study of Variability Heritability and Genetic Advance in Okra [(*Abelmoschus esculentus* L.) Moench] Crop

### ABSTRACT

The study of variability, heritability and genetic advance in okra [*Abelmoschus esculentus* (L.) Moench] crop was conducted during kharif 2019 to ziad 2020 at the research farm of Shri Durgaji, post graduate college Chaneshwar Azamgarh, U.P. with ten genotypes Azad Bhindi1, Azad Bhindi 2, Azad Bhindi 3, KS 312, VRO5, KS 442, KS 439, BO2, Arka Abhay and Prabhani Kranti along with 11 characters viz., days to flowering, height of plant, number of branches per plant/main shoot, number of first fruiting node, height of first fruiting node, number of nodes per plant, length of internodes, length of fruit, width of fruits, number of fruits per plant and fruit yield per plant. All data recorded as per scientific for good genotypes of releasing factors i.e. high yielding and good quality characters. Coefficient of variability for all the characters and the number of first fruiting node coefficients of variability was more in F<sub>1</sub> generations. F<sub>1</sub> generation showed coefficients of variability of greater magnitude for number of first fruiting node, number of nodes in F<sub>1</sub> generation. Number of branches per plant showed maximum coefficient of variabilities in F<sub>2</sub> than parents and F<sub>1</sub> generations and other characters. The High heritability estimates in narrow sense were observed for length of fruit in both the generations and height for all the characters except height of plant, width of fruit, number of fruits per plant and fruit yield per plant in F<sub>1</sub> generation only. Other characters showed moderate heritability in F<sub>2</sub> generations and height of plant in both generation, genetic advance in narrow sense higher in F<sub>2</sub> than F<sub>1</sub> generation for all the characters. The maximum genetic advance was observed for fruit yield per plant in F<sub>1</sub> and F<sub>2</sub> and moderate genetic advance was observed for height of plant in both F<sub>1</sub> and F<sub>2</sub> generation.

Commented [A1]: comma

Commented [A2]: delete

Commented [A3]: comma

Commented [A4]: comma

Commented [A5]: delete fullstop

Commented [A6]: capital 'C'

Commented [A7]: variability

Commented [A8]: add "in"

Commented [A9]: generations

Commented [A10]: than

Commented [A11]: generations.

## 1. INTROUCTION

Okra or *Bhindi* [*Abelmoschus esculentus* (L.) Moench] is an important vegetable crop of India. It belongs to the family Malvaceae and having chromosome number  $2n = 130$ . It behaves as often cross-pollinated crop although it is potential self pollinated crop, 8.75 to 9.61 percent out crossing (Purewal and Randhawa, 1947). Okra is believed to have originated in the Hindustani centre of origin. However, Ethiopia is also considered as its native place from where it disseminated into Arabia down Nile Valley and was introduced in to Europe by the Moors and further into Louisiana during the early 17<sup>th</sup> century by French colonists (Woodruff, 1927). India is also considered its native place as various ancestral wild forms are met with here (Yawalkar, 1965). According to ICMR report per day per capita requirement of vegetables in India is 300g (125g leaf vegetables, 100g root and tuber vegetables and 75g other vegetables), but availability of vegetables in India is about 175 g per day per capita. India stands on second place in vegetable production near about 96.02 million tonnes in the world after China from 6.75 million hectare of land with average productivity of 18.40 tonnes per hectare (NHB report, 2019). Tamilnadu has the highest productivity (25.3 t/ha) followed by Kerala (22.6 t/ha) and Uttar Pradesh (18.5 t/ha). Presently, India's share is 13.60 per cent of total world vegetables production, considering the present situation as well as the growing demand of vegetable consumption to fulfill the need of next coming generation of the country, there is an urgent need to increase the productivity per unit area and overall production of vegetables in the country.

## 2. MATERIAL AND METHODS

The study of variability, heritability and genetic advance in ~~okra~~ okra [*Abelmoschus esculentus* (L.) Moench] crop was conducted during kharif 2019 to ziad 2020 at the research farm of Shri Durgaji, post graduate college, Chaneshwar Azamgarh, U.P. with ten genotypes Azad Bhindi1, Azad Bhindi 2, Azad Bhindi 3, KS 312, VRO5, KS 442, KS 439, BO2, Arka Abhay and Prabhani Kranti along with 11 characters viz., days to flowering, height of plant, number of branches per plant / main shoot, number of first fruiting node, height of first fruiting node, number of nodes per plant, length of

Commented [A12]: Add "s"

Commented [A13]: Not cited in References

Commented [A14]: Not cited in References

Commented [A15]: Not cited in References

Commented [A16]: comma

Commented [A17]: Tamil Nadu

Commented [A18]: is

Commented [A19]: capital letters

Commented [A20]: comma

Commented [A21]: Delete, it is repeated

Commented [A22]: ,

Commented [A23]: ,

Commented [A24]: ,

Commented [A25]:

Commented [A26]: per

Commented [A27]: ,

Commented [A28]:

internodes, length of fruit, width of fruits, number of fruits per plant and fruit yield per plant. All the data recorded as per scientific for good genotypes of releasing factors *i.e.* high yielding and good quality characters.

Commented [A29]: Add :the"

### 3. RESULTS AND DISCUSSION

Commented [A30]: WRIGHT AS MENTIONED

#### A. Variability

Commented [A31]: Capital letters

In the present study, the magnitude of phenotypic coefficient of variability was higher than that of genotypic coefficient of variability for all the characters and the number of first fruiting node coefficients of variability was more in F<sub>1</sub> generations. F<sub>1</sub> generation showed coefficients of variability of greater magnitude for number of first fruiting node, number of nodes in F<sub>1</sub> generation. Number of branches per plant showed maximum coefficient of variabilities in F<sub>2</sub> than parents and F<sub>1</sub> generations and other characters.

Commented [A32]: variability

The difference between phenotypic and genotypic coefficients of variability for all the characters were very small indicating that all the characters were less influenced by environmental factors.

Commented [A33]: the

Heritable variation can't be estimated with the help of genetic coefficient of variation alone Singh *et al.* (1974) had also observed similar variations.

Commented [A34]: Not cited in References

#### B. HERITABILITY

Commented [A35]: capital letters

Heritability denotes the amount of variability, which is heritable, whereas the genetic advance predicts the amount of gain expected by imposing a particular intensity of selection. Therefore, deciding the breeding methodology to be followed, the information on these parameters is important. The heritability facilitates evaluation of genetic and environmental effects in

Commented [A36]: add :s"

phenotypic variation and helps in selection since response to selection for quantitative characters is directly proportionate of the function of its heritability and genetic advance.

Robinson (1949, 1966) categorized the heritability estimate as low (below 10%), medium or moderate (10 to 30%) and high (above 30%) in narrow sense. Several methods have been developed by several workers (Mather, 1949; Warner, 1952; Crumpacker and Allard, 1962; Mather and Jinks, 1971) for estimation of heritability in narrow sense.

According to present study, heritability was high for length of fruit in both F<sub>1</sub> and F<sub>2</sub> generations, high for days to flowering, number of branches per plant, number of first fruiting node, height of first fruiting node, number of nodes per plant, length of internode and length of fruit in F<sub>1</sub> generation; moderate for height of plant, width of fruit, number of fruits per plant and fruit yield per plant in both F<sub>1</sub> and F<sub>2</sub> generations and moderate for all the character except length of fruit in F<sub>2</sub> population. Heritability estimates alone could not be given the real picture of improvement which could be realized during selection. It is only reliable when accomplishment of genetic advance under selection.

### C. GENETIC ADVANCE

In the present study, the magnitude of genetic advance in narrow sense higher in F<sub>2</sub> than F<sub>1</sub> generations for all the characters. The maximum genetic advance was observed for fruit yield per plant in F<sub>1</sub> and F<sub>2</sub> and moderate genetic advance was observed for height of plant in both F<sub>1</sub> and F<sub>2</sub> generations presented in Table 1. The genetic advance was

Commented [A37]: Not cited in References

Commented [A38]: Not cited in References

Commented [A39]: ,

Commented [A40]: s

Commented [A41]: ;

Commented [A42]: s

Commented [A43]: in capitalletters

Commented [A44]: s

Commented [A45]: s

Commented [A46]: 1

found low for all the characters except height of per plant and fruit yield per plant in F<sub>1</sub> and F<sub>2</sub>, which suggested that these characters will not be amenable to selection. Singh and Katiyar (2004), Comstock and Robinson (1952), Yadav *et al.* (2002), Mahajan and Sharma (1979) and Rao *et al.* (1977) have also reported similar results in okra Singh *et al.* (1974), Rao and Satyavathi, (1977).

**Commented [A47]:** Arrange as year-wise

These findings were in agreement with those of Chacko *et al.* (1999), Yadav *et al.* (2002), Singh *et al.* (2004), Mishra *et al.* (2004), Bharagava *et al.* (2006), Yadav *et al.* (2007), Kumar *et al.* (2007). High heritability estimates in narrow sense were observed for length of fruit in both the generations and height for all the characters except height of plant, width of fruit, number of fruits per plant and fruit yield per plant in F<sub>1</sub> generation only. Other characters showed moderate heritability F<sub>2</sub> generations and height of plant in both generations.

**Commented [A48]:** Arrange as year-wise

**Commented [A49]:** Not cited in References

**Commented [A50]:** s

**Table 1. Estimates of variability, heritability and genetic advance for the different characters under study in okra  
[*Abelmoschus esculentus* (L.) Moench]**

Characters	Phenotypic coefficient of variability in per cent			Genotypic coefficient of variability in per cent			Heritability (narrow sense)		Genetic advance in percent of mean (narrow sense)	
	Parent	F <sub>1</sub>	F <sub>2</sub>	Parent	F <sub>1</sub>	F <sub>2</sub>	F <sub>1</sub>	F <sub>2</sub>	F <sub>1</sub>	F <sub>2</sub>
Days to flowering	8.56	8.69	9.21	8.29	8.48	8.92	38.02	28.88	7.25	7.58
Height of plant	8.48	10.29	12.33	8.26	9.98	12.10	22.15	14.52	21.15	26.69
Number of branches per plant	22.42	23.33	32.97	21.14	22.12	32.22	32.61	13.60	1.32	2.09
Number of first fruiting node	20.02	14.22	20.00	19.79	19.95	19.65	51.43	29.41	1.93	2.93
Height of first fruiting node	14.67	12.54	13.19	14.22	12.10	12.60	50.83	29.60	4.76	4.83
Number of nodes per plant	12.71	11.64	15.11	12.27	10.66	14.92	37.28	19.29	4.12	6.30
Length of internode	18.02	18.74	19.56	17.52	18.26	19.14	31.70	25.57	3.88	4.35
Length of fruit	16.52	12.93	13.12	15.47	11.97	13.11	42.04	40.04	2.82	3.05
Width of fruit	17.33	20.58	21.32	15.86	18.94	21.31	22.11	16.98	0.75	0.89
Number of fruits per plant	15.74	17.30	18.42	15.00	16.61	18.41	25.88	19.65	5.40	6.10
Fruit yield per plant	11.59	16.93	16.22	11.03	16.17	16.21	13.43	13.94	61.83	62.22

Commented [A51]: Indicate Unit e. g. cm

Commented [A52]: Indicate Unit

Commented [A53]: Indicate Unit

Commented [A54]: Indicate Unit

Commented [A55]: Indicate Unit

Commented [A56]: Indicate Unit

## REFERENCES

1. Mall RH, Robinson HF. Parent, offspring correlations and heritability of some characters in okra. *Scientia Horti*. 1979; 10(2):135-139.
2. Palaniveluchachamy K, Muthukrishnan CR, Irulappan I. Studies on heritability and genetic advance in *Bhindi* [*Abelmoschus esculentus* (L.) Moench]. *Madras Agric. J.* 1982; 69(9):597-599.
3. Rajani B, Manju P. Variability studies in okra [*Abelmoschus esculentus* (L.) Moench]. *South Indian Horti*. 1997; 45 (1-2):61-62.
4. Rao TS (Swamy Rao T), Sathyavathi GP. Genetic and environmental variability in okra. *Indian J. Agric. Sci.*, 1997; 47(2):80-81.
5. Rath UK, Sahu GS, Mishra RS. Studies on genetic variability in okra. *Orissa, J. Agric. Res.* 1991; 4(1):48-51.
6. Scossirole RE, Ferrania A Hanssimann. Genetic variability for quantitative characters in alfalfa. *State Cent. and Pl. Breed.* 1961; 594-605.
7. Shekhawat AKS, Yadav JR, Srivastava JP, Yadav JK. Studies of heritability and genetic advance in okra [*Abelmoschus esculentus* (L.) Moench]. *Anth. Stress on Envi. And Sust. Deve. A.N.D. College, Kanpur*, 2005; 53-56.
8. Yadav AK, Yadav JR, Kumar S. Study on variability, heritability and genetic advance in okra [*Abelmoschus esculentus* (L.) Moench]. *Nat. Symp. on Emerging Trends in Agrl. Res., Organized by Hi-Tech Hort. Society at PDFSR, Modipuram, Meerut from Sept. 2010*; 11-12(52):187.

Commented [A57]: In capital letters

Commented [A58]: All the references should be as per the journal's style. I have revised all the references as required.

Commented [A59]: Madras