

Effect of Cold plasma seed treatment on physiological and biochemical changes in aged Mustard seed variety-RH0749

Abstract

Low rate of seed germination is a major problem in modern agriculture. Cold activated air plasma has been tested as a pre treatment method to improve the rate and uniformity of the germination. In view of the wide spread cultivation of mustard in India and in other parts of the world, This study was conducted to determine alternative non destructive seed treatment method based on cold plasma chemistry would offer a more viable alternative over traditional seed coating technologies. Seed physiological characteristics were modified in aged low vigour mustard seeds by coating the surface of the seeds with cold plasma process using a rotating plasma reactor during 2023-24 at CCS HAU, College of Agriculture Bawal. The source atmospheric oxygen gas entering the plasma chamber during the reaction process determined the type of coating and coatings were typically. Among the different treatments imposed, seed treatment with 300V @ 10 Min significantly enhanced seed quality attributes viz., germination (92 %), root length (4.37 cm), shoot length (6.98 cm), mean seedling length (11.0 cm), Seedling dry weight (2.67 mg) and SVI-I (1115), SVI-II (254), TDH (0.78 A480 nm) with lower electrical conductivity (14.20 dsm^{-1}) when compared to control (84 %, 3.61cm, 5.39cm, 9.0, 1.97mg, 760, 165, 0.67 A480 nm and 17.73 respectively). The inhibitory effect noticed with higher dose of plasma @ 400v 10 min suggested the need of judicious usage of these plasma frequencies in such applications. The first ever report on the effect of new plasma state on seed germination and establishment of mustard revealed the positive influence and that could be used as seed treatment to enhance seed yield and quality.

Key words: Cold plasma, Mustard, Seed Germination seed vigour

INTRODUCTION

In the last few decades, seed companies invested a lot of efforts in the development of seed treatments, aimed for the improvement of both germination rate and uniformity. These two factors have a major impact on final yield and quality. Even though these treatments are generally effective in enhancement of germination rates and uniformity, they apparently involve the use of expensive materials, costly and bulky procedures, which are sometimes hazardous. The plasma treatment of seeds has been investigated as an alternative to traditional pre-sowing seed treatment in agriculture, such as physical scratching (scarification), heat treatment, chemical treatment with various acids, etc (Filatova *et al.*, 2009). Indeed, multiple studies investigated the possibility of controlling the germination by exposure of seeds to various types of plasma, including atmospheric and low-pressure plasma discharges (Volin *et al.*, 2000, Uta Schnabel *et al.* 2012, Ji *et al.*, 2016).

Indian mustard (*Brassica juncea* (Linn.) is one of the important oilseed crops contributing 25 per cent of the oilseed production of the country. It occupies a prominent place next to groundnut in meeting the oil requirement of about 50 per cent population. Physiological deterioration of seeds

during storage is considered to be one of the major factors preventing seeds from normal germination and vigorous growth (Justice and Bass, 1978). In recent years, low temperature plasma (LTP), also known as non-thermal plasma (NTP) or cold atmospheric plasma (CAP), has been widely applied in biology. It has broad applications in the field of biology, including seed germination, cultivation, surface sterilization, microorganism decontamination, food manufacturing and processing, wound healing, and food storage. In the field of agriculture, plasma agriculture or plasma farming involves comprehensive application of plasma to process from pre-cultivation until the product reaches the kitchen table. In plant sciences, studies on plasma treatment have been focused on exploring the possible applications, standardization of treatment, and characterization of plasma effects in terms of plant biochemistry (Staric., *et al.*, 2020). Recently, molecular mechanisms underlying the effects of plasma on seed germination and plant growth have been explored at the cellular level, including gene expression analysis, transcriptome profiling, protein expression analysis, and epigenetics (Yan D *et al.*, 2022, Ikmal Misnal M.F. *et al.*, 2021).

Cold plasma treatment is a fast, economic and pollution-free method to improve seed performance and crop yield (Zhou *et al.*, 2011). It has essential roles in a broad spectrum of developmental and physiological processes in plants, including reducing the bacterial bearing rate of seeds, changing seed coat structures, increasing the permeability of seed coats, and stimulating seed germination and seedling growth (Sera, *et al.*, 2008., Terumi nishioka *et al.* 2016)

MATERIAL AND METHODS

Seed lots of medium vigour was collected from Regional research station, Bawal, cold plasma frequencies (300v and 400v) were treated to seeds at different duration (Control, 3min, 5 min and 10 min). Seed and seedling quality parameters were analysed as per ISTA (Anon., 2015).

Germination (%)

Four replications of 100 seed from each treatment were kept for germination at $25\pm 1^{\circ}\text{C}$ for 10 days using between paper (BP) method. The germination percentage was expressed on the basis of normal seedlings as described in ISTA Rules (Anon., 2015).

Root length (cm)

From the standard germination test, ten normal seedlings were selected at random from each replication on 7th day and the length of root was measured from the collar region to the tip of the root to base of hypocotyl and the average root length was expressed in centimeter. Dhayala et al. indicated that the effects of a short low-pressure plasma treatment on safflower seed germination were much more effective than a long high-pressure plasma treatment



Fig.1 A. plasma discharge treatment chamber B. mustard seeds undergoing in plasma reactor

Shoot length (cm)

From standard germination test, ten normal seedlings were selected at random from each replication on tenth day and the length of shoot was measured from the collar region to the tip of coleoptile and the average shoot length was expressed in centimeter.

Seedling Vigour Index-I (SVI-I)

The germinated seedlings were evaluated on 5th and 7th day as first and final count, respectively. The percentage of germination was expressed based on the normal seedlings present in the test. Ten normal and healthy seedlings from each replication were selected randomly on 10th day and seedling length (shoot and root) was measured in centimeter. Then the Seedling Vigour Index-I was determined by multiplying standard germination (%) and mean seedling length (cm) and expressed in number (Abdul-Baki and Anderson, 1973).

$$\text{SVI-I} = \text{Germination (\%)} \times \text{Mean seedling length (cm)}$$

Seedling vigour index-II

The seedlings selected for calculating the seedling vigour index-I were oven dried after removing the cotyledon (remnant seed) and the mean the mean seedling dry weight of these seedlings was used for calculating the Seedling Vigour Index-II by using the formula given by Abdul Baki and Anderson (1973) as indicated below:

$$\text{Seedling vigour index II} = \text{Germination (\%)} \times \text{Mean seedling dry weight (mg)}$$

Mean Seedling dry weight (mg)

The seedlings used for measuring the seedling length after removing cotyledons (remnant seed) were dried in hot air oven at $80 \pm 1^{\circ}\text{C}$ for 24 hours and mean seedling dry weight was expressed in milligrams.

Electrical conductivity (dSm^{-1})

Three replicates of 50 seeds each were taken, pre-washed and soaked in 50 ml of distilled water for 8 hours at room temperature. The seed leachate was collected by decanting and EC was measured in digital model conductivity meter (Elicotype Cm-82) possessing electrode at cell constant of 1.1 with calibration on EC mode. The mean value was expressed as dSm^{-1} (Presley, 1958).

Dehydrogenase activity (OD value)

The dehydrogenase activity of the seeds was estimated in duplicate following the method developed by (Kittock and Law, 1968) with minor modifications. About 25 seeds from each treatment were pre conditioned for 6 hours. From that, five embryonic axes were separated and incubated in darkness with five ml of 0.1 per cent tetrazolium chloride solution in glass vials for 2 hrs at 40°C . After incubation, the tetrazolium chloride solution was decanted and the embryos were thoroughly washed with distilled water and surface dried with blotters. The formazan was eluted by soaking the stained embryo in five ml of methyl cellosolve (2 methoxy ethanol) for overnight and the optical density was measured using spectrophotometer model at 470 nm and methyl cellosolve alone was used as a blank. The dehydrogenase activity was expressed as optical density.

RESULT AND DISCUSSION

The results obtained from the laboratory experiments on the physiological and biochemical characters of aged seeds of mustard seeds.

Germination (%)

The data on germination percentage of mustard seeds treated with plasma revealed highly significant differences among the treatments, period of time and their dosage, T_3 : 300 volt with 10 min (92 %) followed by T_3 : 300 volt with 5 min (86.67 %) compared to control (84 %) (Table 1). Immediately after the treatment, it was observed that germination was not influenced much by plasma treatment. During seed germination, many biochemical pathways are activated inside seed such as hydrolysis of starch by amylolytic enzymes, elevated expression of amylolytic enzyme genes, and increase in GA_3 (gibberrellin) level (Chrispeels and Varma1967., S. Miyata *et al.*, 1981., Chandler *et al.*, 1984., Deikman *et al.*, 1985 and Muthukrishnan *et al.*, 1983)




Fig.2 A. Seed germination @ 300v in 10 min

B. Control

Numerous studies found that cold plasma significantly increased seed germination (Dhayal *et al.*, 2006. Zhou *et al.*, 2011, Yin, M. Q *et al.*, 2005., For example, Selcuk *et al.* 2010, found that a plasma treatment significantly increased tomato seed germination. Dhayala *et al.* indicated that the effects of a short low-pressure plasma treatment on safflower seed germination were much more effective than a long high-pressure plasma treatment. Appearance of CO molecules (bands of the Angstrom, the Herzberg and the third positive systems) and ionized O₂ + molecules (the second negative (2-) system) in spectra during seeds treatment confirmed that the plasma chemical etching of seed surface plays an important role in stimulation of biochemical processes that influence on seed germination (Zivkovic, *et al.*, 2004., Tong *et al.*, 2014., Yin *et al.*, 2005)

Shoot length and root length (cm)

The data on shoot length and root length of mustard seeds treated with plasma revealed highly significant differences among the treatments, period of time and their dosage, T₃: 300 volt with 10 min (7.6 cm and 5.27 cm) followed by T₃: 300 volt with 5 min (6.90 cm and 5.55cm) compared to control (5.39 cm and 3.61 cm) (Table 1). Immediately after the treatment, it was observed that germination was not influenced much by plasma treatment.

Previous studies suggested that the plasma induces water uptake of seeds, which results in the increase in seed germination and accelerated seedling growth (Ling *et al.* , 2015, Filatova *et al.* , 2011, Search., 2009) Our results are in line with findings on the influence of the cold plasma treatment on the oilseed rape (*Brassica napus L.*) seed germination under drought stress (Bitarafan *et al.*, 2012, Mostafavi., 2012 and Müller *et al.*, 2010) Changed seed surface in plasma treated seeds could probably enhance and accelerate water uptake. The seeds can also change their dormancy and germination processes (Baskin and Baskin, 1998)

Dry matter production (mg seedling⁻¹⁰)

The data on dry matter production of mustard seeds treated with plasma revealed highly significant differences among the treatments, period of time and their dosage, T₃: 300 volt with 10 min (2.93 mg) followed by T₃: 400 volt with 5 min (2.93 mg) compared to control (1.97mg). Variations among the quantities of dry matter produced by the seedlings were highly significant due to carry-over period, cold plasma seed treatment and its dosages as well as time period (Table 1).

Vigour Index

Significant variation was observed for vigour index-I and II due to activated plasma treatment, and its frequency as well as duration of seed exposure (Table 2). Vigour index value was not influenced significantly plasma treatments immediately after treatment. However treatment T₄: 300 volt with 10 min recorded higher vigour index-I value (1115) and seedling vigour index -II (256). Followed by T₅:400 volt with 3 min (1049, 247) compare to values recorded for control was (760, 165).

4.2.6. Electrical conductivity (dSm⁻¹)

Electrical conductivity of the seed leachate was significantly influenced due to cold plasma (Table 2). However T₄: 300 volt with 10 min treatment recorded the minimum electrical conductivity (14.2 dSm⁻¹) compared to control (17.73 dSm⁻¹). Among the treatments, recorded the lowest electrical conductivity of 16.50 and 16.70 dSm⁻¹ respectively. T₇: 400 volt with 10 min treatment. Seed recorded the highest electrical conductivity of 18.37 dSm⁻¹ followed by control (17.73) Zivkovi proposed and discussed three possible plasma treatment effects on seed outside: etching, surface functionalization, and deposition of small bioactive molecules. We focused on the mechanical outside changes in the seed coat. We confirmed that the plasma treated seeds of *C. album* had a changed surface.

Dehydrogenase activity (OD value)

Cold plasma seed treatments, dosages, and period of resulted in significant difference in dehydrogenase enzyme activity (Table 2). it was observed that control seeds recorded lowest OD value(0.67) for dehydrogenase activity and T₄: 300 volt with 10 min recorded highest Od value (0.78) the values recorded by all other treatments were on par with control.

The quality and vigour of seeds are very often based on the estimation of the viability of seeds with the tetrazolium test using dehydrogenase systems (Jensen et al. 1951; Smith 1952). Generally, the higher DHO activity was determined in embryo, than in root, what might depend on the respiratory or enzymatic activity at an early stage of the germination process.

CONCLUSION

The current research introduces the development of a technology which is based on cold radiofrequency plasma treatment of seeds that can enhance both the rate and uniformity of germination of seeds. The methodology demonstrates a great impact since it proposes an inexpensive and effective solution for seeds as a pre-germination treatment, enabling their permeability increase; replacing and even totally avoiding the need in hazardous acids and/or costly scarification treatments. Despite recent investigations, the relation between the change in the wettability of seeds and the parameters of germination (time and rate) and the effects of plasma treatment on morphology, phenology and quality of plants and fruits after plasma treatment remains obscure. In laboratory germination, the treatments by different currents did present obvious advantage in comparison to the control on the first day of emergence. However, the sprouting percentage of the pretreated seeds improved by about 6 % higher than the control in the present plasma experiment

REFERENCE

Abdul-Baki, A.A and J.D. Anderson. 1973. Vigour deterioration of soybean seeds by multiple criteria. *Crop Sci.*, 13: 630-633.

- Baskin C C, Baskin J M. 1998. Seeds. Ecology, biogeography, and evolution of dormancy and germination. San Diego: Academic Press Limited
- Bitarafan, A. Hossein, and S. Rad, "Water stress effect on spring rapeseed cultivars yield and yield components in winter planting," *Int. J. Phys. Sci.*, vol. 7, no. 19, pp. 2755–2767, 2012.
- Dhayal, M., Lee, S. Y. and Park, S. U. 2006. Using low-pressure plasma for *Carthamus tinctorium* L. seed surface modification. *Vacuum* 80, 499–506
- Filatova I, Azharonok V, Kadyrov M, Beljavsky V, Gvozdo A, Shik A, 2011. The effect of plasma treatment of seeds of some grain and legumes on their sowing quality and productivity. *Rom J Phys.* 56:139–43
- Filatova, V. Azharonok, E. Gorodetskaya, L. Mel, O. Shedikova, and a Shik, 2009. "Plasma-radiowave stimulation of plant seeds germination and inactivation of pathogenic microorganisms," *Ispc*, vol. 19, no. 10, pp. 4–7.
- Ikmal Misnal M.F., Redzuan N., Firdaus Zainal M.N., Raja Ibrahim R.K., Ahmad N., Agun L. Emerging Cold Plasma Treatment on Rice Grains: A Mini Review. *Chemosphere.* 2021;274:129972.doi: 10.1016/j.chemosphere.2021.129972. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
- ISTA (International Seed Testing Association). 2015. International Rules for Seed Testing: Ph.D. Thesis edition 2015. ISTA, Bassersdorf, Switzerland.
- Jensen C.O., Sacks W. and Baldauski F.A. 1951. The reduction of triphenyltetrazolium chloride by dehydrogenases of corn embryos. *Science* 113: 65–66.
- Ji S.H., Choi K.H., Pengkit A., Im J.S., Kim J.S., Kim Y.H., Park Y., Hong E.J., Jung S.k., Choi E.H., 2016. Effects of High Voltage Nanosecond Pulsed Plasma and Micro DBD Plasma on Seed Germination, Growth Development and Physiological Activities in Spinach. *Arch. Biochem. Biophys.* ;605:117–128. doi: 10.1016/j.abb.2016.02.028. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]
- Kittock, D.L and A.G. Law. 1968. Relationship of seed vigour to respiration and tetrazolium reduction by germinating wheat seeds. *Agron. J.*, 60: 286-288.
- Ling, L. Jiangang, S. Minchong, Z. Chunlei, and D. Yuanhua, 2015. "Cold plasma treatment enhances oilseed rape seed germination under drought stress," *Sci. Rep.*, vol. 5, pp. 1–10.
- Ling, L.; Jiafeng, J.; Jiangang, L.; Minchong, S.; Xin, H.; Hanliang, S.; Yuanhua, D. Effects of cold plasma treatment on seed germination and seedling growth of soybean. *Sci. Rep.* 2014, 4, 5859. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)] [[Green Version](#)]
- Mostafavi, 2012. "Effect of Salt Stress on Germination and Early Seedling Growth Stage of Sugar Beet," *Am.-Eurasian J. Sustain.*, vol. 6, no. 2, pp. 120–125.
- Müller, D. Lüttschwager, and P. Lentzsch, 2010 "Recovery from drought stress at the shooting stage in oilseed rape (*Brassica napus*)," *J. Agron. Crop Sci.*, vol. 196, no. 2, pp. 81–89.
- Presley, J. T. 1958. Relation of protoplast permeability of cotton seed viability and predisposition of seedling diseases. *Plant Disease Report*, 42(7): 582.

- Search, C. Journals, A. Contact, M. Iopscience, P. Sci, and I. P. Address, 2009. “Does Cold Plasma Affect Breaking Dormancy and Seed Germination ? A Study on Seeds of Lamb ’s Quarters (*Chenopodium album* agg.),” vol. 750.
- Selcuk, M., Oksuz, L. and Basaran, P. 2008. Decontamination of grains and legumes infected with *Aspergillus* spp. and *Penicillium* spp. by cold plasma treatment. *Bioresource Technol.* 99, 5104–5109
- Sera, B., Stranak, V., Sery, M., Tichy, M. and Spatenka, P. 2008. Germination of *Chenopodium Album* in response to microwave plasma treatment. *Plasma Sci. Technol.* 10, 506–511
- Smith F.G. 1952. The mechanism of the tetrazolium reaction in corn embryos. *Plant Physiol.* 27: 445–456.
- Staric P., Vogel-Mikus K., Mozetic M., Junkar I. 2020. Effects of Nonthermal Plasma on Morphology, Genetics and Physiology of Seeds: A Review. *Plants*;9:1736. doi: 10.3390/plants9121736. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Terumi Nishioka, Yuichiro Takai , Tomoko Mishima, Mitsuo Kawaradani, Hideo Tanimoto , Kiyotsugu Okada, Tatsuya Misawa , And Shinichi Kusakari ., 2016. “Low-Pressure Plasma Application for the Inactivation of the Seed-borne Pathogen *Xanthomonas campestris*,” *Biocontrol Sci.*, vol. 21, no. 1, pp. 37–43.
- Tong, J. Y., He, R., Zhang, X. L., Zhan, R. T., Chen, W. W. and Yang, S. Z. 2014. Effects of atmospheric pressure air plasma pretreatment on the seed germination and early growth of *Andrographis paniculata*. *Plasma Sci. Technol.* 16, 260
- Uta Schnabel, Oliver Handorf, Kateryna Yarova, Björn Zessin, Susann Zechlin, Diana Sydow, Elke Zellmer, Jörg Stachowiak, Mathias Andrasch, Harald Below, and Jörg Ehlbeck 2012. “Decontamination of microbiologically contaminated specimen by direct and indirect plasma treatment,” *Plasma Process. Polym.*, vol. 9, no. 6, pp. 569–575.
- Volin, F. S. Denes, R. A. Young, and S. M. T. Park, 2000. “Modification of Seed Germination Performance through Cold Plasma Chemistry Technology,” *Crop Sci.*, vol. 40, no. 6, p. 1706.
- Yan D., Lin L., Zvansky M., Kohanzadeh L., Taban S., Chriqui S., Keidar M. 2022. Improving Seed Germination by Cold Atmospheric Plasma. *Plasma*;5:98–110. doi: 10.3390/plasma5010008. [CrossRef] [Google Scholar]
- Yin, M. Q., Huang, M. J., Ma, B. Z. and Ma, T. C. 2005. Stimulating effects of seed treatment by magnetized plasma on tomato growth and yield. *Plasma Sci. Technol.* 7, 3143–3147 .
- Zhou, Z. W., Huang, Y. F., Yang, S. Z. and Chen, W. 2011. Introduction of a new atmospheric pressure plasma device and application on tomato seeds. *Agri. Sci.* 2, 23–27
- Zivkovic, N. Puac, Z. Giba, D. Grubišić and Z. Lj. Petrovic, 2004. *Seed Sci. technol.*, 32, 693–701

Table: 1 Effect of cold plasma seed treatment on Seed germination, root length, shoot length, and dry weight of mustard under lab condition.

Treatments	Seed germination (%)	Root length (cm)	Shoot Length (cm)	Mean seedling length (cm)	Seedling dry weight (mg)
T1: Control	84.33	3.61	5.39	9.0	1.97
T2: 300 volt with 3 min	86.33	4.03	6.40	10.0	2.30
T3: 300 volt with 5 min	86.67	4.37	6.98	11.0	2.67
T4: 300 volt with 10 min	92.00	5.27	7.60	13.0	2.93
T5: 400 volt with 3 min	84.33	5.55	6.90	12.0	2.93
T6: 400 volt with 5 min	79.33	3.21	4.58	8.0	1.98
T7:400 volt with 10 min	73.67	2.99	4.48	7	1.90
Mean	83.80	4.14	6.04	10.19	2.38
S.Em±	1.5886	0.1314	0.3202	0.39	0.1
CD (0.05)	6.68804	0.5532	1.3484	1.62	0.6
CV (%)	3.28318	5.4878	9.1725	6.55	10.3

Table: 2 Effect of cold plasma seed treatment on Seedling Vigour index-I, Seedling vigour index-II, Electrical conductivity, and Total dehydrogenase activity o mustard under lab condition

Treatments	Seedling vigour index -I	Seedling Vigour Index –II	Electrical Conductivity (dSm⁻¹)	Total Dehydrogenase Activity (OD value)
T₁: Control	760	165	17.73	0.67
T₂: 300 volt with 3 min	900	198	15.70	0.70
T₃: 300 volt with 5min	1044	246	15.87	0.70
T₄: 300 volt with 10 min	1115	254	14.20	0.78
T₅:400 volt with 3 min	1049	247	14.53	0.73
T₆: 400 volt with 5 min	615	157	16.80	0.70
T₇: 400 volt with 10 min	551	139	18.37	0.65
Mean	862	201	16.17	0.70
S.Em±	36.197	14.87	0.272	0.021
CD (0.05)	152.38	62.63	1.148	0.090
CV (%)	7.268	12.79	2.922	5.282

UNDER PEER REVIEW