

## **Studies on Genetic Divergence in Cabbage Genotypes in the mid hills of Solan district of H.P**

### **ABSTRACT**

Cabbage (*Brassica oleracea* var. *capitata* L.) is one of the most important vegetables of the family Brassicaceae and is grown under temperate to tropical climatic conditions for its head in more than ninety countries throughout the world. The diversity in cabbage is immense due to the presence of variation in shape, size, weight and maturity of heads. The greater the variability in the available germplasm, more would be the chances for selecting superior genotypes for use in breeding programmes or direct use as cultivars. Twenty three genotypes of cabbage including check variety, Golden Acre were evaluated in a Randomized Complete Block Design (RCBD) with three replications. The results concluded that twenty-three diverse genotypes of cabbage were grouped into four clusters. The resultant four clusters showed genetic diversity and maximum number of genotypes were accommodated in cluster I (12) followed by II (7), IV (3) and III (1). The maximum inter cluster distance was recorded between cluster II and III, therefore, the hybridization between the genotypes of cluster II and III can be performed to get superior hybrids or recombinants in segregating populations for further breeding programmes. Based on the mean performance, genetic distance and clustering pattern, inter-varietal crosses can be made which may be advantageous in creating wider variability for better heterotic combinations and transgressive segregants in cabbage.

### **INTRODUCTION**

Cabbage (*Brassica oleracea* var. *capitata* L.) is one of the most important vegetables of the family Brassicaceae and is grown under temperate to tropical climatic conditions for its head in more than ninety countries throughout the world. Modern compact-head cabbage cultivars have descended from wild, non-heading cabbage types, originating in the eastern Mediterranean region and Asia Minor (Dickson and Wallace,

1986). The vegetable Brassicas are consumed for its nutritional values, minerals, carotenoids, and vitamins. Cabbage is a rich source of Vitamin A, B<sub>1</sub>, B<sub>2</sub> and C, besides being fairly rich in protein, carbohydrates and minerals, viz., phosphorus, potassium, sodium and sulphur. In addition to being nutrient-dense, cabbage provides medicinal benefits like anti-inflammatory and anticarcinogenic characteristics, since it contains chemical substances like glucosinolates, glutathione and isothiocyanates.

The diversity in cabbage is immense due to the presence of variation in shape, size, weight and maturity of heads. The greater the variability in the available germplasm, more would be the chances for selecting superior genotypes for use in breeding programmes or direct use as cultivars. Due to the high market demand for the crop, it has become necessary to develop new and promising hybrids of cabbage. Plant breeders must assess genetic diversity among germplasms while selecting possible parental lines for two reasons:

- (i) In the hybridization programme, genetically diverse parents are likely to provide a significant heterotic effect, and
- (ii) Genetically distant parents may produce a wide range of variability in the segregating generation.

#### **MATERIALS AND METHODS:**

Twenty three genotypes of cabbage including check variety, Golden Acre were evaluated in a Randomized Complete Block Design (RCBD) with three replications. The observations were recorded on plant height (cm), days to fifty per cent marketable maturity, number of non wrapper leaves, stalk length (cm), head colour, leaf margin, shape of blade, attitude of outer leaves, polar diameter of head (cm), equatorial diameter of head (cm), head shape index, head compactness ( $\text{g}/\text{cm}^3$ ), core length (cm), grosshead weight (g), net head weight (g), yield per plot (kg) & per hectare (q), TSS ( $^{\circ}\text{B}$ ), ascorbic acid content ( $\text{mg}/100\text{g}$ ) and anthocyanin content ( $\text{mg}/100\text{g}$ ). The standard growing practices of cabbage recommended in the Package of Practices for Vegetable Crops-2020 by Dr YS Parmar University of Horticulture and Forestry, Nauni, Solan (HP) were followed to raise a healthy crop. Many cabbage researchers (Singh, 2011 and Meena et al. 2013) have emphasised the use of Mahalanobis  $D^2$  statistics (Rao, 1952) for evaluating genetic divergence since it allows accurate comparison of all potential pairs of populations in any given group effecting actual crosses. List of cabbage

**Commented [RR1]:** How many sample every variables?

genotypes along with their sources have been presented in table 1 :-

**Commented [RR2]:** Please mention your software that used to process this data

**Table 1 :List of cabbage genotypes and their sources**

Sr. No.	Genotypes	Source
1.	CB – 1	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
2.	CB – 3	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
3.	CB – 4	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
4.	CB – 5	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
5.	CB – 6	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
6.	CB – 7	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
7.	CB – 8	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
8.	CB – 9	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
9.	CB – 10	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
10.	CB – 11	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
11.	CB – 12	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
12.	CB – 13	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
13.	CB – 14	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
14.	CB – 15	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
15.	CB – 16	ICAR-IARI Regional Research Station, Katrain, Kullu, HP
16.	UHF-M-1	Dept of Vegetable Science, Dr YSPUHF – Nauni, Solan, HP
17.	UHF-M-2	Dept of Vegetable Science, Dr YSPUHF – Nauni, Solan, HP
18.	UHF-M-3	Dept of Vegetable Science, Dr YSPUHF – Nauni, Solan, HP
19.	UHF-M-4	Dept of Vegetable Science, Dr YSPUHF – Nauni, Solan, HP
20.	UHF-M-7	Dept of Vegetable Science, Dr YSPUHF – Nauni, Solan, HP
21.	UHF-M-9	Dept of Vegetable Science, Dr YSPUHF – Nauni, Solan, HP
22.	UHF-2	Dept of Vegetable Science, Dr YSPUHF – Nauni, Solan, HP
23.	<b>Golden Acre (Check)</b>	ICAR-IARI Regional Research Station, Katrain, Kullu, HP

## RESULTS AND DISCUSSION:

### Composition of clusters

The grouping pattern of twenty-three cabbage genotypes has been provided in Table 2

based on the performance of various attributes. All genotypes were divided into four (I-IV) groups. Cluster I had the most genotypes (12), followed by Cluster II (7), Cluster IV (3) and Cluster III (1). The genotypes that appeared in the same cluster did so because they shared genetic homogeneity. Group constellation through genetic divergence has also been reported by Khan et al. (2009) in kale, Meena et al. (2013), Deep (2019) and Kumari (2019) in cabbage, Quamuruzzaman et al. (2007), Dey et al. (2010), Santhosha et al. (2011) and Kumar et al. (2017) in cauliflower.

**Table 2 Clustering pattern of genotypes of cabbage on the basis of genetic divergence**

Cluster	Number of genotypes	Name of genotypes
I	12	CB-1, CB-3, CB-4, CB-6, CB-7, CB-9, CB-10, CB-11, CB-12, CB-15, CB-16, UHF-2
II	7	CB-5, CB-9, CB-12, UHF-M-1, UHF-M-3, UHF-M-4, Golden Acre
III	1	CB-8
IV	3	CB-13, CB-14, UHF-M-2

**Intra and inter-cluster genetic divergence ( $D^2$ )**

Table 3 displays the average intra and inter-cluster divergence ( $D^2$ ) values. The intra-cluster distances are depicted by the diagonal figures in the table. Cluster I had the greatest intra-cluster distance (14.16) while Cluster IV had the lowest (12.50). The high intra-cluster distance suggested that the genotypes in cluster I were genetically diverse to a large extent. The maximum inter-cluster distance (36.24) was recorded between Cluster II and III, showing significant variability between these two clusters, while the least distance (21.05) was reported between Cluster II and IV, indicating a close association.

**Table 3 Average intra and inter-cluster distance among 23 genotypes of cabbage**

**Commented [RR3]:** What is the purpose intra and inter cluster in this research?

**Commented [RR4]:** Why do you choose this value 14.16, 12.99, 12.50, and 0?

Cluster	I	II	III	IV
I	<b>14.16</b>			
II	22.97	<b>12.99</b>		
III	23.45	36.24	<b>0</b>	
V	22.07	21.05	24.01	<b>12.50</b>

#### 4.3.1 Mean performance of clusters

Furthermore, dependable conformity based on cluster means was determined for several horticultural traits and is shown in Table 4. Cluster III (17.50 cm) had the tallest plants, followed by cluster IV (16.83 cm), cluster II (16.42 cm) and cluster I (15.86 cm). Cluster II (96.67 days) had the earliest marketable maturity, followed by cluster IV (103.56 days), with a progressive delay in cluster I (104.33 days) and cluster III (115.33 days). The number of non-wrapper leaves orderly increased through cluster II (15.81), clusters I (16.75), cluster IV (18.78) and cluster III (26.00). Similarly, cluster-wise increment in stalk length was observed as; cluster III (6.50 cm), cluster I (7.12 cm), cluster IV (7.80 cm) and cluster II (8.28 cm). The highest value of polar diameter of head was recorded in cluster I (11.37 cm) followed by cluster II (11.15 cm), cluster IV (10.64 cm) and the lowest value was recorded in cluster III (9.00 cm). Highest to lowest values of equatorial diameter of head were scaled through cluster II (10.97 cm), cluster IV (10.48 cm), cluster I (10.30 cm) and cluster III (8.43 cm). Head shape index was recorded maximum in cluster I (1.11) followed by cluster III (1.07), cluster II (1.02) and cluster IV (1.02). The highest gross head weight was observed in cluster IV (1214.44 g) followed by cluster II (1191.67 g), and cluster I (1031.33 g), while lowest was observed in cluster III (723.33 g). The highest value of net head weight was depicted in Cluster IV (685.56 g) followed by Cluster II (684.29 g), cluster I (555.60 g) and cluster III (366.67). Maximum head compactness was observed in cluster IV (56.88 g/cm<sup>3</sup>) followed by cluster III (55.80 g/cm<sup>3</sup>), cluster II (51.77 g/cm<sup>3</sup>) and cluster I (43.83 g/cm<sup>3</sup>). The smallest core length was observed in cluster III (4.33 cm) followed by cluster II (5.35 cm), cluster IV (5.62 cm) and cluster I (5.74 cm). The highest total soluble solids were estimated in the order viz., cluster III (7.50 °B), cluster IV (6.50 °B), cluster I (6.34°B) and cluster II (6.26 °B).

Maximum ascorbic acid content was observed in cluster II (28.64 mg/100 g) followed by cluster IV (27.63 mg/100 g), cluster I (15.36 mg/100 g) and cluster III (12.37 mg/100 g). The maximum anthocyanin content was recorded in cluster III (8.29 mg/100 g) followed by cluster IV (7.49 mg/100 g), cluster I (5.47 mg/100 g) and cluster II (5.05 mg/100 g). Cluster IV (10.28 kg) recorded the highest yield per plot, followed by cluster II (10.26 kg), cluster I (8.33 kg) and cluster III (5.50 kg).

**Table 4 Cluster means for different characters among 23 genotypes of cabbage**

Sr. No.	Characters	Clusters			
		I	II	III	IV
1	Plant Height	15.86	16.42	17.50	16.83
2	Days to 50% marketable maturity	104.33	96.67	115.33	103.56
3	Number of non-wrapper leaves	16.75	15.81	26.00	18.78
4	Stalk length (cm)	7.12	8.28	6.50	7.80
5	Polar diameter (cm)	11.37	11.15	9.00	10.64
6	Equatorial diameter (cm)	10.30	10.97	8.43	10.48
7	Head Shape Index	1.11	1.02	1.07	1.02
8	Gross Head weight (g)	1031.33	1191.67	723.33	1214.44
9	Net Head weight (g)	555.60	684.29	366.67	685.56
10	Head Compactness (g/cm <sup>3</sup> )	43.83	51.77	55.80	56.88
11	Core Length (cm)	5.74	5.35	4.33	5.62
12	TSS (°B)	6.34	6.26	7.50	6.50
13	Ascorbic Acid (mg/100 g)	15.36	28.64	12.37	27.63
14	Anthocyanin (mg/100 g)	5.47	5.05	8.29	7.49
15	Yield per plot (kg)	8.33	10.26	5.50	10.28

## CONCLUSION

In the present study, based on genetic divergence, twenty-three diverse genotypes of cabbage were grouped into four clusters. The resultant four clusters showed genetic diversity and maximum number of genotypes were accommodated in cluster I (12)

followed by II (7), IV (3) and III (1). The maximum inter cluster distance was recorded between cluster II and III, therefore, the hybridization between the genotypes of cluster II and III can be performed to get superior hybrids or recombinants in segregating populations for further breeding programmes. Based on the mean performance, genetic distance and clustering pattern, inter-varietal crosses can be made which may be advantageous in creating wider variability for better heterotic combinations and transgressive segregants in cabbage.

#### REFERENCES :

- Dickson MH and Wallace DH. Cabbage breeding. in: Bassett MJ ed. Breeding vegetable crops. 1986;395-432.
- Dey SS, Sharma SR, Prakash C, Barwal RN and Bhatia R. Genetic divergence in snowball cauliflower (*Brassica oleracea* L. var. *botrytis* L.). *Indian Journal of Plant Genetic Resources*.2010; 24: 48-51.
- Deep N.Genetic evaluation of exotic germplasm of red cabbage (*Brassica oleracea* var. *capitata* f. *rubra*) (MSc Thesis). Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, HP, India. 2019 ;49-51.
- Kumar V, Singh DK, Panchbhaiya A and Singh N. Genetic divergence studies in mid-season cauliflower (*Brassica oleracea* var. *botrytis* L.) through Principal Component Analysis (PCA) and D<sup>2</sup> analysis. *International Journal of Chemical Studies*.2017; 5(4): 989-993.
- Kumari S. Genetic Variability and Correlation studies in Cabbage (*Brassica oleracea* var. *capitata* L.) (MSc Thesis). Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, HP, India. 2019; 57-58.
- Khan SH, Ahmed N, Jabeen N, Mushtak F and Hossain K. Genetic divergence in Kale (*Brassica oleracea* L. var. *acephala* L.). *Asian Journal of Horticulture* 2009;4: 134-137.
- Meena ML, Ram RB, Dwivedi DH and Bharti N. Estimation of biological divergence in cabbage (*Brassica oleracea* L. var. *capitata* L.). *Progressive Horticulture* 2013; 45: 351-354.

Quamruzzaman AKM, Rahman MM, Uddin MN, Siddiky MA and Prodhann MDH. Genetic diversity in cauliflower (*Brassica oleracea* L. var. *botrytis* L.). *Indian Journal of Horticulture* 2007; 64: 50-52.

Rao CR. *Advanced Statistical Method in Biometrical Research*. John Wiley and Sons Inc., New York. 1952; 357-363.

Santhosha HM, Varalakshmi B and Narase Gowda NC. Genetic diversity in early cauliflower (*Brassica oleracea* L. var. *botrytis* L.) germplasms. *Journal of Horticultural Sciences* 2011; 24: 21-24.

UNDER PEER REVIEW