

Original Research Article

Chemical Fungicides and Antagonists as Potential Treatments for *Ganoderma lucidum*-Associated Basal Stem Rot in Mandarin Trees

ABSTRACT

Ganoderma lucidum causes basal stem rot disease in mandarin trees, resulting in a variety of symptoms including dieback, foliage discoloration, an unhealthy appearance and eventual tree declines or death. Prior to field application, it is imperative to assess the efficacy of newer molecules and possible potential antagonists through *in vitro* testing. In this study, ten chemical fungicides and four antagonists were evaluated for their effectiveness against *G. lucidum* using the poison food method and dual culture technique, respectively. Among the chemical fungicides tested, Copper oxychloride 50% WP (@3g/l), Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L SC (@0.6 g/l), and Metiram 55% + Pyroclostrubin 5% WG (@1g/l) demonstrated the highest effectiveness, entirely suppressing (100%) pathogen growth. In the dual culture technique, the fungal antagonists *Trichoderma harzianum* (79.63%) and *Trichoderma asperellum* (77.57%) exhibited superior effectiveness compared to *Pseudomonas fluorescense* and *Bacillus subtilis* against *G. lucidum*. These findings highlight the efficacy of chemical fungicides and antagonists in treating *G. lucidum*-induced basal stem rot in mandarin trees.

Keywords: Antagonist, fungicides, *Ganoderma lucidum*, Mandarin

1. INTRODUCTION

The root parasite *Ganoderma lucidum* (Leys) causes basal stem rot disease, which affects a variety of hard wood-fruit, ornamental, social forest and forest trees [1]. It causes white spongy root rot, also known as trunk rot, root rot, basal rot and *Ganoderma* root rot. Affected trees show varying degrees and combinations of dieback, foliage colour loss and generally unhealthy appearance. The tree eventually declines or dies. The fungi can spread through the air via basidiospores or by root grafting from infected to healthy tree roots. When *Ganoderma* spp. infect main or lateral roots, a whitish mycelium mat forms beneath the bark, gradually transforming into a brown hue. This white mycelium progresses upward from crown roots to the trunk, forming strands or ribbons known as rhizomorphs [2]. Near or at the base of tree, these structures produce fan-shaped fruiting bodies, commonly referred to as brackets

or semi-circular conks. These brackets may be either stalked or sessile. Over time, infected roots and trunk wood become spongy, decompose and disintegrate [3].

Citrus encompasses several economically significant species, with only a few being commercially cultivated in India, including mandarins, lemons, limes, sweet oranges, and pomelos. In India, citrus cultivation spans approximately 10.91 lakh hectares, yielding production of 141.1 lakh metric tons. Maharashtra state contributes to this total with citrus cultivation covering 2.10 lakh hectares and producing 18.76 MT. Specifically, Nagpur mandarin (*Citrus reticulata*) is cultivated on 1.48 lakh ha in the Vidarbha region of Maharashtra, generating an annual yield of 13.56 lakh tons [4]. The successful cultivation of citrus faces challenges from various diseases, including fungal, viral and bacterial pathogens. Citrus is susceptible to more than 150 diseases and disorders, and the current concern among growers is the basal rot or trunk rot caused by *Ganoderma lucidum*. This particular disease, also known as Ganoderma rot, affects various crops and is caused by different Ganoderma species, posing a significant threat to plantation crop production and productivity [5]. Knorr [6] identified *Ganoderma sessilis*, now known as *G. lucidum*, as a fungal agent capable of invading healthy citrus wood and it was also isolated from trees with heart rot in Florida. Additionally, [7] reported cases in south Texas where 'Marrs' early orange trees on sour orange rootstock were infected with *G. lucidum*.

Currently, there is no effective method for managing basal stem rot in the field. Given the inadequacy of existing management systems, the exploration of novel molecules or bio-agents becomes crucial. Testing these newer biocontrol agents and molecules *in vitro* before field application is essential. Evaluating bioagents or newer fungicides molecules *in vitro*, allows for their potential inclusion as alternatives to current recommendations under natural conditions. Additionally, the utilization of both efficient bioagents and chemical solutions may contribute to minimizing the development of resistance. Furthermore, the utilization of both efficient bioagents and chemical solutions may contribute to minimizing the development of resistance.

Various researchers worldwide have explored methods to control *Ganoderma* basal rot in crops like oil palm, arecanut and coconut, but limited attention has been given to fruit crops such as mandarin. Notably, there is a research gap in the Vidarbha region regarding fungicides and bio-agents for combating mandarin basal rot. In light of these considerations, this study was undertaken to identify effective fungicides and antagonists for controlling

Ganoderma basal rot in Nagpur mandarin. In light of these considerations, this study was undertaken to identify effective fungicides and antagonists for controlling *Ganoderma* basal rot in Nagpur mandarin.

2. MATERIALS AND METHODS

The basidiocarps were obtained from infected trees of Nagpur mandarin growing in the AICRP on Fruits scheme field. The fruiting bodies were then brought to the laboratories, and subjected for isolation as method proposed by [8] with some modifications. The purified culture was then identified as *G. lucidum* based on morphological and cultural characteristics as described by [9] and [10].

2.1 Evaluation chemical fungicides

Ten chemical fungicides were evaluated by employing poisoned food method [11] against *Ganoderma* basal rot of Nagpur mandarin. Potato dextrose agar (PDA) was prepared and distributed at the rate of 100 ml in 250 ml conical flask; autoclaved at 1.5 kg/cm² for 15 minutes. Requisite quantity of each of the chemical fungicides as per concentration was added in sterilized melted PDA separately so as to obtain desired concentration. Flask containing poisoned medium was shaken well to have even and uniform distribution of fungicides. About 20 ml of poisoned PDA was poured in each pre sterilized Petri plate and allow to solidifying. These Petri plates were inoculated by respective pathogens separately. Six mm discs of ten days old fungus cultures were cut with sterilized cork borer and transferred to the centre of plates in inverted position containing poisoned medium. Suitable controls without fungicides were maintained by placing fungal discs in untreated plates. The whole procedure was carried out under aseptic conditions.

Experiment conducted in Completely Randomized Design (CRD) with four replications. The Petri plates were then incubated in an incubator at $25 \pm 2^{\circ}\text{C}$ for 7 days. The diameter of the colony of the respective fungal pathogens on medium was measured after 10 days of incubation (DAI) and per cent inhibition was calculated by using the following formula [12].

2.2 Evaluation of antagonists

The dual culture technique [13] was used to test the antagonist's activity of *Trichoderma harzianum*, *Trichoderma asperellum*, *Bacillus subtilis* and *Pseudomonas fluorescens* received from Center for Organic Research and Training, Dr. PDKV, Akola. The

pathogen (*G. lucidum*) and antagonists were grown on PDA (for fungal antagonists) and NA (for bacterial antagonists) media for a week at $25 \pm 2^{\circ}\text{C}$. Two culture disc (each 6 mm for fungal antagonists) one each of the test pathogens and test bioagents were cut out with sterilized cork borer and placed equidistance, exactly opposite to each other on autoclaved and cooled solid surface of PDA medium in Petri plates, and incubated at $25 \pm 2^{\circ}\text{C}$. To test bacterial antagonists, a test pathogen mycelium plug was placed in the centre of PDA medium on a 90 mm Petri plates. Bacterial suspension loops were streaked on inoculated plates 1 cm away from test pathogen disc. Petri plates were then incubated at $25 \pm 2^{\circ}\text{C}$. The experiment was conducted in Completely Randomized Design (CRD) with five treatments and five replications. Linear mycelial growth of test pathogen was tested after 24 hours, continued till untreated control plates were fully covered with mycelial growth of test pathogen and averaged finally. Per-cent inhibition of the test pathogen with test antagonists, over control was calculated by applying formula suggested by [14].

2.3 Statistical analysis

The data of each character were subjected to statistical test by applying of variance for different treatments. Fisher's protected critical difference (CD) test was used to indicate the difference between the treatments at the probability level of $p= 0.01$ following the procedure described by [15].

3. RESULTS AND DISCUSSION

3.1 Evaluation of fungicides

The *in vitro* fungitoxic effects of ten fungicides were assessed against *G. lucidum* at their respective concentrations. The obtained results are outlined in Table 1.

The findings indicated that ten different chemical fungicides significantly impede the radial growth of *G. lucidum*. Notably, Copper oxychloride 50% WP (@3g/l), Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L SC (@0.6 g/l), and Metiram 55% + Pyroclostrobin 5% WG (@1 g/l) emerged as the most effective, completely inhibiting the growth of the test fungus at their recommended concentrations. For Kasugamycin 5% + Copper oxychloride 45% WP (@ 2 g/l), Metalaxyl 8% + Mancozeb 64% WP (@ 2.5 g/l), Mandipropamid 23.4% SC (@ 0.8 ml/l), Propineb 70% WP (@ 3 g/l), Zineb 75% WP, (@ 2 g/l) and Cymoxanil 8% + Mancozeb 64% WP (@ 2.5 g/l), the results showed radial mycelial growth of 1.32, 1.66, 1.77, 2.64, 2.78 and 2.97 cm, respectively, accompanied by growth inhibition percentages of 86.64%,

81.47%, 80.24%, 70.63%, 69.06%, 67.36%. Fosetyl-AL 80% WP (@2.5g/l) exhibited a radial mycelial growth of 6.66 cm with least growth inhibition (26.55%). The control plate recorded the highest radial mycelial growth at 9.0 cm.

Researchers have effectively utilized chemical fungicides to control *Ganoderma*, as demonstrated by studies conducted by [16]; [17]; [18] and [19].

This finding aligns with the observations made by [20] and [1], who similarly noted the inhibitory effectiveness of copper oxychloride 50% WP against *Ganoderma lucidum* growth. Additionally, [19] documented that metalaxyl + mancozeb and carbendazim + mancozeb were also successful in impeding the growth of *G. lucidum*.

Table 1. Efficacy of chemical fungicides against *G. lucidum* (10 DAI)

SN	Treatments	Formulation	Conc. used (%)	Radial mycelium growth (cm)	% Growth inhibition
T ₁	Propineb 70%	Wettable powder	3 g/l	2.64	70.63
T ₂	Zineb 75%	Wettable powder	2 g/l	2.78	69.06
T ₃	Copper oxychloride 50%	Wettable powder	3 g/l	0.00	100.00
T ₄	Fosetyl-AL 80%	Wettable powder	2.5 g/l	6.66	26.55
T ₅	Cymoxanil 8% + Mancozeb 64%	Wettable powder	2.5 g/l	2.97	67.36
T ₆	Metalaxyl 8% + Mancozeb 64%	Wettable powder	2.5 g/l	1.66	81.47
T ₇	Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L	Suspension concentrates	0.6 g/l	0.00	100.00
T ₈	Metiram 55% + Pyroclostrobin 5%	Wettable granules	1 g/l	0.00	100.00
T ₉	Kasugamycin 5% + Copper oxychloride 45%	Wettable powder	2 g/l	1.32	86.64
T ₁₀	Mandipropamid 23.4%	Suspension concentrates	0.8 ml/l	1.77	80.24
T ₁₁	Control		-	9.00	
	SE (m) ±		-	0.57	-
	CD (P=0.01)		-	2.28	-

In the present investigation, Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L SC and Metiram 55% + Pyroclostrobin 5% WG emerged as the most potent fungicides in suppressing the mycelial growth of *G. lucidum* in laboratory conditions. A review of existing

literature suggests that these results may contribute novel information, as the effectiveness of Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L SC and Metiram 55% + Pyraclostrobin 5% WG against *G. lucidum* might be reported here for the first time. These findings could prove valuable for citrus growers grappling with Ganoderma basal rot management. However, a further investigation is needed to determine the safety and efficacy of these fungicides in field conditions.

Copper oxychloride 50% WP, Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L SC, and Metiram 55% + Pyraclostrobin 5% WG were found to be fungitoxic to *G. lucidum* based on the study's results. Copper oxychloride is broad-spectrum copper based fungicides, effective against a wide range of fungi. It is commonly used in agriculture to protect crops from fungal diseases. The fungicidal action of copper oxychloride is attributed to its ability to release copper ions, which interfere with the enzymatic processes of the fungi. This disruption inhibits fungal growth and reproduction. Pre-formulated combination fungicides, such as Fluxapyroxad + Pyraclostrobin, are commercially accessible, offering both preventive and curative effects against a wide range of fungal pathogens. Fluxapyroxad is classified as a succinate dehydrogenase inhibitor (SDHI), while Pyraclostrobin is categorized as a strobilurin fungicide. The distinct modes of action exhibited by Fluxapyroxad and Pyraclostrobin may contribute to effective control of test pathogen by disrupting energy production in the mitochondria and interferes with the electron transport chain, affecting energy metabolism in fungi. Another effective pre-mixed fungicide was Metiram 55% + Pyraclostrobin 5% WG, recognized for its effective against adiversespectrum of fungal pathogens. Metiram is an ethylene bis-dithiocarbamate (EBDC) fungicide, functions as a protective agent by inhibiting spore germination and mycelial growth on plant surfaces. Concurrently, Pyraclostrobin disrupts fungal respiration by inhibiting the mitochondrial electron transport chain, there by exerting control of fungal diseases. In this context, these specific concentrations of the mentioned chemicals demonstrated effectiveness in inhibiting the radial growth of *G. lucidum*, indicating their potential as antifungal agents against this fungus. The mechanism of fungitoxicity may involve interference with the fungus's metabolic processes, or other vital functions, leading to the observed inhibition of growth [21].

3.2 Evaluation of antagonists

In present study, four antagonists were subjected to *in vitro* evaluation against *G. lucidum* using the dual culture technique, and percent inhibition of the respective pathogens over control was calculated. The results indicated that all the evaluated antagonists

exhibited fungistatic activity, and significantly inhibited the mycelial growth of *G. lucidum* (Table 2). Of the four bio-agents tested, *T. harzianum* and *T. asperellum* exhibited the highest effectiveness, recorded minimal mycelial growth of 1.83 cm and 2.01 cm, respectively. This corresponded to the highest mycelial growth inhibition, with values of 79.63% and 77.57%, respectively, compared to the control (9.0 cm). *Pseudomonas fluorescens* and *Bacillus subtilis* demonstrated comparatively lower effectiveness, with radial mycelial growth of 4.67 cm and 6.28 cm along with inhibition percentage of 48.08% and 30.15% against the test pathogen, respectively. The control plate exhibited the maximum growth of *G. lucidum*, measuring 9.00 cm.

Thus, all the antagonists' agents tested *in vitro* demonstrated fungistatic activity against *G. lucidum*, with the fungal antagonists proving more effective than bacterial antagonists in inhibiting the tested pathogen. These findings are in consistent with those documented by previous researchers such as [22], [23]; and [24]. Their studies reported that *Trichoderma* antagonists significantly suppressed the growth of *G. lucidum* in comparison to *Pseudomonas fluorescens* and *Bacillus subtilis*. According to [1], the research findings revealed significant growth reduction of *Trichoderma* compared to the control *in vitro* when exposed to *G. lucidum*. Among the *Trichoderma* species tested, *T. harzianum* exhibited the maximum reduction in radial growth of the test fungus (68.5%), followed by *T. viride* (65.7%) and *T. pseudokoningii* (61.11%), in comparison to the control.

Siddiquee [22] explored the antagonistic characteristics of 48 local isolates of *T. harzianum* against *Ganoderma boninense* (strain PER 71) using dual culture techniques. The findings revealed that every *T. harzianum* isolate inhibited the growth of PER 71, exhibited percentage values for the inhibition radial growth ranging from 47.86 to 72.06%. Various antagonists employ distinct mechanisms to combat pathogens. These mechanisms can include direct parasitism and lysis of the pathogen, competition with the pathogen for space and nutrients, enzymatic degradation, and the production of inhibitory compounds, like antibiotic compounds that are toxic to the pathogen, collectively contributing to the inhibition or suppression of *G. lucidum* growth [25].

Table 2. Effect of antagonists on radial mycelial growth of *G. lucidum*

SN	Treatments	Radial mycelium growth (cm)	% Growth inhibition
T ₁	<i>Trichoderma harzianum</i>	1.83	79.63
T ₂	<i>Trichoderma asperellum</i>	2.01	77.57

T ₃	<i>Bacillus subtilis</i>	6.28	30.15
T ₄	<i>Pseudomonas fluorescens</i>	4.67	48.08
T ₅	Control	9.00	-
	SE (m) ±	0.39	-
	CD (P=0.01)	1.58	-

Trichoderma species are recognized as myco-parasites that target various fungi, displaying an active hyphal parasitic nature [26]. The efficacy of *Trichoderma* as a biocontrol agent is attributed to several mechanisms, including competition for space and nutrients, the release of chitinolytic enzymes, myco-parasitism, and the production of an inhibitory compound [27] and [28]. The dual culture experiments likewise demonstrated that the presence of *Pseudomonas fluorescens* inhibited the growth of *G. lucidum* mycelium (48.08%), but to a lesser extent than the fungal antagonist. Pseudomonades have been shown to produce a diverse range of chemicals with antimicrobial activity against phyto-pathogenic fungi and bacteria [29]. Pseudomonades strains produce many antimicrobial compound, including various phenazines, phenazine-1-carboxylic acid and 2, 4-diacetylploroglucinol, pyoluteorin and pyrrolnitrin [30], which may contribute to the antifungal activity against *G. lucidum* in our study. New quinone and hydroquinone antibiotics generated by *Pseudomonas* spp. have been shown to be effective against Gram positive bacteria, certain fungi, and yeasts [31]. These antibiotics' antifungal efficacy was defined by their particular inhibition of fungal cell wall production. Lim [32] found that *Pseudomonas aeruginosa* B6 isolate had the highest percent growth inhibition (75.76%) against *Ganoderma boninense*, with a minimum inhibitory dose of 0.04 mg/ml. The compound that may contribute to this biocontrol function was identified as 3-demethylubiquinone-9 utilizing LC-MS.

4. CONCLUSION

Hence, the following conclusions can be drawn from the obtained results across various aspects. Among the chemicals examined, Copper oxychloride 50% WP, Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L SC and Metiram 55% + Pyroclostrubin 5% WG demonstrated greater effectiveness in inhibiting the growth of *G. lucidum*. In terms of bio-agents, the fungal antagonist *T. harzianum* and *T. asperellum* outperformed the bacterial antagonist's *P. fluorescens* and *B. subtilis* in suppressing the growth of *G. lucidum*.

FUTURE SCOPE

According to the findings of the current laboratory investigation, fungicides such as Copper oxychloride 50% WP or Fluxapyroxad 167 G/L + Pyraclostrobin 333 G/L SC or Metiram 55% + Pyroclostrobin 5% WG, as well as antagonists *T. harzianum* and *T. asperellum*, can be used in the field to manage *G. lucidum*.

5. REFERENCES

1. Nirwan B, Choudhary S, Sharma K, Singh S. *In vitro* studies on management of root rot disease caused by *Ganoderma lucidum* in *Prosopis cineraria*. Current Life Sciences. 2016; 2(4):118-126.
2. Alexopoulos CJ, Mims CW, Blackwell M. Introductory Mycology. John Wiley and Sons, Inc. 1996; p 869.
3. Skaria M, Smith GS, Gilbertson RL. Root rot of young citrus caused by a species of *Ganoderma lucidum* complex. Phytopathology. 1990;80(10): 974.
4. Anonymous. Fruit crops estimates. Database of National Horticulture Board, Ministry of Agriculture, 2020-21; Govt. of India. <https://nhb.gov.in>
5. Palanna KB, Narendrappa T, Basavaraj S, Shreenivasa KR. Virulence analysis and influence of soil pattern and agronomic practices with respect to *Ganoderma* foot rot of arecanut in southern Karnataka. J. Plant. Crops. 2018; 46(1): 21-30.
5. Knorr LC. *Citrus Diseases and Disorders*. Gainesville, FL: University Presses of Florida.1973; pp. 103–105.
6. Skaria M, Farrald C. Biological and cultural factors associated with Citrus replant problems in Texas. Phytopathology. 1989: 79(10): 1193.
7. Assyiffa NMR, Natasha NZ, Hamzah AZ, Rasdli MZ. *In vitro* management of *Ganoderma* Basal stem rot disease on oil palm using wild Basidiomycetes fruiting on the ground after rainfall. Paper presented in 7th International Conference on Chemical, Agricultural, Biological and Environmental Sciences, held at Kuala Lumpur (Malaysia) during Dec. 14-15, 2017.
9. Idris AS, Ariffin D, Swinburne TR, Watt TA. The identity of *Ganoderma* species responsible for basal stem rot (BSR) disease of oil palm in Malaysia – pathogenicity test. MPOB Information Series No. 102. 2000; MPOB, Bangi.
10. Ryvardeen L. Studies in neotropical polypores2: a preliminary key to neotropical species of *Ganoderma* with laccate pileus. Mycologia. 2000; 92(1):180-191.
11. Nene YL, Thapliyal PN. Fungicides in Plant Disease Control. Second Edition Oxford & IBH Publishing, New Delhi. 1993; pp.531-532.
12. Vincent JM. Distortion of fungal hyphae in presence of certain inhibitors. Nature. 1947; 159:239-241.
13. Dennis C, Webster J. Antagonistic properties of species-groups of *Trichoderma*: Production of non-volatile antibiotics. Transactions of the British Mycological Society. 1971; 57(1), pp.25-IN3.

14. Bivi MR, Farhana SMN, Khairulmazmi A, Idris AS. Control of *Ganoderma boninense*: A causal agent of basal stem rot disease in oil palm with endophyte bacteria *in vitro*. *Int. J. Agric. Biol.* 2010; 12:833-839.
15. Gomez KA, Gomez AA. *Statistical Procedures for Agricultural Research*, 2edn, John Wiley and Sons, New York, NY, USA. 1984.
16. Bhaskharan R. Integrated management of basal stem rot disease of coconut. *Indian Coconut J.* 1993; 24:5-8.
17. George S., G. Chung and K Zakaria, 1996. Updated results on trunk injection of fungicides for the control of *Ganoderma* basal stem rot. In: *Proceedings of the 1996 PORIM International Palm Oil Congress-Agriculture Conference*, Kuala Lumpur, Malaysia. 1996; pp. 508-515
18. Naik RG, Venkatesh. Management of basal stem rot of coconut. *Ind. J. Agric. Res.* 2001; 35(2): 115-117.
19. Anonymous. Annual Progress Report for 2012-2013. Central Plantation Crops Research, Kasaragod, Kerala India. 2013; pp-142.
20. Srinivasulu B, Kumar KVK, Aruna K, Lakshmi MV, Rao DVR. Biointensive IDM approach against basal stem rot and stem bleeding diseases of coconut. *J. Plant. Crops.* 2006; 34(3): 502-507.
21. Jochen P, Zubrod, MircoBundschuh, Gertie Arts, Carsten A, Bruhl Gwenaël Imfeld, Anja Knabel, Sylvain Payraudeau, Jes J Rasmussen, Jason Rohr, Andreas Scharmüller, Kelly Smalling, Sebastian Stehle, Ralf Schulz, and Ralf B Schafer. Fungicides: An overlooked pesticide class. *Environmental Science and Technology.* 2019; 53 (7): 3347-3365.
22. Siddiquee S, Yusuf UK, Hossain K, Jahan S. *In vitro* studies on the potential *Trichoderma harzianum* for antagonistic properties against *Ganoderma boninense*. *J. Food. Agric. Environ.* 2009; 7(3&4): 970-976.
23. Bhansali R. *Ganoderma* diseases of woody plants of Indian arid zone and their biological control. In: Merillon JM, Ramawat KG, eds. *Plant Defence: Biological control*. Springer Verlag, The Netherlands, 2012;209-240.
24. Palanna KB, Shreenivasa KR, Basavaraj S, Narendrappa T. Review of genus *Ganoderma* causing basal stem rot (Coconut) and foot rot (Areca nut) with respect etiology and management. *Int. J. Curr. Microbiol. App Science.* 2020; 9(4): 1434-1455.
25. Japanis Feodora Grace, Yow San Chan and Khim Phin Chong. Evaluation on the effectiveness of combination of biocontrol agents in managing *Ganoderma boninense* of oil palm. *Malays. J. Microbiology.* 2021;17(1):1-10.
26. Susanto A, Sudharto PS, Purba RY. Enhancing biological control of basal stem rot disease (*Ganoderma boninense*) in oil palm plantations. *Mycopathologia.* 2005; 159: 153–157.
27. Iyer Rohini, Parvathy, Meera G, Lekha Hegde, Vinayak, Gunasekharan M. Management of basal stem rot disease of *Areca catechu* L. in India. *J. Plantn Crops.* 2004; 32(1): 25-27.
28. Chakrabarty R, Ray AK. *In vitro* studies on basal stem rot of areca nut caused by

- Ganoderma lucidum* (Curtis Ex. Fr.) Karst. J. Plant. Crops. 2007; 35 (1):39-41.
29. Cazorla FM, Duckett SB, Bergstroem ET, Noreen S, Odijk R, Lugtenberg BJ, Thomas-Oates JE, Bloemberg GV. Biocontrol of avocado dematophora root rot by antagonistic *Pseudomonas fluorescens* PCL1606 correlates with the production of 2-hexyl 5-propyl Resorcinol. Mol. Plant Microbe Interaction. 2006; 19: 418-428.
 30. Paul D, Sarma YR. Plant growth promoting rhizobacteria (PGPR) mediated root proliferation in black pepper (*Piper nigrum* L.) as evidenced through GS root software. Arch. Phyto. Plant Protection. 2006; 39: 311-314.
 31. Kumagai K, Kato M, Nabeshima S, Tsuyumu S. T 1801 A, B, C, and D, new quinone and hydroquinone antibiotics produced by a strain of *Pseudomonas*. Biosci, Biotechnol. and Biochemistry. 1992; 56(9): 1439-1442.
 32. Lim Pei Hui, Jualang Azian Gansau, Khim Phin Chong. Biocontrol of basal stem rot pathogen *Ganoderma boninense* by *Pseudomonas aeruginosa*. Bangladesh J. Botany. 2019; 48 (2): 209-215.