

“Assessment of Physico-chemical properties of Tea Garden soils of Darjeeling, West Bengal, India.”

ABSTRACT

The experiment was conducted in 2023 to assess the physical and chemical properties of tea garden soils of Darjeeling, West Bengal within its geographical coordinates ranging from 27° 2' 9.6252"N latitude to 88° 15' 45.6192"E longitude. Nine distinct villages located in the Darjeeling district covering Glenburn, Soureni, and Selimbong tea gardens ~~were~~ chosen for the study. The physical ~~properties~~ of tea garden soils, soil colour, texture, bulk density (Mg m^{-3}), particle density (Mg m^{-3}), percent pore space and percent water holding capacity were analysed ~~with the following standard procedure Munsell, (1971), Bouyoucos (1927), Muthuvel et al., (1992) and chemical properties pH, EC (dSm^{-1})_{at 25°C}, percent organic carbon, available nitrogen, phosphorus and potassium (kg ha^{-1}), Ca^{+3} $\text{C mol(p}^+ \text{) kg}^{-1}$ and Mg^{+3} $\text{C mol(p}^+ \text{) kg}^{-1}$ were analysed by following Jackson (1958), Wilcox (1950), Walkley and Black (1947), Subbiah and Asija (1956), Olsen et al., (1954), Toth and Prince (1949) and Jackson (1973) at 0-15, 15-30 and 30-45 cm depths. The results showed that the soil colour varied from brown to yellowish brown in dry condition and dark brown to dark reddish brown in wet ~~conditions~~. The soils of Selimbong were sandy loam to sandy clay loam, Soureni ~~were had~~ clay loam ~~soils~~, and soils of Glenburn were clay loam to sandy clay loam in texture. The variation in bulk density and particle density ~~was/were~~ recorded non-significant whereas percent pore space and percent water holding capacity were found significant. While organic carbon, available NPK, exchangeable calcium and magnesium were recorded significant, the pH and EC were found ~~to be non-significant~~.~~

Keywords: Physico-chemical properties, soil analysis, tea garden soil, Darjeeling.

INTRODUCTION

One of the most vital natural resources for human well-being is soil. (Zubber et al. 2021). The Darjeeling hill region of West Bengal, India, is renowned for producing the world's best-quality tea due to its optimal combination of altitude, sloping terrain, fertile soil, and climate (Datta, 2010). Tropical to subtropical climates with more than 200 cm of annual

Commented [A1]: From the journal's website: Abstract The abstract should be concise and informative. It should not exceed 300 words in length. It should briefly describe the purpose of the work, techniques and methods used, major findings with important data and conclusions. **Different sub-sections**, as given below, should be used. **No references should be cited in this part.**

Commented [A2]: There are three accepted ways of giving GPS coordinates. Either E and N, OR degrees, minutes and seconds or the +/- digital way. But you are not allowed to mix them, as you have done.

Commented [A3]: Nine villages = plural were

Commented [A4]: And chemical?

Commented [A5]: Mg? Don't you mean kg? Stick to SI units.

Commented [A6]: List the elements also, but not the methods

Commented [A7]: Why is this in subscript?

Commented [A8]: You see why you should stick to kg and not use megagrams? What is Mg? Magnesium or megagram/

Commented [A9]: You are following the methods, not the people.

Commented [A10]: 1927 in reference list

Commented [A11]: 1934 in reference list

Commented [A12]: 1927 in reference list

Commented [A13]: Two types of density

Commented [A14]: Which significance level did you use? 95%? You have to say. Better still, give the *P*-values.

Commented [A15]: You need to rewrite this whole abstract, omitting the references. You do not need to say which methods you used – that is why there is a Materials and Methods section. You need to give a summary. See the guidelines of the journal.

Commented [A16]: You need to add a concluding remark here.

Commented [A17]: Wrong reference style. From the journal's website: References must be listed at the end of the manuscript and numbered in the order that they appear in the text. Every reference referred in the text must also present in the reference list and vice versa. In the text, citations should be indicated by the reference number in brackets [3].

Commented [A18]: Not in reference list

Commented [A19]: Not in reference list

precipitation are ideal for the growth of tea (Ray et al., 2012). Tea grows well in sandy loam, diverse soil, slope, acidic, and well-drained environments (Tarun et al. 2018). The region's soil makeup is mostly made up of rocky, clay, and loamy soils, each of which has unique qualities (Pramanik, 2016). The world-renowned Darjeeling tea is of exceptional quality and is renowned for its muscatel taste (Anil et al. 2016). Organic tea cultivation in Darjeeling hills is getting popularity day by day due to growing health awareness of consumers, fetching good prices and high demand in international market: (Bisen & Singh, 2012).

MATERIALS AND METHODS

Soil samples were collected from nine different villages which lie in the Selimbong, Soureni, and Glenburn tea gardens of Darjeeling District at three respective depths of 0-15, 15-30 and 30-45 cm for the analysis of physical and chemical properties of soil. The total of 81 soil samples were collected from three tea gardens, in each of nine villages and at three depths out of which 27 samples are representing three depths of a tea garden to analyse the physical and chemical properties. These samples were air dried in shade for one week to obtain constant weight then crushed with wooden hammer, after that it was sieved with 0.2 mm sieve to obtained composite samples of each site and each depth. The physical properties of soils, soil colour, texture, bulk density

(Mg m^{-3}), particle density (Mg m^{-3}), percent pore space and percent water holding capacity were analysed with the following standard procedures: Munsell, (1971), Bouyoucos (1927), Muthuvel et al., (1992) and chemical properties pH, EC (dSm^{-1}) at 25°C , percent organic carbon, available nitrogen, phosphorus and potassium (kg ha^{-1}), Ca^{+3} C mol(p^+) kg^{-1} and Mg^{+3} C mol (p^+) kg^{-1} were analysed by following Jackson (1958), Wilcox (1950), Walkley and Black (1947), Subbiah and Asija (1956), Olsen et al., (1954), Toth and Prince (1949) and Jackson (1973) at 0-15, 15-30 and 30-45 cm depths. The data recorded during the course of the investigation was collected according to a completely randomized design, as per the method "Analysis of Variance technique" as given described by Fisher (1960).

RESULTS AND DISCUSSION

The results depicted in table 1 shows that the soil colour in dry condition varied from brown (7.5YR 5/4) to dark yellowish brown (10YR 3/4) in 0-15, brown (7.5YR 5/4) to yellowish brown (10YR 5/4) in 15-30 and brown (7.5YR 5/4) to pale brown (7.5YR 6/3) in 30-45 cm

Commented [A20]: Not in reference list

Commented [A21]: Makeup and made up are synonyms.

Commented [A22]: Not in reference list

Commented [A23]: Why did you decide on this research? What was the need you addressed? Since Rei et al did a similar and recent study. How did your work differed from theirs?

Commented [A24]: Because you did more than one analysis, you need to use the plural – analyses.

Commented [A25]: Only one sample per garden per depth? Not good. You should have taken at least 3 samples per plot and then bulked them. Much more representative than only one.

Commented [A26]: If that is what you mean?

Commented [A27]: What exactly do you mean? Maybe break the sentence in two and make this another sentence. Do you mean that there are 27 gardens with 3 depths each, thus 81 samples in total? Not clear.

Commented [A28]: Do you mean weight? I think you mean mass. From the internet: Mass (g) is the measure of the amount of matter in a body. Weight (N) is the measure of the amount of force acting on a mass due to the acceleration due to gravity

Commented [A29]: So you did take more than one sample per garden per depth? It is unclear.

Commented [A30]: You need to put the method next to the analysis: soil colour (Munsell, 1971) but of course, if you follow the guidelines of this journal, you will only have the reference number instead of the name and year.name and year.

Commented [A31]: Why is this in subscript?

Commented [A32]: You do not mention chemical analyses, but you measured chemicals. Amend accordingly.

Commented [A33]: The only Jackson in reference list is 1927

Commented [A34]: 1934 in reference list

Commented [A35]: The only Jackson in reference list is 1927

depth and in wet condition from dark brown (7.5YR 4/4) to dark reddish brown (5YR 3/3) in 0-15, brown (7.5YR 5/4) to dark brown (7.5YR 4/4) in 15-30 and yellowish brown (10YR 5/4) to dark brown (7.5YR 4/4) in 30-45cm depth. The Yellow-yellow colour of the soil was probably due to hydrated iron oxides. The brown colour indicates that the soil contains a high concentration of iron oxides and the dark colour suggests a high level of organic matter in the soil. Wet soils are darker due to similarity in refractive properties of water and soil (Anushka *et al.* 2021). Similar results were also reported by Nayak *et al.*, (2002). The results depicted in table 2 shows that the soil texture of Selimbong ranged from sandy clay loam (sand 64.3%, silt 13.9% and clay 21.8%) to sandy loam (sand 67.0%, silt 17.8% and clay 15.2%). The soil texture of Soureni was found to be clay loam (sand 26.2%, silt 39.4% and clay 32.6). The soil texture of Glenburn ranged from clay loam (sand 48.9%, silt 21.9% and clay 29.2) to sandy clay loam (sand 53.8%, silt 26.1% and clay 21.1%). It may be due to igneous and metamorphic rocks at an altitude of 2500 m in Darjeeling that have been eroded by various physical, chemical and biological weathering disorders to form such type of soil (Zubber *et al.* 2021). The results were found similar to that of Majumdar *et al.*, (2014). The bulk density of soil at 0-15, 15-30 and 30-45 cm depths respectively, was found to be non-significant at 5% critical difference, ~~the~~ The maximum bulk density was 1.22 Mg-kg m⁻³ at 0-15, 1.13 Mg-kg m⁻³ at 15-30 and 1.11 Mg-kg m⁻³ at 30-45 cm depths respectively found in at Selimbong. The minimum bulk density was 1.01 Mg-kg m⁻³ at 0-15 and 30-45 found in Selimbong and Glenburn respectively and 1.02 Mg-kg m⁻³ at 15-30 cm depth found in Selimbong mentioned in figure 1. The higher values of bulk density indicates that the soil is widely composed of clay and aggregated loams and the lower values indicate the presence of high organic matter (Pratistha *et al.*, 2020). The results were found to be similar to that of Wankhade *et al.*, (2015). The results graphically presented in figure 1 also shows the particle density of the soils at 0-15, 15-30 and 30-45 cm depths respectively and was found to be non-significant at 5% critical difference, ~~the~~ The maximum particle density was 2.58 Mg-kg m⁻³ at 0-15, 2.63 Mg-kg m⁻³ at 15-30 and 2.64 Mg-kg m⁻³ at 30-45 cm depth found in Soureni. The minimum particle density was 2.20 Mg-kg m⁻³ at 0-15 found in Glenburn and 2.21 Mg-kg m⁻³ in 15-30 as well as 30-45 cm depths found in Selimbong and Glenburn respectively. The high particle density values indicates that the soil has comparatively low organic matter content and the low particle density values indicates the presence of organic matter content (Anushka et al., 2021). Similar results were obtained by Barthwal et al., (2019). The results mentioned in table 3 shows that the percent pore space of the soil at 0-15, 15-30 and 30-45 cm depths respectively, was found to be significant at 5% critical difference, ~~the~~ The

Commented [A36]: Did you measure the iron oxide? If not, you need a reference here as well as the next two mentions about the reason for a specific colour. And remember, you cannot say it IS so, just that it may be so.

Commented [A37]: This should be Rai et al. Anushka is a first name, NOT a surname.

Commented [A38]: Results = plural = show

Commented [A39]: Not in reference list

Commented [A40]: If it were non-significant, I would not bother to add the graphs. I would suggest mention the bulk density range for each village and nothing more. Since it is not significant.

Commented [A41]: You do not have the names of the villages in the graph, so it is difficult to correlate the text with the graphs. Another reason I would omit the graph.

Commented [A42]: See my comment for bulk density: Ditto

Commented [A43]: You say there is no significance. So how can you talk about the reasons for high and low particle density? Are these non-significant values deemed high or low? Since it cannot be both. Once you say if it is high or low you can go on and discuss why it is high or low.

Commented [A44]: If this is not Rai et al, you need to list Anushka in the reference list.

maximum percent pore space 68.96% at 0-15, 62.96% at 15-30 cm were found in Glenburn and 59.86% at 30-45 cm depth found in Soureni. The minimum percent pore space was 52.96% at 0-15 cm and 50.30% at 15-30 cm found in Glenburn ~~and while~~ 50.53% at 30-45 cm depth ~~was~~ found ~~in at~~ Selimbong. The pore space ~~decreases-decreased~~ with increase in depth and it may be due to the lower content of organic matter present in the sub-surface soils than that of the surface soils (Pratistha *et al.*, 2020). ~~The s~~Similar findings were also reported by Pandey *et al.*, (2013). The water holding capacity of soil at 0-15, 15-30 and 30-45 cm depths respectively, was found to be significant at 5% critical difference, ~~the~~The maximum water holding capacity was 79.43% at 0-15, 78.83% at 15-30 and 72.66% at 30-45 cm depths respectively was recorded in Soureni. The minimum 60.23% at 0-15 cm was found in Glenburn, 53.36% at 15-30 cm depth found in Selimbong and 61.01% at 30-45 cm depth was found in Glenburn (~~presented in~~ tableTable 3). The sandy loam and clay loam soils indicate high clay content which can absorb a considerable amount of water for a long period in the micropores (Anushka *et al.*, 2021). Water holding capacity depends upon number, capillary spaces of the soil and size of the soil's pore spaces (Shirsath, 2021). These findings were similar with to that of Deb *et al.*, (2013). The results mentioned in figure 2 shows the soil pH and EC dSm^{-1} at 25°C at 0-15, 15-30 and 30-45 cm depths repectively ~~was~~ were found to be non-significant at 5% critical difference, the maximum soil pH 5.16 at 0-15, 5.28 at 15-30 as well as 30-45 cm depth was found in Selimbong. The minimum soil pH 4.40 at 0-15 as well as 15-30 and 4.46 at 30-45 cm depth was found in Soureni. The results indicated that the soil is highly acidic in nature. This level of high acidity may be due to heavy rainfall and leaching persisting in the region (Majumdar *et al.*, 2014). Similar findings were recorded by Ganorker *et al.*, (2017). Similarly, the maximum EC 0.13 at 0-15 as well as 15-30 and 0.14 at 30-45 cm depth was found in Glenburn. The minimum EC 0.03 at 0-15, 0.05 at 15-30 and 0.06 at 30-45 cm depth was found in Selimbong. The low values of EC might be due to leaching of soluble salts by heavy rainfall (Ray and Mukhopadhyay~~et al.~~, 2012). These findings were similar with to that of Santhi *et al.*, (2017). The per cent organic carbon of soil at 0-15, 15-30 and 30-45 cm depths respectively was found to be significant at 5% critical difference, ~~The~~ The maximum per cent organic carbon 2.02 at 0-15 cm was found in Selimbong, while 1.45 at 15-30 cm and 1.42 at 30-45 cm depths was found in Soureni. The minimum per cent organic carbon was recorded in Selimbong with 1.46 at 0-15 cm, 0.74 at 15-30 and 0.63 at 30-45 cm depths respectively ~~was recorded in Selimbong~~ presented (in tableTable 3). The high per-centage organic carbon ~~seem~~ present in the surface soils may be due to the undecomposed and partially decomposed organic matter and lower values seen in the sub-surface soils may be due to decomposed organic matter which has

Commented [A45]: 2018 in reference list

Commented [A46]: Is sandy loam also high in clay?

Commented [A47]: Not in reference list

Commented [A48]: 2018 in reference list

Commented [A49]: Results=plural show

Commented [A50]: Why subscript?

Commented [A51]: If not significant, why bother to give the numbers?

Commented [A52]: Not in reference list

Commented [A53]: Not in reference list

undergone chemical and biological changes already (Anushka *et al.* 2021). These findings were similar ~~with to~~ that of Arya, (2014). ~~The~~ table 4 shows the available nitrogen ~~in~~ kg ha⁻¹ of soil at 0-15, 15-30 and 30-45 cm depth ~~repectively, where it~~ was found ~~to be~~ significant at 5% critical difference. ~~The~~ maximum available nitrogen ~~of~~ 696.3 kg ha⁻¹ at 0-15 ~~and~~ 340.7 kg ha⁻¹ at 15-30 ~~cm respectively~~ was recorded in Soureni, ~~and while~~ 209.73 kg ha⁻¹ at ~~the~~ 30-45 cm depth was recorded in Selimbong. The minimum available nitrogen ~~of~~ 441.3 kg ha⁻¹ at 0-15 ~~cm depth was~~ found in Selimbong, ~~while 212.7 kg ha⁻¹ was found at~~ 15-30 ~~cm~~ and 177.70 kg ha⁻¹ at ~~the~~ 30-45 ~~cm depth was found in depth~~ at Glenburn. The high nitrogen content may be due to leaf litter or application of organic ~~amendments~~ in the soil. It may also be because of the high organic matter content in soil and addition of ~~chemical~~ fertilizers by the tea growing farmers- (Pratistha *et al.*, (2020). These findings were similar with that of Mishra *et al.*, (2000). The available phosphorus of soil at 0-15, 15-30 and 30-45 cm depths ~~respectively~~, was found ~~to be~~ significant at 5% critical difference. ~~the~~ ~~The~~ maximum available phosphorus ~~of~~ 29.54 kg ha⁻¹ at 0-15, 27.79 kg ha⁻¹ at 15-30 and 25.02 kg ha⁻¹ at 30-45 cm depths ~~s~~ was found in Glenburn. The minimum available phosphorus was ~~found in Soureni, with~~ 17.23 kg ha⁻¹ at 0-15, 16.55 kg ha⁻¹ at 15-30 and 16.13 kg ha⁻¹ at 30-45cm depth ~~found in Soureni (presented in table~~ Table 4). The available phosphorus content may be due to the influence of soil pH which is acidic in nature and often limits nutrients ~~that remains~~ present in plant nuclei and act as an energy storage (Kekane *et al.*, 2015). These findings were similar with that of Ram *et al.*, (2014). ~~The~~ table 4 also shows the available potassium of soil at 0-15, 15-30 and 30-45cm depths ~~respectively~~ was found ~~to be~~ significant at 5% critical difference. ~~the~~ ~~The~~ maximum available potassium was 334.92 kg ha⁻¹ at 0-15, 322.32 kg ha⁻¹ at 15-30 and 320.32 kg ha⁻¹ at 30-45 cm depths ~~respectively and it was~~ found ~~in~~ at Selimbong. The minimum available potassium was ~~found in Glenburn, with~~ 101.43 kg ha⁻¹ at 0-15, 99.90 kg ha⁻¹ at 15-30 and 97.43 kg ha⁻¹ at 30-45 cm depths ~~respectively found in Glenburn. High~~ ~~The~~ ~~high~~ proportion of potassium may be due to the feldspar and muscovite mica found in the Darjeeling region (Laitpharlang Cajee, 2018). These findings were similar with that of Ram *et al.*, (2014). The results ~~mentioned~~ in figure 3 ~~shows~~ ~~that~~ the exchangeable calcium of soil at 0-15, 15-30 and 30-45cm depths ~~s~~ was found significant at 5% critical ~~difference~~, ~~with~~ the maximum exchangeable calcium 0.42 C mol (p⁺) kg⁻¹ at 0-15, 0.39 C mol (p⁺) kg⁻¹ at 15-30 and 0.38 C mol (p⁺) kg⁻¹ at 30-45 cm depths ~~respectively~~. ~~found in~~ at Selimbong. The minimum exchangeable calcium was ~~found at Soureni, with values of~~ 0.03 C mol (p⁺) kg⁻¹ at ~~both~~ 0-15 ~~as well as and~~ 15-30 cm depths, and 0.02 C mol (p⁺) kg⁻¹ at 30-45 cm depth ~~found in Soureni~~. Very low values of calcium ~~were~~

Commented [A54]: Check this reference that is not in the list

Commented [A55]: Not in reference list

Commented [A56]: Differences are significant or not. How was this significance measured – between depths or between sites?

Formatted: Superscript

Commented [A57]: If this is a common practice in the area, you need to mention it and also what organic amendments, since that can influence the minerals in the soil.

Commented [A58]: See my above comment

Commented [A59]: Not in reference list

Commented [A60]: Wat was significant. The differences between the depths or between the towns? You need to be more precise.

Commented [A61]: It ease reading if you refrain from using the exact same sentence structure.

Commented [A62]: 2016 in reference list

Commented [A63]: Difference between what? Depth or sites?

Commented [A64]: What is deemed high?

Commented [A65]: Proportion?

Commented [A66]: It is not common to refer to an author's first name. Stick with the surname.

Commented [A67]: 2016 in reference list

Commented [A68]: Results = plural = show

Commented [A69]: Difference between what exactly?

Commented [A70]: Isn't this supposed to be centimol cmol? Because then the c should not be a capital letter. And the p? What does that stand for?

Commented [A71]: What is deemed 'normal', if you say this is very low?

recorded which may be due to the leaching of calcium as hydrogen is added to the soil by decomposition of organic matter as well as due to heavy rainfall (Pratistha *et al.*, 2020). These findings were similar with that of Ray *et al.*, (2012). The exchangeable magnesium of soil at 0-15, 15-30 and 30-45 cm depths was found significant at 5% critical difference, the maximum exchangeable magnesium 0.22 C mol(p⁺) kg⁻¹ at 0-15, 0.20 C mol(p⁺) kg⁻¹ at 15-30 and 0.19 C mol(p⁺) kg⁻¹ at 30-45cm depth found in Selimbong. In the case of Soureni, the minimum exchangeable magnesium was 0.02 C mol(p⁺) kg⁻¹ at the 0-15 as well as the 15-30 cm depths and while it was 0.01 C mol(p⁺) kg⁻¹ at 30-45 cm depth found in Soureni presented graphically in Figure 3). Very low magnesium was recorded in all sites which may be due to heavy rainfall prevailing in the area and surface leaching of basic cations (Mg⁺²) by the excessive rainfall, as also reported by (Ray *et al.*, (2012). These findings were similar with that of Ray *et al.*, (2012).

Commented [A72]: And the natural acidic nature of the soils?

Commented [A73]: Again: was the different depths the significant factor or was it the different sites/villages?

Commented [A74]: Your readers all know the reference to a figure means graphic info.

Commented [A75]: What is deemed normal Mg levels?

Table 1: Soil colour of different villages in dry and wet condition of tea garden soils of Darjeeling district at 0-15cm, 15-30 and 30-45cm

Village	Dry condition			Wet condition		
	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm
Pokhringung Khasmahal (V1)	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown	7.5YR 5/4 Brown	5YR 3/3 Dark Reddish Brown	7.5YR 5/4 Brown	7.5YR 4/4 Dark Brown
Soolbongs Tea Garden (V2)	7.5YR 5/4 Brown	7.5YR 6/3 Pale Brown	7.5YR 6/3 Pale Brown	10YR 5/4 Yellowish Brown	10YR 6/4 Light Yellowish Brown	10YR 5/4 Yellowish Brown
Selimbong (V3)	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown	7.5YR 5/4 Brown	2.5YR 4/4 Reddish Brown	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown
Phuguri Forest (V4)	10YR 5/4 Yellowish Brown	10YR 6/6 Brownish Yellow	7.5YR 6/3 Pale Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown
Sourinibasti (V5)	10YR 3/4 Dark Yellowish Brown	10YR 5/4 Yellowish Brown	10YR 8/6 Pale Yellow	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown	7.5YR 5/4 Brown
Singbulli Tea Garden (V6)	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	10YR 6/6 Brownish Yellow	7.5YR 5/4 Brown	7.5YR 6/3 Pale Brown	7.5YR 4/4 Dark Yellow
Singringtam (V7)	7.5YR 5/4 Brown	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown
Kambal Tea Garden (V8)	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	7.5YR 5/4 Brown	7.5YR 4/4 Dark Brown	5YR 3/4 Dark Reddish Brown	7.5YR 4/4 Dark Brown

Commented [A76]: You can give the cm at each number or at the end, but not at the first and the last. Thus: 0-15c, 15-30 and 30-45 cm OR 0-15 cm, 15-30 cm and 30-45 cm. And please leave a space between number and unit. From the internet: Include a space between numbers and units: for example, 75 kg, but no space with percentages, e.g. 75%.

Manedara (V9)	7.5YR 5/4 Brown	10YR 5/4 Yellowish Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown
----------------------	--------------------	--------------------------------	-------------------------	-------------------------	-------------------------	--------------------

Table 2: Soil texture of different villages of tea garden soils of Darjeeling district at 0-15 cm, 15-30 and 30-45 cm depths.

Tea Garden	Village	Sand %	Silt %	Clay %	Textural Class
Selimbong	V1	64.3	13.9	21.8	Sandy Clay Loam
	V2	67.0	17.8	15.2	Sandy Loam
	V3	61.5	22.0	16.5	Sandy Loam
	V4	26.2	39.2	34.6	Clay Loam
Soureni	V5	29.4	37.5	33.1	Clay Loam
	V6	23.9	37.1	39	Clay Loam
	V7	48.9	21.9	29.2	Clay Loam
Glenburn	V8	45.9	29.3	24.8	Clay Loam
	V9	53.8	26.1	21.1	Sandy Clay Loam

Table 3: Pore space, water holding capacity and percent organic carbon of soil in different villages of Darjeeling at 0-15 cm, 15-30 cm and 30-45 cm depths

Villages	Pore space			Water holding capacity			Organic carbon			
	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	
Selimbong	V1	57.16	54.73	50.53	69.70	65.06	68.13	1.46	0.74	0.71
	V2	54.63	57.83	56.14	71.03	53.36	63.16	2.02	1.02	0.63
	V3	55.50	54.36	52.13	69.93	70.53	68.03	1.76	1.02	0.86
	V4	53.13	52.30	56.73	71.96	71.16	70.69	1.87	1.30	1.25
Soureni	V5	56.43	55.63	55.33	79.43	78.83	70.86	1.81	1.45	1.42
	V6	57.66	55.40	59.86	74.13	70.05	72.66	1.71	1.27	1.27
	V7	52.96	50.30	52.06	70.56	68.68	68.40	1.59	1.44	1.31
Glenburn	V8	68.96	62.67	54.36	65.96	63.86	66.25	1.51	1.27	1.03
	V9	58.73	62.96	53.23	61.01	62.51	60.23	1.53	1.15	1.09
F-test	S	S	S	S	S	S	s	s	s	
S. Em. (±)	4.84	4.28	2.87	5.27	7.08	3.74	0.18	0.22	0.27	
C.D @5%	1.24	4.39	4.4	1.92	0.57	8.73	3.10	1.41	1.59	

Table 4: Available potassium, available nitrogen, and available phosphorus of soil in different villages of Darjeeling at 0-15 cm, 15-30 cm and 30-45 cm depths respectively.

Villages	Available potassium			Available nitrogen			Available phosphorus			
	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	
Selimbong	V1	314.44	306.67	302.16	484.0	268.5	209.73	25.35	25.02	24.35
	V2	334.92	322.32	320.32	441.3	241.5	194.56	20.12	19.90	19.25
	V3	278.84	273.00	268.47	442.9	279.6	193.86	17.86	17.29	16.13
Soureni	V4	125.76	124.02	120.88	612.1	282.7	189.90	18.72	17.28	16.89
	V5	110.92	106.10	100.64	679.4	314.2	189.03	24.03	23.43	22.84
	V6	149.90	145.31	141.08	696.3	340.7	182.30	17.23	16.55	16.20
Glenburn	V7	219.46	214.37	210.83	543.5	217.5	183.16	28.79	27.79	25.02
	V8	143.84	138.48	136.76	553.8	212.7	185.76	24.26	22.87	21.73
	V9	101.43	99.90	97.43	574.8	275.4	177.70	29.54	27.50	24.13
F-test	s	s	s	s	s	s	s	s	s	
S. Em. (±)	91.29	88.54	88.53	93.30	41.96	9.35	4.61	4.39	3.65	
C.D @5%	5.63	9.58	5.99	1.10	2.81	3.16	0.001	0.008	0.138	

Fig 1: Graphical representation of bulk density ($Mg\ kg\ m^{-3}$) and particle density ($Mg\ kg\ m^{-3}$) of soil in different villages of tea garden soil of Darjeeling at 0-15cm, 15-30 cm and 30-45 cm depths respectively.





Fig 2: Graphical representation of soil pH and EC (dSm^{-1}) at 25°C of soil in different villages of tea garden soil of Darjeeling at 0-15 cm, 15-30 cm and 30-45 cm depths respectively. And please add where V1-V9 is located.

Commented [A77]: Why subscript?





Fig 3: Graphical representation of exchangeable calcium ($\text{C mol}(\text{p}^+) \text{kg}^{-1}$) and exchangeable magnesium ($\text{C mol}(\text{p}^+) \text{kg}^{-1}$) of soil in different villages of tea garden soil of Darjeeling at 0-15cm, 15-30 cm and 30-45cm depth



CONCLUSION

It is evident that the Darjeeling district soils are in good physical condition, which is favourable for tea cultivation. The sub-surface layer showed darker-coloured soil while the surface layer soil was found to be lighter in colour. The soil texture that was found to be dominant ~~is~~ was clay loam. The bulk density and particle density of the soil ~~was~~ were found to

Commented [A78]: Two different kinds of density

be in low levels. The high clay component of the soil was discovered to contribute to its good pore space and water-holding capacity. Low pH and EC values have been recorded. Mostly all the villages showed high organic carbon content as well as high available nitrogen concentration which may be due to the adoption of organic farming in the tea gardens. The soils had a medium to high range of phosphate and potassium, with values decreasing as depth increased. Very low values of exchangeable calcium and magnesium was found in all the regions which may be due to heavy rainfall prevailing in the area and surface leaching of basic cations (Ca^{+2} and Mg^{+2}) by the excessive rainfall. The results of the study indicated that maintaining the sustainability of tea crop output and quality may benefit from the implementation of good agriculture practises, which involve the application of organic amendments, herbicides, and micronutrients. However, application of more labile organic inputs, liming materials and suitable inorganic fertilizers (N-P-K) would be effective for sustainable management and improving fertility status of the soils under tea growing region.

References:

- Ahamed, Z. (2021). A study on Soil Problems in North Bengal, West Bengal, India. *Mazedan Chemical Research Journal*, 2(1), 1-3.
- Barthwal, A., Swaroop, N., Rao, P. S., & Thomas, T. (2019). Assessment of physical properties of soil in Dehradun district, Uttarakhand, India. *International Journal of Chemical Studies*, 7(3): 1623-1625.
- Bisen, J.S. and Singh A.K. (2012). Impact of inorganic to organic cultivation practices on yield of tea in Darjeeling hills – A case study. *Indian Journal of Horticulture* 69(2): 288-291
- Black, C.A. (1965). Methods of soil analysis. *American Society of Agronomy*, Madison, Wisconsin, USA, 2.
- Bouyoucos, G. J. (1927). The hydrometer as a new method for the mechanical analysis of soils. *Soil Science*, 23: 343-353. region, Sikkim, India.
- Cajee L., (2018). Physical Aspects of the Darjeeling Himalaya: Understanding from a Geographical Perspective. *IOSR Journal Of Humanities And Social Science (IOSR-JHSS)*. Volume 23, Issue 3, pp 66-79.
- Deb, P., Debnath, P., Denis, A.F and Lepcha, O.T. (2018). Variability of soil physico-chemical properties at different agro-ecological zones of Himalayan region, Sikkim, India. *Environmental Development and Sustainability*, Springer science + Business Media B.V, part of Springer Nature 2018.
- Fisher, R. A. (1960). The design of experiments. Seventh edition. *Hafner publishing company*, New York.
- Ghosh, H. N. and Kumar, R. (1999) Seasonal variation in photosynthesis and productivity of young tea. *Experimental Agriculture*. 35:71-85.

Commented [A79]: How did you discover this? I thought you read it. Discover implicates that you found it out during your study.

Commented [A80]: You said in the text: The high nitrogen content may be due to leaf litter or application of organic amendments in the soil. It may also be because of the high organic matter content in soil and addition of chemical fertilizers by the tea growing farmers

Commented [A81]: You never mentioned the application of herbicides, micronutrients and only mentioned once that some fertilizer is applied. How can you make a conclusion about things you never even mentioned? Your conclusion must be based on YOUR research.

Commented [A82]: This would increase the pH and I thought tea grew in acidic soils? From the internet: The most comfortable soil pH level for tea plants is between 4.5 to 5.5. So why would you want to increase the pH with lime?

Commented [A83]: The reference style is not according to the journal's website. From the journal's guidelines: note the abbreviated article names.

For Published paper:

1. Hilly M, Adams ML, Nelson SC. A study of digit fusion in the mouse embryo. *Clin Exp Allergy*. 2002;32(4):489-98.

Note: List the first six authors followed by et al.
Note: Use of DOI number for the full-text article is encouraged. (if available).

Commented [A84]: Not in text

Commented [A85]: Not in text

Commented [A86]: 2013 in text

Commented [A87]: Deb, Proloy & Debnath, Prankanu & Denis, Anjelo & Lepcha, Ong. (2019). Variability of soil physicochemical properties at different agroecological zones of Himalayan region: Sikkim, India. *Environment, Development and Sustainability*. 21. 10.1007/s10668-018-0137-8. If this is the same, your date is wrong.

Commented [A88]: Not in text

10. Jackson, M.L. (1927) Soil Chemical analysis. *Prentice hall of India Pvt. Limited*, New Delhi, 38:226.
11. Kekane S.S., Chavan R.P., Shinde D.N., Patil C.L., and Sagar S.S. (2015). A review on physico-chemical properties of soil. *International Journal of Chemical studies*. 3(4): pp 29-32.
12. Majumdar K, Ray DP, Chakraborty S, Pandit T. (2014) Change of nutrient status of hilly soil in Darjeeling district within five years. *International Journal of Bioresource Science*. 1(1):25-30.
13. Misra, T. K., Nanda, A.K., Mandal, P, and Saha, A. (2017) Physiochemical Properties of Soils under Different Tea Growing Regions of North Bengal: A study from 2006 to 2010. *International Journal of Research in Chemistry and Environment*. Vol 8 Issue 1, pp 44-48
14. Munsell, A. H. (1954). Munsell Soil Color Charts. *Munsell Color Company Inc.*, Baltimore.
15. Muthuvel, P., Udayasoorian, C., Natesan, R., & Ramaswami, P. R. (1992). *Introduction to Soil Analysis*. *Tamil Nadu Agricultural University*, Coimbatore.
16. Nayak, R.K., Sahu, G.C and Nanda, S.S.K (2002). Characterization and classification of soils of Central Research Station, Bhubaneswar. *Agropedology*, 12:1-8.
17. Olsen, S.R., Cole, C.V., Watanable, F.S and Dean, L.A (1954). Estimation of available phosphorus in soil by extraction with sodium bicarbonate, USDA, *Circular*, 939: 1-19.
18. Pandey V, Gautam P, Singh AP (2018). Assessment of physical properties of soil under different land use systems in a Mollisol. *Journal of Pharmacognosy and Phytochemistry*. 7(6):2645-2648.
19. Pramanik M.K (2016). Site suitability analysis for agricultural land use of Darjeeling district using AHP and GIS techniques. *Modeling Earth Systems and Environment* 2(2)
20. Pratistha, P., Swaroop, N., David, A. A., & Thomas, T. (2020) Assessment of physical properties of soil in Kalimpong district, West Bengal, India. *International Journal of Chemical Studies*. 8(4): 1718-1721.
21. Rai, A. and Thomas, T. (2021) Assessment of Physical Properties of Soils of Darjeeling District, West Bengal, India. *Biological Forum – An International Journal*. 13 (2). pp 481- 487.
22. Ram R, Chatterjee S, Maji C, Sharma PK, Rani P. (2016) Effect of soil health, nutrient management and soil test-based doses of lime on mulberry leaf yield (*Morus alba* L.) in acid soils of Kalimpong hills. *International Journal of Agriculture Sciences*. 8(59):3333-3337.
23. Ray, S.K and Mukhopadhyay, D. (2012) A Study on Physicochemical Properties of Soils under different tea growing regions of West Bengal (India). *International Journal of Agriculture Sciences*. Volume 4, Issue 8, pp 325-329.
24. Shirsath, W.B (2021). Quantitatively Physico-Chemical Analysis of Some Soil Samples of Satana (Baglan)Tahsil, District Nashik, Maharashtra (India). *Journal of Scientific Research*. Volume 65, Issue 7, pp 143-148.
25. Singh, A.K and Chauhan, R. (2016) Tea Research for Darjeeling Tea Industry: *Various Aspects. Tea: Technological Initiative*. pp 195-239.
26. Subba, B., Mondal, S., Mandal, U.K. and Tamang, L. (2024). Small-scale tea farming sustainability: A case study of Darjeeling Hill Region, India. *Farming System*. Volume 2, Issue 2. pp 110-117.

Commented [A89]: 1958 and 1973 in text

Commented [A90]: Not in text

Commented [A91]: 2013 in text

Commented [A92]: Not in text

Commented [A93]: You don't have all the authors.
Amend: Anushka Rai*, Tarence Thomas, Arun Alfred David and Raghunandan Singh

Commented [A94]: 2014 in text

Formatted: Font: Italic

Commented [A95]: Not in text

Commented [A96]: Not in text

27. Subbiah, B.V. and Asija, G.L. (1956). A rapid procedure for the estimation of available nitrogen in soils. *Current Science*, 25: 259-260.
28. Toth, S.J and Prince, A.L. (1949). Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by Flame Photometer method. *Soil Science*. 67:437445.
29. Walkley, A and Black, C.A (1934). An estimation for different methods for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*. 37:29-38.
30. Wankhade, R. R. (2015). Physico-chemical analysis of soil from some farms of Digras region of district Yavatmal in Maharashtra, India. *International Journal of Multidisciplinary & Allied Studies*. 2(10): 1-4.
31. Wilcox, L.V. (1950). Electrical Conductivity. *American Water Work Association Journal*. 42:775-776.

Commented [A97]: 1947 in text

UNDER PEER REVIEW