

# Assessment of Physico-chemical properties of Tea Garden soils of Darjeeling, West Bengal, India

## ABSTRACT

The experiment was conducted in 2023-2024 to assess the physical and chemical properties of tea garden soils of Darjeeling, West Bengal within its geographical coordinates ranging from 27° 2' 9.6252" N latitude to 88° 15' 45.6192"E longitude. Nine distinct villages located in the Darjeeling district covering Glenburn, Soureni, and Selimbong tea gardens were chosen for the study. The physical properties of tea garden soils, soil colour, texture, bulk density ( $\text{Mg m}^{-3}$ ), particle density ( $\text{Mg m}^{-3}$ ), percent pore space and percent water holding capacity were analysed at 0-15, 15-30 and 30-45 cm depths. The results showed that the soil colour varied from brown to yellowish brown in dry condition and dark brown to dark reddish brown in wet. The soils of Selimbong were sandy loam to sandy clay loam, Soureni were clay loam, and soils of Glenburn were clay loam to sandy clay loam in texture. The variation in bulk density and particle density was recorded non-significant with the values ranging from 1.01-1.22  $\text{Mg m}^{-3}$  and 2.21-2.64  $\text{Mg m}^{-3}$  respectively. The percent pore space and percent water holding capacity were found significant with the values ranging from 50.53-68.96% and 53.36-79.43% respectively. The soil pH ranged from 4.32-5.28 and electrical conductivity (EC) 0.03-0.97  $\text{dS m}^{-1}$  which were found to be non-significant. The organic carbon and available nitrogen were found to be significant with values ranging from 0.63-2.02% and 177.70-696.3  $\text{kg ha}^{-1}$  respectively. The available phosphorus ranged from 16.13-29.54  $\text{kg ha}^{-1}$  and available potassium 100.64-334.92  $\text{kg ha}^{-1}$  which was found to be significant. The values of exchangeable calcium and magnesium were recorded significant ranging from 0.02-0.42  $\text{C mol(p}^+) \text{ kg}^{-1}$  and 0.016-0.223  $\text{C mol(p}^+) \text{ kg}^{-1}$  respectively.

**Keywords:** Physico-chemical properties, soil analysis, tea garden soil, Darjeeling.

## INTRODUCTION

“One of the most vital natural resources for human well-being is soil. Among the eight districts of North Bengal, Darjeeling and Kalimpong are located in the foothills of the Himalayas [31-34]. The soil here is completely different from other parts of the state. This area is mainly found in hilly soils, forest soils, old alluvium and red soils” (Ahamed Z., 2021).

“Tropical to subtropical climates with more than 200 cm of annual precipitation are ideal for the growth of tea. Evaluations of soil fertility status under tea growing soils are necessary to make a proper management and sound fertilizer recommendation for optimizing the yield of tea” (Ray et al., 2012). “Tea grows well in sandy loam, diverse soil, slope, acidic, and well-drained environments. It is well known fact that for the plant growth i.e. optimizing the yield of tea and its quality-quantity of secondary metabolites is highly influenced by the soil physicochemical characters” (Mishra et al. 2018). “The region's soil makeup is of rocky, clay, and loamy soils, each of which has unique qualities” (Pramanik, 2016). “The world-renowned Darjeeling tea is of exceptional quality and is renowned for its muscatel taste. The tea is available in the form of black, green, white and oolong” (Anil et al. 2016). “Organic tea cultivation in Darjeeling hills is getting popularity day by day due to growing health awareness of consumers, fetching good prices and high demand in international market. Maintenance of the present tea crop productivity in the changing scenario of climate and mode of cultivation is a big challenge for every stakeholder of Darjeeling tea industry, while it has been established that the 12-30% reduction in yield is quite natural after conversion to organic” (Bisen & Singh, 2012).

## MATERIALS AND METHODS

Soil samples were collected from nine different villages which lie in the Selimbong, Soureni, and Glenburn tea gardens of Darjeeling District at three respective depths of 0-15, 15-30 and 30-45 cm for the analysis of physical and chemical properties of soil. A total of 81 soil samples were collected from three tea gardens, in each of nine villages and at three depths out of which 27 samples are representing three depths of a tea garden to analyse the physical and chemical properties. These samples were air dried in shade for one week to obtain constant weight then crushed with wooden hammer, after that it was sieved with 0.2mm sieve to obtain composite samples of each site and each depth. The physical properties of soils, soil colour, texture, bulk density ( $\text{Mg m}^{-3}$ ), particle density ( $\text{Mg m}^{-3}$ ), percent pore space and percent water holding capacity were analysed with the following standard procedures: Munsell, (1971), Bouyoucos (1927), Muthuvel et al., (1992) and chemical properties pH, EC ( $\text{dSm}^{-1}$ ) at 25°C, percent organic carbon, available nitrogen, phosphorus and potassium ( $\text{kg ha}^{-1}$ ),  $\text{Ca}^{+3} \text{C mol(p}^{+}) \text{kg}^{-1}$  and  $\text{Mg}^{+3} \text{C mol(p}^{+}) \text{kg}^{-1}$  were analysed by following Jackson (1958), Wilcox (1950), Walkley and Black (1947), Subbiah and Asija (1956), Olsen et al., (1954), Toth and Prince (1949) and Jackson (1973) at 0-15, 15-30 and 30-45 cm depths. The data recorded during the course of

investigation was subjected to statistical analysis by the method of analysis of Completely Randomized Design (CRD) as per the method of "Analysis of Variance technique" (ANOVA) (Fisher, 1960).

## RESULTS AND DISCUSSION

The results depicted in table 1 shows that the soil colour in dry condition varied from brown (7.5YR 5/4) to dark yellowish brown (10YR 3/4) in 0-15, brown (7.5YR 5/4) to yellowish brown (10YR 5/4) in 15-30 and brown (7.5YR 5/4) to pale brown (7.5YR 6/3) in 30-45cm depth and in wet condition from dark brown (7.5YR 4/4) to dark reddish brown (5YR 3/3) in 0-15, brown (7.5YR 5/4) to dark brown (7.5YR 4/4) in 15-30 and yellowish brown (10YR 5/4) to dark brown (7.5YR 4/4) in 30-45cm depth. The yellow colour of the soil was probably due to hydrated iron oxides. The brown colour indicates that the soil contains a high concentration of iron oxides and the dark colour suggests a high level of organic matter in the soil. Wet soils are darker due to similarity in refractive properties of water and soil (Rai *et al.* 2021). Similar results were also reported by Nayak *et al.*, (2002). The results depicted in table 2 show that the soil texture of Selimbong ranged from sandy clay loam (sand 64.3%, silt 13.9% and clay 21.8%) to sandy loam (sand 67.0%, silt 17.8% and clay 15.2%). The soil texture of Soureni was found to be clay loam (sand 26.2%, silt 39.4% and clay 32.6). The soil texture of Glenburn ranged from clay loam (sand 48.9%, silt 21.9% and clay 29.2) to sandy clay loam (sand 53.8%, silt 26.1% and clay 21.1%). It may be due to igneous and metamorphic rocks at an altitude of 2500 m in Darjeeling that have been eroded by various physical, chemical and biological weathering disorders to form such type of soil (Ahamed Z., 2021). The results were found similar to that of Majumdar *et al.*, (2014). The bulk density of soil at 0-15, 15-30 and 30-45 cm depths respectively, was found to be non-significant at 5% critical difference. The maximum bulk density was 1.22 Mg m<sup>-3</sup> at 0-15, 1.13 Mg m<sup>-3</sup> at 15-30 and 1.11 Mg m<sup>-3</sup> at 30-45 cm depths found in Selimbong. The minimum bulk density was 1.01 Mg m<sup>-3</sup> at 0-15 and 30-45 found in Selimbong and Glenburn respectively and 1.02 Mg m<sup>-3</sup> at 15-30 cm depth found in Selimbong mentioned in figure 1. The higher values of bulk density indicates that the soil is widely composed of clay and aggregated loams and the lower values indicate the presence of high organic matter (Pratistha *et al.*, 2020). The results were found to be similar to that of Wankhade *et al.*, (2015). The results graphically presented in figure 1 also shows the particle density of the soils at 0-15, 15-30 and 30-45 cm depths respectively and was found to be non-significant

at 5% critical difference. The maximum particle density was  $2.58 \text{ Mg m}^{-3}$  at 0-15,  $2.63 \text{ Mg m}^{-3}$  at 15-30 and  $2.64 \text{ Mg m}^{-3}$  at 30-45 cm depth found in Soureni. The minimum particle density was  $2.20 \text{ Mg m}^{-3}$  at 0-15 found in Glenburn and  $2.21 \text{ Mg m}^{-3}$  in 15-30 as well as 30-45 cm depths found in Selimbong and Glenburn respectively. The high particle density values indicates that the soil has comparatively low organic matter content and the low particle density values indicates the presence of organic matter content (Anushka *et al.*, 2021). Similar results were obtained by Barthwal *et al.*, (2019). The results mentioned in table 3 shows that the percent pore space of the soil at 0-15, 15-30 and 30-45 cm depths was found to be significant at 5% critical difference. The maximum percent pore space 68.96% at 0-15, 62.96% at 15-30 cm were found in Glenburn and 59.86% at 30-45 cm depth found in Soureni. The minimum percent pore space was 52.96% at 0-15 cm and 50.30% at 15-30 cm found in Glenburn while 50.53% at 30-45 cm depth was found at Selimbong. The pore space decreased with increase in depth and it may be due to the lower content of organic matter present in the sub-surface soils than that of the surface soils (Pratistha *et al.*, 2020). Similar findings were also reported by Pandey *et al.*, (2013). The water holding capacity of soil at 0-15, 15-30 and 30-45 cm depths respectively was found to be significant at 5% critical difference. The maximum water holding capacity was 79.43% at 0-15, 78.83% at 15-30 and 72.66% at 30-45 cm depths respectively was recorded in Soureni. The minimum 60.23% at 0-15 cm was found in Glenburn, 53.36% at 15-30 cm depth found in Selimbong and 61.01% at 30-45 cm depth was found in Glenburn (Table 3). The sandy loam and clay loam soils indicate high clay content which can absorb a considerable amount of water for a long period in the micropores (Rai *et al.*, 2021). Water holding capacity depends upon number, capillary spaces of the soil and size of the soil's pore spaces (Shirsath, 2021). These findings were similar to that of Deb *et al.*, (2013). The results mentioned in figure 2 show the soil pH and EC  $\text{dSm}^{-1}$  at  $25^\circ\text{C}$  at 0-15, 15-30 and 30-45 cm depths respectively were found to be non-significant at 5% critical difference. The maximum soil pH 5.16 at 0-15, 5.28 at 15-30 as well as 30-45cm depth was found in Selimbong. The minimum soil pH 4.40 at 0-15 as well as 15-30 and 4.46 at 30-45cm depth was found in Soureni. The results indicated that the soil is highly acidic in nature. This level of high acidity may be due to heavy rainfall and leaching persisting in the region (Majumdar *et al.*, 2014). Similar findings were recorded by Ganorker *et al.*, (2017). Similarly, the maximum EC 0.13 at 0-15 as well as 15-30 and 0.14 at 30-45cm depth was found in Glenburn. The minimum EC 0.03 at 0-15, 0.05 at 15-30 and 0.06 at 30-45cm depth was found in Selimbong. The low values of EC might be due to leaching of soluble salts by heavy rainfall (Ray and Mukhopadhyaya, 2012). These findings were similar to that of Rai *et*

al., (2021). The per cent organic carbon of soil at 0-15, 15-30 and 30-45 cm depths was found to be significant at 5% critical difference. The maximum per cent organic carbon 2.02 at 0-15 cm found in Selimbong while 1.45 at 15-30 cm and 1.42 at 30-45 cm depths was found in Soureni. The minimum per cent organic carbon was recorded in Selimbong with 1.46 at 0-15 cm, 0.74 at 15-30 and 0.63 at 30-45 cm depths respectively (Table 3). The high per-centage organic carbon present in the surface soils may be due to the undecomposed and partially decomposed organic matter and lower values seen in the sub-surface soils may be due to decomposed organic matter which has undergone chemical and biological changes already (Rai *et al.* 2021). These findings were similar to that of Arya, (2014). Table 4 shows the available nitrogen in  $\text{kg ha}^{-1}$  of soil at 0-15, 15-30 and 30-45 cm depth respectively where it was found to be significant at 5% critical difference. The maximum available nitrogen of  $696.3 \text{ kg ha}^{-1}$  at 0-15 and  $340.7 \text{ kg ha}^{-1}$  at 15-30 cm respectively was recorded in Soureni, while  $209.73 \text{ kg ha}^{-1}$  at the 30-45 cm depth was recorded in Selimbong. The minimum available nitrogen of  $441.3 \text{ kg ha}^{-1}$  at 0-15 cm depth was found in Selimbong, while  $212.7 \text{ kg ha}^{-1}$  was found at 15-30 cm and  $177.70 \text{ kg ha}^{-1}$  at the 30-45 cm depth at Glenburn. The high nitrogen content may be due to leaf litter or application of organic amendments in the soil. It may also be because of the high organic matter content in soil and addition of chemical fertilizers by the tea growing farmers (Pratistha *et al.*, 2020). These findings were similar with that of Mishra *et al.*, (2000). The available phosphorus of soil at 0-15, 15-30 and 30-45 cm depths respectively, was found to be significant at 5% critical difference. The maximum available phosphorus of  $29.54 \text{ kg ha}^{-1}$  at 0-15,  $27.79 \text{ kg ha}^{-1}$  at 15-30 and  $25.02 \text{ kg ha}^{-1}$  at 30-45 cm depths was found in Glenburn. The minimum available phosphorus was found in Soureni, with  $17.23 \text{ kg ha}^{-1}$  at 0-15,  $16.55 \text{ kg ha}^{-1}$  at 15-30 and  $16.13 \text{ kg ha}^{-1}$  at 30-45 cm depth (Table 4). The available phosphorus content may be due to the influence of soil pH which is acidic in nature and often limits nutrients that remain present in plant nuclei and act as an energy storage (Kekane *et al.*, 2015). These findings were similar with that of Ram *et al.*, (2014). Table 4 also shows the available potassium of soil at 0-15, 15-30 and 30-45 cm depths respectively was found to be significant at 5% critical difference. The maximum available potassium was  $334.92 \text{ kg ha}^{-1}$  at 0-15,  $322.32 \text{ kg ha}^{-1}$  at 15-30 and  $320.32 \text{ kg ha}^{-1}$  at 30-45 cm depths respectively and it was found at Selimbong. The minimum available potassium was found in Glenburn, with  $101.43 \text{ kg ha}^{-1}$  at 0-15,  $99.90 \text{ kg ha}^{-1}$  at 15-30 and  $97.43 \text{ kg ha}^{-1}$  at 30-45 cm depths respectively. The high proportion of potassium may be due to the feldspar and muscovite mica found in the Darjeeling region (Laitpharlang Cajee, 2018). These findings were similar with that of Ram *et al.*, (2014). The results in figure 3 show that the exchangeable calcium of soil at 0-15, 15-30 and 30-45 cm depths was found significant

at 5% critical difference, with the maximum exchangeable calcium  $0.42 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 0-15,  $0.39 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 15-30 and  $0.38 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 30-45 cm depths respectively found at Selimbong. The minimum exchangeable calcium was found at Soureni, with values of  $0.03 \text{ C mol (p}^+) \text{ kg}^{-1}$  at both 0-15 and 15-30 cm depths, and  $0.02 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 30-45 cm depth. Very low values of calcium were recorded which may be due to the leaching of calcium as hydrogen is added to the soil by decomposition of organic matter as well as due to heavy rainfall (Pratistha *et al.*, 2020). These findings were similar with that of Ray *et al.*, (2012). The exchangeable magnesium of soil at 0-15, 15-30 and 30-45 cm depths was found significant at 5% critical difference, the maximum exchangeable magnesium  $0.22 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 0-15,  $0.20 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 15-30 and  $0.19 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 30-45 cm depth found in Selimbong. In the case of Soureni, the minimum exchangeable magnesium was  $0.02 \text{ C mol (p}^+) \text{ kg}^{-1}$  at the 0-15 as well as the 15-30 cm depths while it was  $0.01 \text{ C mol (p}^+) \text{ kg}^{-1}$  at 30-45 cm depth (Figure 3). Very low magnesium was recorded in all sites which may be due to heavy rainfall prevailing in the area and surface leaching of basic cations ( $\text{Mg}^{+2}$ ) by the excessive rainfall, as also reported by Ray *et al.*, (2012).

## CONCLUSION

It is evident that the Darjeeling district soils are in good physical condition, which is favourable for tea cultivation. The soil colour varied from brown to yellowish brown in dry condition and dark brown to dark reddish brown in wet condition. The soils of Selimbong were sandy loam to sandy clay loam in texture, soils of Soureni were clay loam in texture, and soils of Glenburn were clay loam to sandy clay loam in texture. The bulk density and particle density of soil was found to be in low levels ranging from  $1.22 \text{ Mg m}^{-3}$  to  $1.01 \text{ Mg m}^{-3}$  and  $2.64 \text{ Mg m}^{-3}$  to  $2.20 \text{ Mg m}^{-3}$  respectively. The high clay component of the soil was discovered to contribute to its good pore space and water-holding capacity. Low pH and EC values have been recorded. Mostly all the villages showed high organic carbon content as well as high available nitrogen concentration which may be due to the adoption of organic farming in the tea gardens. The soils had a medium to high range of phosphate and potassium, with values decreasing as depth increased. Very low values of exchangeable calcium and magnesium was found in all the regions which may be due to heavy rainfall prevailing in the area and surface leaching of basic cations ( $\text{Ca}^{+2}$  and  $\text{Mg}^{+2}$ ) by the excessive rainfall. The main aim of the study was to determine the physical and chemical properties of the soils in the selected tea gardens. Not only would

this be beneficial to make practical decisions in adopting farming techniques and incorporate fertilizers but also add to the database on soils in these areas for reference for further scientific research. The findings suggested that implementing excellent agricultural practises, such as the use of organic amendments, herbicides, and micronutrients, could help to ensure the sustainability of tea crop yield and quality. However, applying additional labile organic inputs, liming materials, and appropriate inorganic fertilizers (N-P-K) would be useful for long-term management and boosting soil fertility in the tea-growing region.

## **ACKNOWLEDGEMENT**

I sincerely acknowledge my advisor Dr. Ram Bharose, the Head of Department Prof. (Dr.) Tarence Thomas for their guidance, support and expertise and the Department of Soil Science and Agriculture Chemistry, Sam Higginbottom University of Agricultural Science and Technology, Prayagraj, U.P. for providing necessary support and desired equipment's for this research work.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

**Table 1:** Soil colour of different villages in dry and wet condition of tea garden soils of Darjeeling district at 0-15 cm, 15-30 cm and 30-45 cm depths.

Village	Dry condition			Wet condition		
	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm
<b>Pokhribung</b>	7.5YR 4/4	7.5YR 5/4	7.5YR 5/4	5YR 3/3	7.5YR 5/4	7.5YR 4/4
<b>Khasmahal (V1)</b>	Dark Brown	Brown	Brown	Dark Reddish Brown	Brown	Dark Brown
<b>Soolbongs Tea Garden (V2)</b>	7.5YR 5/4 Brown	7.5YR 6/3 Pale Brown	7.5YR 6/3 Pale Brown	10YR 5/4 Yellowish Brown	10YR 6/4 Light Yellowish Brown	10YR 5/4 Yellowish Brown
<b>Selimbong (V3)</b>	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown	7.5YR 5/4 Brown	2.5YR 4/4 Reddish Brown	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown
<b>Phuguri Forest (V4)</b>	10YR 5/4 Yellowish Brown	10YR 6/6 Brownish Yellow	7.5YR 6/3 Pale Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown
<b>Sourinibasti (V5)</b>	10YR 3/4 Dark Yellowish Brown	10YR 5/4 Yellowish Brown	10YR 8/6 Pale Yellow	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown	7.5YR 5/4 Brown
<b>Singbulli Tea Garden(V6)</b>	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	10YR 6/6 Brownish Yellow	7.5YR 5/4 Brown	7.5YR 6/3 Pale Brown	7.5YR 4/4 Dark Yellow
<b>Singringtam (V7)</b>	7.5YR 5/4 Brown	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown
<b>Kambal Tea Garden (V8)</b>	10YR 5/4 Yellowish Brown	10YR 5/4 Yellowish Brown	7.5YR 5/4 Brown	7.5YR 4/4 Dark Brown	5YR 3/4 Dark Reddish Brown	7.5YR 4/4 Dark Brown
<b>Manedara (V9)</b>	7.5YR 5/4 Brown	10YR 5/4 Yellowish Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown	7.5YR 4/4 Dark Brown	7.5YR 5/4 Brown

**Table 2:** Soil texture of different villages of tea garden soils of Darjeeling district at 0-15cm, 15-30 cm and 30-45 cm depths.

Tea Garden	Village	Sand %	Silt %	Clay %	Textural Class
Selimbong	V1	64.3	13.9	21.8	Sandy Clay Loam
	V2	67.0	17.8	15.2	Sandy Loam
	V3	61.5	22.0	16.5	Sandy Loam
	V4	26.2	39.2	34.6	Clay Loam
Soureni	V5	29.4	37.5	33.1	Clay Loam
	V6	23.9	37.1	39	Clay Loam
	V7	48.9	21.9	29.2	Clay Loam
Glenburn	V8	45.9	29.3	24.8	Clay Loam
	V9	53.8	26.1	21.1	Sandy Clay Loam

**Table 3:** Pore space, water holding capacity and percent organic carbon of soil in different villages of Darjeeling at 0-15 cm, 15-30 cm and 30-45 cm depths


Villages		Pore space			Water holding capacity			Organic carbon		
		0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm
Selimbong	V1	57.16	54.73	50.53	69.70	65.06	68.13	1.46	0.74	0.71
	V2	54.63	57.83	56.14	71.03	53.36	63.16	2.02	1.02	0.63
	V3	55.50	54.36	52.13	69.93	70.53	68.03	1.76	1.02	0.86
	V4	53.13	52.30	56.73	71.96	71.16	70.69	1.87	1.30	1.25
Soureni	V5	56.43	55.63	55.33	79.43	78.83	70.86	1.81	1.45	1.42
	V6	57.66	55.40	59.86	74.13	70.05	72.66	1.71	1.27	1.27
	V7	52.96	50.30	52.06	70.56	68.68	68.40	1.59	1.44	1.31
Glenburn	V8	68.96	62.67	54.36	65.96	63.86	66.25	1.51	1.27	1.03
	V9	58.73	62.96	53.23	61.01	62.51	60.23	1.53	1.15	1.09
F-test		S	S	S	S	S	S	s	s	s
S. Em. (±)		4.84	4.28	2.87	5.27	7.08	3.74	0.18	0.22	0.27
C.D @5%		1.24	4.39	4.4	1.92	0.57	8.73	3.10	1.41	1.59

**Table 4:** Available potassium, available nitrogen and available phosphorus of soil in different villages of Darjeeling at 0-15 cm, 15-30 cm and 30-45 cm depths respectively.



Villages	Available potassium	Available nitrogen	Available phosphorus
----------	---------------------	--------------------	----------------------

		0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm
Selimbong	V1	314.44	306.67	302.16	484.0	268.5	209.73	25.35	25.02	24.35
	V2	334.92	322.32	320.32	441.3	241.5	194.56	20.12	19.90	19.25
	V3	278.84	273.00	268.47	442.9	279.6	193.86	17.86	17.29	16.13
	V4	125.76	124.02	120.88	612.1	282.7	189.90	18.72	17.28	16.89
Soureni	V5	110.92	106.10	100.64	679.4	314.2	189.03	24.03	23.43	22.84
	V6	149.90	145.31	141.08	696.3	340.7	182.30	17.23	16.55	16.20
	V7	219.46	214.37	210.83	543.5	217.5	183.16	28.79	27.79	25.02
Glenburn	V8	143.84	138.48	136.76	553.8	212.7	185.76	24.26	22.87	21.73
	V9	101.43	99.90	97.43	574.8	275.4	177.70	29.54	27.50	24.13
F-test		s	s	s	s	s	s	s	s	s
S. Em. ( $\pm$ )		91.29	88.54	88.53	93.30	41.96	9.35	4.61	4.39	3.65
C.D @5%		5.63	9.58	5.99	1.10	2.81	3.16	0.001	0.008	0.138



**Fig 1:** Graphical representation of bulk density ( $\text{Mg m}^{-3}$ ) and particle density ( $\text{Mg m}^{-3}$ ) of soil in different villages of tea garden soil of Darjeeling at 0-15 cm, 15-30 cm and 30-45 cm depths respectively.



**Fig 2:** Graphical representation of soil pH and EC ( $\text{dSm}^{-1}$ ) at  $25^\circ\text{C}$  of soil in different villages of tea garden soil of Darjeeling at 0-15cm, 15-30 cm and 30-45cm depth



**Fig 3:** Graphical representation of exchangeable calcium ( $C \text{ mol}(p^+) \text{ kg}^{-1}$ ) and exchangeable magnesium ( $C \text{ mol}(p^+) \text{ kg}^{-1}$ ) of soil in different villages of tea garden soil of Darjeeling at 0-15 cm, 15-30 cm and 30-45 cm depths respectively.



## References

1. Ahamed, Z. (2021) A study on Soil Problems in North Bengal, West Bengal, India. *Mazedan Chemical Research Journal*; 2(1), 1-3.

2. Arya, M.K. (2014) Assessment of physico-chemical properties of soil along altitudinal gradients in a protected forest in the Kumaun Himalayas, India. *Nature and Science*; 12(2): 32-37.
3. Barthwal, A., Swaroop, N., Rao, P. S., & Thomas, T. (2019) Assessment of physical properties of soil in Dehradun district, Uttarakhand, India. *International Journal of Chemical Studies*; 7(3): 1623-1625.
4. Bisen, J.S. and Singh A.K. (2012) Impact of inorganic to organic cultivation practices on yield of tea in Darjeeling hills – A case study. *Indian Journal of Horticulture*; 69(2): 288-291
5. Black, C.A. (1965) *Methods of soil analysis*. American Society of Agronomy, Madison, Wisconsin, USA, 2.
6. Bouyoucos, G. J. (1927) The hydrometer as a new method for the mechanical analysis of soils. *Soil Science*; 23: 343-353. region, Sikkim, India.
7. Cajee L., (2018) Physical Aspects of the Darjeeling Himalaya: Understanding from a Geographical Perspective. *IOSR Journal of Humanities And Social Science (IOSR-JHSS)*; 23(3): pp 66-79.
8. Deb, P., Debnath, P., Denis, A.F and Lepcha, O.T. (2018) Variability of soil physico-chemical properties at different agro-ecological zones of Himalayan region, Sikkim, India. *Environmental Development and Sustainability*, Springer science + Business Media B.V, part of Springer Nature 2018.
9. Fisher, R. A. (1960) *The design of experiments*. Seventh edition. Hafner publishing company, New York.
10. Ganorker, R.P., Hole, H.A and Pund, D.A (2017) Assessment of soil nutrients and physico-chemical parameters in the region of Hiwarkhed village of Amaravati district, Maharashtra, India. *Rasayan J Chem*; 10(2): 429-433.
11. Jackson, M.L. (1927) *Soil Chemical analysis*. Prentice hall of India Pvt. Limited, New Delhi ;38:226.
12. Kekane S.S., Chavan R.P., Shinde D.N., Patil C.L., and Sagar S.S. (2015) A review on physico-chemical properties of soil. *International Journal of Chemical studies*; 3(4): pp 29-32.
13. Laitpharlang Cajee, (2018) Physical aspects of the Darjeeling Himalaya: Understanding from a Geographical Perspective. *IOSR Journal of Humanities And Social Science (IOSR-JHSS)*; 23(3): pp 66-79
14. Majumdar K, Ray DP, Chakraborty S, Pandit T. (2014) Change of nutrient status of hilly soil in Darjeeling district within five years. *International Journal of Bioresource Science*; 1(1):25-30.
15. Mishra, T. K., Nanda, A.K., Mandal, P, and Saha, A. (2017) Physiochemical Properties of Soils under Different Tea Growing Regions of North Bengal: A study from 2006 to 2010. *International Journal of Research in Chemistry and Environment*; 8(1), pp 44-48
16. Munsell, A. H. (1954) *Munsell Soil Color Charts*. Munsell Color Company Inc., Baltimore.
17. Muthuvel, P., Udayasoorian, C., Natesan, R., & Ramaswami, P. R. (1992) *Introduction to Soil Analysis*. Tamil Nadu Agricultural University, Coimbatore.
18. Nayak, R.K., Sahu, G.C and Nanda, S.S.K (2002) Characterization and classification of soils of Central Research Station, Bhubaneswar. *Agropedology*; 12:1-8.

19. Olsen, S.R., Cole, C.V., Watanable, F.S and Dean, L.A (1954) Estimation of available phosphorus in soil by extraction with sodium bicarbonate, USDA, Circular; 939: 1-19.
20. Pramanik M.K (2016) Site suitability analysis for agricultural land use of Darjeeling district using AHP and GIS techniques. *Modeling Earth Systems and Environment*; 2(2)
21. Pratistha, P., Swaroop, N., David, A. A., & Thomas, T. (2020) Assessment of physical properties of soil in Kalimpong district, West Bengal, India. *International Journal of Chemical Studies*; 8(4): 1718-1721.
22. Rai, A. and Thomas, T. (2021) Assessment of Physical Properties of Soils of Darjeeling District, West Bengal, India. *Biological Forum – An International Journal*; 13 (2). pp 481- 487.
23. Ram R, Chatterjee S, Maji C, Sharma PK, Rani P. (2016) Effect of soil health, nutrient management and soil test-based doses of lime on mulberry leaf yield (*Morus alba* L.) in acid soils of Kalimpong hills. *International Journal of Agriculture Sciences*; 8(59):3333-3337.
24. Ray, S.K and Mukhopadhyay, D. (2012) A Study on Physicochemical Properties of Soils under different tea growing regions of West Bengal (India). *International Journal of Agriculture Sciences*; 4(8), pp 325-329.
25. Shirsath, W.B (2021) Quantitatively Physico-Chemical Analysis of Some Soil Samples of Satana (Baglan)Tahsil, District Nashik, Maharashtra (India). *Journal of Scientific Research*; 65(7), pp 143-148.
26. Subbiah, B.V and Asija, G.L. (1956) A rapid procedure for the estimation of available nitrogen in soils. *Current Science*; 25: 259-260.
27. Toth, S.J and Prince, A.L. (1949) Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by Flame Photometer method. *Soil Science*; 67:437445.
28. Walkley, A and Black, C.A (1934) An estimation for different methods for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*; 37:29-38.
29. Wankhade, R. R. (2015) Physico-chemical analysis of soil from some farms of Digras region of district Yavatmal in Maharashtra, India. *International Journal of Multidisciplinary & Allied Studies*; 2(10): 1-4.
30. Wilcox, L.V. (1950) Electrical Conductivity. *American Water Work Association Journal*; 42:775-776.
31. Muzib S, Rahman MH, Haque HA, Tanu FZ, Hakim A. Quality of Tea Soil Induced by Cultivation Period. *Asian Soil Res. J.* [Internet]. 2023 Sep. 9 [cited 2024 Jun. 10];7(3):30-42. Available from: <https://journalasrj.com/index.php/ASRJ/article/view/134>
32. Bhuyan K, Kalita R, Saikia GK, Chandra Barua S, Borah N. Effect of Gas Flaring on Physico-Chemical Properties of Soil under Tea Plantation in Assam, India. *Int. J. Plant Soil Sci.* [Internet]. 2022 Mar. 9 [cited 2024 Jun. 10];34(7):16-23. Available from: <https://journalijpss.com/index.php/IJPSS/article/view/1706>
33. Karak T, Paul RK, Boruah RK, Sonar I, Bordoloi B, Dutta AK, Borkotoky B. Major soil chemical properties of the major tea-growing areas in India. *Pedosphere*. 2015 Apr 1;25(2):316-28.
34. Baruah BK, Das B, Medhi C, Misra AK. Fertility status of soil in the tea garden belts of golaghat district, Assam, India. *Journal of Chemistry*. 2013;2013(1):983297.