

Bioprospecting of Indigenous Microbial Species for Bioethanol Production from the Fruits of Some Selected Exotic Ornamental Plants

ABSTRACT

The study determined the ability of some selected indigenous microbial isolates, namely *Bacillus subtilis*, *Zymomonas mobilis* and *Saccharomyces cerevisiae*, in the production of bioethanol from the fruits of some exotic plants (i.e. *Ixora coccinea*, *Duranta repens* and *Syzygium guineense*). *B. subtilis* was isolated from soil, while *Z. mobilis* and *S. cerevisiae* were isolated from palm wine using cultural methods. The organisms were identified based on their colony characteristics and morphological features, which were observed by microscopic examination of the stained smears of the isolates. The identity of the organisms was confirmed through the molecular study of the 16S rRNA and 18S rRNA regions of the microbial genomes using Polymerase Chain Reaction – Denaturing Gradient Gel Electrophoresis (PCR - DGGE). The fruit samples were collected, washed, blended, filtered using a clean piece of cloth and pretreated using ultrasonication and autoclave. Concentrations of reducing sugars in the fruit filtrates were determined using DNSA-UV-visible spectrophotometry. Seven (7) conical flasks (500ml capacity) were arranged for each fruit sample in duplicate, and 250ml of the fruit filtrates were dispensed in the flasks accordingly. The fermenting organisms were inoculated in the flasks containing the fruit filtrates singly and in combination. Flasks were incubated in shaking incubator for seven days at 35°C. After incubation, the concentrations of bioethanol produced were determined using the UV-visible spectrophotometry. Highest reducing sugars concentrations of 43.7% and 43.6% were found in *I. coccinea* and *S. guineense*, respectively, while 36.4% was recorded for *D. repens*. Highest bioethanol yield of 27.5m/L was found in *I. coccinea* fermented by *S. cerevisiae* and *Z. mobilis*, while the lowest yield of 17.25m/L was found *D. repens*. The yield in bioethanol production was higher in samples containing a combination of fermenting organisms compared to ones with individual cells. It was recommended that the fruits of these exotic plants should be considered for commercial production of bioethanol.

Keywords: Indigenous microbial species, Exotic plants, Fruits, Bioethanol, Fermentation.

Introduction

The incessant global population growth, coupled with industrialization, is increasing the energy demands all over the world. Biomass-derived fuels are known to provide promising alternative; they are renewable, non-toxic and their combustion do not release carbon dioxide (CO₂) into the atmosphere. It is estimated that biomass will contribute up to 50% of the world's primary energy consumption by the year 2050 (Kumar, 2010); it is one of the world's largest sustainable energy source (Kour *et al.*, 2019). The global community has acknowledged bioethanol for providing energy security worldwide. This has to do with the fact that bioethanol has a great potential of becoming a future fuel of choice due to its renewable nature and efficiency. Bioethanol is a liquid

biofuel that it can be produced from the microbial fermentation of various biodegradable feedstock, including fruits (Kour *et al.*, 2019; Dandasena and Shahi, 2023). Of great interest in the present study are the fruits of some exotic ornamental plants such as scarlet jungle flame (*Ixora coccinea*), golden newdrop (*Duranta repens*) and malmo (*Syzygium guineense*). The spread of these plants in our residential and institutional areas is increasing; they produce fruits that cause various environmental inconveniences if not properly managed.

Microorganisms ferment a vast array of carbohydrates from fruits to produce bioethanol. The use of microorganisms to produce biofuels provide a double environmental benefit in addition to reducing greenhouse gases emissions (Zuluaga-Mazo *et al.*, 2020). The production processes also reduce environmental waste by harnessing the ability of various microorganisms to utilize and break down complex carbohydrate into fermentable sugars (El-Imam, 2012; Alam *et al.*, 2022). Microorganisms varied in their abilities to degrade sugars; microbial species have some types of sugars they can hydrolyze, not necessarily all; thus, different organisms are needed in fermentation medium with a variety of sugars, so that absolute degradation of these substrates can be fully attained to achieve highest bioethanol yield (Al-Makishah, 2017). These organisms can be isolated from their various habitats and identified; their abilities to produce bioethanol from different sugars can be determined (Sawicka *et al.*, 2022). This study determined the bioethanol production ability of some indigenous microbial isolates using the fruits of some selected exotic plants, namely *I. coccinea*, *D. repens* and *S. guineense*.

MATERIALS AND METHODS

Isolation of Microorganisms

The organisms isolated and used in the study were *Bacillus subtilis*, *Zymomonas mobilis* and *Saccharomyces cerevisiae*. *B. subtilis* was isolated from garden soil samples, while *Z. mobilis* and *S. cerevisiae* were isolated from palm wine according to the method employed by Amin *et al.* (2015). Ten (10) garden soil samples and 10 palm wine samples were collected from different locations in clean polythene bags and plastic bottles, respectively. One milligram (1g) of the soil sample was dissolved in 10 ml of nutrient broth and incubated at 35°C for 24 hours. After incubation, 1ml of the cultured broth was streaked on fresh nutrient agar and incubated under the same conditions as above. In the same manner, one milliliter (1ml) of wine sample was introduced into 10 ml of nutrient broth and incubated anaerobically at 35°C for 24 hours. After incubation, 1ml from the broth culture was streaked on fresh nutrient agar and incubated anaerobically to isolate *Z. mobilis*. To isolate *S. cerevisiae*, 1ml of the wine sample was aseptically transferred into 10 ml of Sabouraud Dextrose Broth and incubated at 35°C for 48 hours, and after incubation, 1ml was streaked on fresh Sabouraud Dextrose Agar (SDA) and incubated under the same conditions (Sukirtha and Abinaya, 2019).

Identification of the Microbial Isolates

Colony characteristics were recorded for all the isolates to facilitate identification (Caprette, 2022). The colonies expected to be the target organisms were sub-cultured on fresh media to produce the axenic culture of the individual cells. These were used for microscopic examinations and molecular analysis to confirm their identities. For microscopy, a portion of the colony from the

axenic culture was collected using a wire loop and smeared on clean glass slide. Both the bacterial and yeast isolates were Gram-stained to examine their Gram reactions and shapes (Pryshchepa *et al.*, 2022).

The molecular analysis was carried out by genomic study of the 16S rRNA and 18S rRNA regions of the bacterial and fungal (yeast) DNAs, respectively. The microbial DNAs were extracted using Nucleo-pore gDNA Fungal/Bacterial Mini Kit (Genetix Biotech Asia Manual, 2018), amplified using universal primers for bacteria (27F/1492R) and fungi (NS1/NS4), and identified using BLAST programme, as described in Alhassan *et al.* (2020).

Pretreatment of Fruit Samples

The clean fruit samples were separately blended with a blender (Preethi Taurus, India) to accelerate microbial degradation and enhance fermentation. A duplicated setups of seven (7) conical flasks (500ml-capacity) were arranged for each fruit and a known quantity (20g) of the blended fruit sample was accordingly placed in each flasks and diluted with 250ml of distilled water. The conical flasks were covered appropriately with cotton wool and aluminum foil, and then placed in ultrasonicator (SW 30H, Nickel-Electro Ltd., Switzerland) to enhance pretreatment. The samples were then sterilized in an autoclave (STV-I, Bioevopeak, China) at a pressurized temperature of 121°C for 15 minutes.

Bioethanol Production

The production of bioethanol from the pretreated fruit samples was conducted through fermentation as described in Alhassan (2023). Each sample in the first setup was inoculated with 25ml of the suspension of a pure broth culture of *Saccharomyces cerevisiae*; while the second was inoculated with that of *B. subtilis*, the third with *Z. mobilis*, the 4th with a mixed culture of *S. cerevisiae* and *B. subtilis*, the 5th with a mixed culture of *S. cerevisiae* and *Z. mobilis*, the 6th with a mixed culture of *B. subtilis* and *Z. mobilis*, and the last (7th) was inoculated with a mixed culture of *S. cerevisiae*, *Z. mobilis* and *B. subtilis*. The flasks were incubated in shaking incubator (SI – 200, Major Science Co. Ltd., Singapore) at 45°C for seven (7) days (Ziemlewska *et al.*, 2021), and bioethanol produced was recovered from the fermentation media using distillation (Alhassan, 2023).

Estimation of Reducing Sugars and Bioethanol

Reducing sugars concentrations in the pretreated fruit samples and bioethanol yield were determined using UV-spectrophotometry as described in Alhassan (2023).

RESULTS AND DISCUSSION

Results

Results of Microbial Isolation: Table 1 below shows the result of the microbial isolation from the soil and palm wine samples. *B. subtilis* was isolated in four out of the 10 soil samples investigated. *Z. mobilis* was isolated from two out of 10 samples, while *S. cerevisiae* was isolated

from six (6) out of 10 samples. The identification of these organisms was confirmed by molecular study of the 16S and 18S RNA examination.

Table 1: Results of the Microbial Isolation and Identification

Organism	Yield	Cultural features	Morphological features	Molecular confirmation
<i>B. subtilis</i>	40%	White rounded colonies with flat and dry surfaces.	Short chains Gram positive large rod-shaped cells	Confirmed
<i>Z. mobilis</i>	20%	Whitish convex colonies	Gram negative rod-shaped cells	Confirmed
<i>S. cerevisiea</i>	60%	Smooth creamy colonies with circular margin	Circular budding yeast cells	Confirmed

Results of Reducing Sugars Determination: The results of the reducing sugars determination after pretreatment is presented in Figure 1 below. It was found that *I. coccinea* had the highest reducing sugars concentration (28.6%) with significant difference ($p < 0.05$) compared to *S. guineese* (23.2%) and *D. repens* (with 5.3%).




Figure 1: The Result of the Reducing Sugars Concentrations in the Fruit Samples

Results of Bioethanol Yield: Figure 2 presents the results of the bioethanol concentrations, where it was observed that *I. coccinea* had the highest yield of bioethanol with significant difference ($p > 0.05$) when compared to *S. guineense* and *D. repens*, but there was no significant difference regarding the fermentation ability of the microbial isolates. Moreover, there was no significant difference in bioethanol yield observed between the single culture and the mixed culture of the microbial isolates.




Figure 2: The Result of bioethanol yield from the fermentation of fruit samples using various combinations of fermenting organisms

Discussion

Microorganisms are diverse in their ability to degrade organic matter to generate various important products such as bioethanol. Microorganisms can be found in different environment, including soil, fermented beverages and fruits, among others (Sukirtha and Abinaya, 2019). *S. cerevisiae* and *Z. mobilis* are usually isolated from alcoholic beverages, and are extensively used in the production of bioethanol. The isolation of these organisms from palm wine in the present study was in line with previous studies, including Frances *et al.* (2023) and Migap *et al.* (2023), where the organisms were isolated from this source to produce bioethanol. Some species of bacteria such as *Bacillus* and *Clostridium* are usually isolated from soil samples and they are capable of producing a variety of alcohols such as ethanol and butanol. The isolation of *B. subtilis* from soil samples in this study was in line with previous studies, including the work of Nithya and Kumar (2022), where the organism was isolated from garden soil to produce ethanol. *S. cerevisiae*, *Z. mobilis* and *B. subtilis* isolated in this study have been described as promising organisms in the production of bioethanol.

The proliferation of exotic plants is on the increase in Kano and other Northern states in Nigeria. These plants produce a considerable amount of fruits that are usually wasted; their consumption is not common. Of particular interest in this study are *I. coccinea*, *D. repens* and *S. guineense*; their

fruits are described to have high sugar content (Jordaan and Downs, 2012). Considering the results of fermentation using the identified organisms in this study, the utilization of these fruits in the production of bioethanol will have great economic and sustainable impacts.

Ultrasonication has been used in the recent times to enhance pretreatment in bioethanol production processes. The use of ultrasonication in the pretreatment of the fruit samples was found to be very effective, as it increased the yield of reducing sugars in the samples, higher than what was found in Alhassan (2023), in which autoclave was employed as the only pretreatment method. Similar findings were reported in Subhedar and Gogate (2016), where ultrasonication was found to enhance pretreatment of various biomass in bioethanol production processes.

With regards to bioethanol yield, the mixed cultures containing two fermenting organisms produced the higher ethanol concentrations compared to ones produced by a single isolate. Moreover, the mixed culture containing the three organisms was found to have the highest yield, though the differences in each case were not statistically significant (at $p > 0.05$). This finding is similar to that of Chibuzor *et al.* (2016) and Na-Allah *et al.* (2021), where a combination of the fermenting organisms was found to produce higher bioethanol yield compared to individual microbial cells. The overall bioethanol yield in this study was in agreement with that of Alhassan (2023).

CONCLUSION

The fruits of exotic plants, specifically *I. coccinea*, *D. repens* and *S. guineense*, are produced in large quality every season, and usually found everywhere in the environment around cities of developing countries, including Kano (Nigeria) without proper management and utilization. These fruits can be fermented by a variety of microorganisms including yeast (such as *S. cerevisiae*) and bacteria (such as *B. subtilis* and *Z. mobilis*) to produce bioethanol. If commercial production of this fuel and other important bioproducts will be considered from these fruits, it will have a great socioeconomic and environmental impact, especially in developing countries.

References

- Alam, A., Bibi, F., Deshwal, K. (2022). Biofortification of primary crops to eliminate latent hunger: an overview. *Natural Resources for Human Health*, 2(1): 91 – 99. <https://doi.org/10.53365/nrfhh/142883>.
- Alhassan A.S. (2023). Assessment of Fruits of Some Exotic Plants in the Production of Bioethanol using *Saccharomyces cerevisiae*. *Bayero Journal of Pure and Applied Sciences*, 14(1): 416 – 420.
- Alhassan, A.S., Yahaya, S., Musa, H., Mohammed, K. and Mukhtar, M.D. (2020). Utilization of Rumen Content by Indigenous Microorganisms in a Modified Anaerobic Digester. *Energy Research Journal*, Volume 11: 12.21.
- Al-Makishah, N.H. (2017). Bioenergy: Microbial Biofuel Production Advancement. *International Journal of Pharmaceutical Research & Allied Sciences*, 6(3):93-106
- Amin, M., Rakhisi, Z. Ahmady, A.Z. (2015). Isolation and Identification of *Bacillus* Species from Soil and Evaluation of Their Antibacterial Properties. *Avicenna J Clin Microb Infec.*, 2(1): e23233.
- Caprette, D.R. (2022). Describing Colony Morphology [Internet]. Available from: <https://bit.ly/324cqkA>
- Chibuzor, O., Uyoh, E.A and Igile, G. (2016). Bioethanol production from cassava peels using different microbial inoculants. *African Journal of Biotechnology*, 15(30), pp. 1608-1612.
- Dandasena, T.K. and Shahi, S. (2023). A Renewable Biofuel-Bioethanol: A Review. *Journal of Advanced Zoology* 44(S3):1698-1706. DOI: 10.17762/jaz.v44iS3.2388.
- EL-Imam, A M. (2012). Microbial Biofuels: Fast, Present and Future Trends: A Review. *International Journal of Phytofuels and Allied Sciences*, 1(1), p. 130.
- Frances, E.C., Enoch, N.N., Johnson, O.O., Eziamaka, A.E.C., and Ann, M.O. (2023). Isolation and Characterization of Yeast Associated with Palm Wine Fermentation. *Asian Journal of Food Research and Nutrition*, 2(3), 117 – 127.
- Jordaan, L.A. and Downs, C.T. (2012). Nutritional and Morphological Traits of Invasive and Exotic Fleshy-fruits in South Africa. *Biotropica* 0(0): 1–6.
- Kour, D., Rana, K.L., Yadav, N., Yadav, A.N., Rastegari, A.A., Singh, C., Negi, P., Singh, K. and Saxena, A.K. (2019). Technologies for Biofuel Production: Current Development, Challenges, and Future Prospects (Chapter 1), *Prospects of Renewable Bioprocessing in Future Energy Systems, Biofuel and Biorefinery Technologies*, Vol. 10, Springer Nature Switzerland AG.
- Kumar A, Kumar K, Kaushik N, Sharma S, Mishra S (2010) Renewable energy in India: current status and future potentials. *Renew Sust Energy Rev* 14:2434–2442.

Migap, H.H., Abdullahi, I.O., Whong, C.M.Z., Benschak, J.A., Jurbe, F.D., Adamu, R.R., Dalhatu, H.A. and Uzoma, S. (2023). Isolation and molecular characterization of *Zymomonas mobilise* for bioethanol production from palm saps in some parts of Jos, Plateau state, Nigeria. *International Journal of Bioscience and Biochemistry*, 5(1): 37-41

Na-Allah, A.J., Iliyasu, M.Y., Haruna, U.S., Ahmad, A., Oguche, S.O., Karamba, K.I., Abba, M. and Agbo, E.B. (2021). Sustained production of bioethanol by *Zymomonas mobilis* and *Saccharomyces cerevisiae* using Rice Husk and Groundnut Shell as Substrates. *Microbiology Research Journal International*, 31(9): 15-26.

Nithya, K.V. and PrasannaKumar, C. (2022). Isolation and Characterization of *Bacillus subtilis* from the Garden Soil. National College, Tiruchirappali. <https://doi.org/10.13140/RG.2.2.16986.62404>

Pryshchepa, O., Złoch, M., Pomastowski, P., Railean-Plugaru, V., Rodzik, A., Szultka-Młyńska, M. and Buszewski, B. (2022). Modern Approaches for Microorganisms' Identification. In: Buszewski, B., Baranowska, I. (eds) *Handbook of Bioanalytics*. Springer, Cham. https://doi.org/10.1007/978-3-030-95660-8_40

Subhedar, P.B. and Gogate, P.R. (2016). Use of Ultrasound for Pretreatment of Biomass and Subsequent Hydrolysis and Fermentation (Chapter 6): Biomass Fractionation Technologies for a Lignocellulosic Feedstock Based Biorefinery, Elsevier. P. 127-149.

Sawicka, B., Pszczółkowski, P., Barbaś, P., Skiba, D. and Bienia, B. (2022). The Role and Importance of Microorganisms in Environmental Sustainability (Chapter 7), Wiley Online Library. <https://doi.org/10.1002/9781119834489.ch7>

Sukirtha, K. and Abinaya, K. (2019). Isolation and Identification of Microorganisms from Palm Wine for Ethanol Production. *International Journal of Pharmacy and Biological Sciences*, 9(2): 74-81

Zuluaga-Mazo, C., Arango-Bermudez, D., Salas-Zapata, W.A. (2020). Profile of the Use of Microorganisms within Environmental Management: Systematic Review 2012-2017. *Gestión y Ambiente*, 23(1). DOI: <https://doi.org/10.15446/ga.v23n1.83464>