

**Original Research Article**  
**Effect of salinity and substrate on the  
emergence and growth of propagules of the  
mangrove species *Rhizophora racemosa* in the  
Sassandra-Dagbego Ramsar complex (Côte  
d'Ivoire)**

---

**ABSTRACT**

This study aimed to evaluate the behaviour of seeded propagules of *Rhizophora racemosa* on different substrates and under different salinity levels. Three substrates including sand, mud and a mixture of the two were tested together with three salinity levels (low 5%, moderate 10% and high 25%). *R. racemosa* seedlings were more likely to emerge and grow under moderate and low salinity conditions. The propagules had significant early growth in mud compared to sand and mud-sand mixture. The combined effect of salt and substrate influenced significantly propagules performance in nursery ( $P < 0.001$ ). High propagules emergence and growth was observed in the combination mud and salt treatments as compared to sand and sand-mud mixture substrates. These results provide valuable information for the management and restoration of mangroves, highlighting the optimal environmental conditions for the successful regeneration of this species.

*Keywords:* Climate resilience, *Rhizophora racemosa*, Mangrove restoration, Early growth, Côte d'Ivoire

**1. INTRODUCTION**

Mangroves are primarily found in tropical and subtropical coastal areas, usually in sheltered estuaries and bays where wind and wave energy are lower [1,2]. They provide various ecosystem functions and benefits, such as carbon sequestration, storm and wave buffering, and the preservation of marine and coastal biodiversity [3-5]. Despite the ecological importance of mangrove forests, their global coverage has been steadily declining. In Côte d'Ivoire, efforts have been made to restore mangroves, albeit minimally [6], by planting hundreds of hectares of degraded mangroves using direct seeding techniques and transplanting seedlings from nurseries [7]. As mangroves are easy to rehabilitate, restoration efforts often involve hasty or belated plantations of inappropriate species in unsuitable locations [8]. These activities are frequently coupled with poor post-planting monitoring [9] and lack of active involvement of local communities [7]. Consequently, these endeavors often result in failures characterized by high mortality rates and stunted plant growth [10], suggesting effective strategy be deployed. The design of effective conservation and management plans usually requires knowledge of the system's functioning [11]. Beyond the inefficacy of conservation measures, the failure of rehabilitation or restoration projects necessitates scientific investigations into the system's functioning and species' physiological responses to key stress factors, including salinity [9]. Although mangrove forests are typically located in sheltered coasts with tranquil environments, the establishment and early root development of seedlings are sensitive to the physical composition of sediments in


mangroves [12]. Recent research suggests that environmental properties (soil and water) influence mangrove vegetation, specific composition, and structure [13]. According to Sinsinet *et al.*, [14], variations in NaCl also affect mangrove seedling growth. Salinity is one of the most critical factors influencing mangrove health, distribution, growth, and productivity [15]. It often exhibits strong spatial and temporal fluctuations caused by freshwater and seawater inputs, flooding, groundwater infiltration, and evaporation [16-18]. Salinity was reported to determine the survival and growth of planted mangrove seedlings in restoration projects across various sites [19-22]. However, these factors are rarely considered in mangrove regeneration initiatives and their subsequent monitoring and evaluation operations. In Côte d'Ivoire, numerous studies have previously highlighted the species richness, anthropogenic pressures leading to mangrove degradation [23,7], mangrove rehabilitation, and the ecosystem services they provide to surrounding populations [7]. However, little is known about how substrate and salinity affect the emergence and growth of mangrove seedlings. Yet, this knowledge is crucial for a sound decision-making and a better monitoring of nurseries in mangrove restoration program. Therefore, the present study focusing on *R. racemosa* nursery production was initiated to assess the behavior of propagules sown on different substrates and subjected to varying salinity levels. More specifically, the study aimed to (i) identify the appropriate substrate type for the emergence and early growth of *R. racemosa* propagules and (ii) evaluate the effect of different salinity levels on the emergence and early growth of *R. racemosa* propagules and (iii) determine the combined effect of substrate type and salinity level on the emergence and early growth of *R. racemosa* propagules.

## 2. MATERIAL AND METHODS

### 2.1 Materials

#### 2.1.1 Study site

This study was carried out in Niéga, located in the town of Sassandra and part of the Sassandra-Dagbego Ramsar complex (Fig. 1). Niéga has sandy soil, although in mangrove stands it is mainly silty-clay soil. Salinity varies greatly in the river depending on the geographical location and the measurement season. The tropical climate with average monthly temperatures ranging from 23 to 28°C [24] and an average annual rainfall of about 1,600 mm [25] is characterized by four seasons: two rainy seasons and two dry seasons. The main rainy season is from April to July. The short rainy season is from October to November. The main dry season runs from December to March and the small from August to September. Two of the three mangrove species found in the country are present : *Rizophoraracemosa* (red mangrove), *Avicennia germinans* (white mangrove).



## **2.2Methods**

### **2.2.1 Seedling Production**

The nursery was set up near the lagoon at low tide close to brackish water. The experiment was conducted in a semi-controlled environment for better management of predators and watering [26,14]. The fruiting peak of *R. racemosa* species in the region varies between November and March. Given this, propagules were collected for this study in February. All propagules were collected on the tree according to different criteria used by Sinsin *et al.*, (2021) [14]. The length ranged from (12 cm to 40 cm), circumference (3 cm), colour (brown or green) and the presence of a bud. Mature propagules that are still attached to the tree are identified because they drop off easily on contact. Three salinity levels were tested : high salinity (sea water), moderate (lagoon water) and low (mixture lagoon water - drilling water). The high level (H) corresponds to 25% salinity against 10% and 5% respectively for the moderate level (M) and low level (L). Propagules were planted in polyethylene pots filled with one of three types of substrates all taken from the vicinity of the lagoon: sand-mud mixture (Mi), sand (S), mud (Mu), according to the volumetric ratio of 1:1 (MI). One propagule was planted directly in the substrate of each pot.

### **2.2.2Experimental Design, Sampling, Measurement**

The experiment was set in completely randomized and based on a 3 x 3 factors, three salinity levels and three substrate types with 3 repetitions. The nursery spanned 8 m<sup>2</sup>, and we utilized 270 polyethylene pots (13 cm in diameter and 24 cm in height). We employed 90 pots for each repetition, totaling 30 pots per treatment. Spacing of 50 cm between repetitions and 43 cm between treatments were observed.

### **2.2.3Statistical Analysis**

The collected data were analyzed using the RStudio software. All data were tested for normality. Descriptive statistics were performed and histograms were drawn to illustrate the evolution of emergences and growth parameters as a function of time, treatments and factors. The differences in emergence between the three salinity levels and between the three substrates were explored by a non-parametric alternative to the single-factor ANOVA test (the Kruskal-Wallis test). Dunn test was done to identify different groups. The leaf number, stem height variables were compared between the fixed salinity and substrate factors using a two-way ANOVA. The outputs are presented as means  $\pm$  standard deviations, while differences between treatments were considered significant at the 5% ( $p < 0.05$ ) level.


## **3. RESULTS AND DISCUSSION**

### **3.1Results**


#### **3.1.1 Emergence of propagules following the substrate type**

For the different substrate types, emergence rates were substantially equal, showing that the emergence of propagules was not significantly influenced by the substrate type. From the 30 days observation period, the rate of emergence of propagules was higher in mud, followed by mud-sand mixture and sand (Fig. 2). The emergence on sand-mud mixture and mud started simultaneously earlier (7th day) to reach respectively a higher rate of 69 and 86%,

respectively at 28 and 25th days. The emergence of sand seeded propagules was low and reached its maximum (51%) just at 21th day (Fig. 3).




**Fig. 2. Emergence of the seedlings of *R. racemosa* with different substrate**




**Fig.3. Evolution of emergence rates of the seedlings of *R. racemosa* with different substrate**

### 3.1.2 Emergence of propagules as a function of salinity

After 30 days of observation, the rate of emergence of propagules had varied with the treatments though no significant between treatments . The emergence of propagules was lower in high salinity treatment (Fig. 4). The emergence of high salinity propagules began late and a low rate (34%) was achieved. On the other hand, moderate and low salinity levels started a little earlier (6th and 9th days) to reach respectively a higher rate of 82 and 89%, respectively (Fig. 5).



**Fig. 4. Emergence of the seedlings of *R. racemosa* with different salinities**



**Fig. 5. Evolution of emergence rates of the seedlings of *R. racemosa* with different salinities**

### 3.1.3 Combined effect of salinity and substrate on propagule emergence

The emergence of propagules in the sand substrate reached its maximum, 73 and 77% for low salinity (5%) and moderate salinity (10%), respectively. High salinity gave a very low emergence rate (3%). The emergence of the propagules began on the 7th and 10th day for moderate and low salinity levels, respectively. On the other hand, it started only after 14th day for the high salinity level (Fig. 6a). Similarly, for the mud-sand mixture, the lowest emergence rate was recorded for high salinity (25%). The emergence rates varied and reached its maximum between 80-100% for the moderate and low salinity levels. The emergence was earlier and began on the 6th day for the moderate salinity and 9th day for low salinity. The emergence rate did not change from the 21st day for the low salinity level and from 24th day for the moderate salinity level (Fig. 6b). Moreover, the mud substrate presented remarkable rates of emergence irrespective of the levels of salinity. The emergence of propagules from low and moderate salinity levels began earlier (6th day). They reached 90% compared to 73% for the high salinity level that began 7 days later (Fig. 6c). Therefore, sand and mixture sand - mud, are not suitable for the emergence of propagules with high salinity. These substrates are suitable only for moderate and low salinity levels. Conversely, the mud was suitable for all three salinity levels.




Fig. 6a. Evolution of the emergence of *R. racemosa* propagules in sand substrates following different salinity levels for 30 days





Fig. 6b. Evolution of the emergence of *R. racemosa* propagules in mixture substrates following different salinity levels for 30 days



**Fig. 6c. Evolution of the emergence of *R. racemosa* propagules in mud substrates following different salinity levels for 30 days**


**3.1.4 Effect of treatments on propagule height growth**

Significant variations were observed for *R. racemosa* seedlings height regardless of the type of treatments (Table 1). The smallest increase in height were recorded for treatments (HS, HMi, HM) between 0 and 1.80 cm at 5 weeks. The highest increase in height were observed for treatments (LMu and MS) with 9 and 8 cm achieved at 5 weeks (Fig. 7). Thus, a treatment consisting of low salinity with mud as substrate is more suitable for *R. racemosa* seedling growth at an early stage of its life cycle.

**Table 1. Results of the ANOVA for effect of duration and treatments on seedling height**

Source of variation	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Weeks	1	103.23	103,23	982,46	< 2e-16 ***
Treatments	8	218.66	27.33	260.12	< 2e-16 ***
Weeks : Treatments	8	41.83	5.23	49.77	3.55e-14 ***
Residuals	27	2.84	0.11		
---					
Signif. codes	0 '****'	0.001 '***'	0.01 '**'	0.05 '.'	0.1 ' ' 1

**Height-Times**




**Fig. 7. Height growth of *R. racemosa* for 5 weeks after propagule emergence**

### 3.1.5 Effect of treatments on propagule height growth

There were significant variations for seedling height among the different treatments (Table 2). High numbers of leaves (Table 2) were observed for treatments LS, LMi, LMu, MS, MMi, MMu (up to 4 leaves). Although having given less leaves, the appearance of the leaves under the treatment HMu, HS happened at the same time as the others. Only HMi treatment showed late onset of leaves (Fig. 8)

**Table 2. Results of the ANOVA for effect of duration and treatments on seedling height**

Source of variation	Df	Sum Sq	Mean Sq	F value	Pr(>F)
<b>Weeks</b>	1	19.6	19.600	63.00	1.57e-08 ***
<b>Treatments</b>	8	30.4	3.800	12.21	3.36e-07 ***
<b>Weeks :</b>	8	5.6	0.700	2.25	0.0551
<b>Treatments</b>					
<b>Residuals</b>	27	8.4	0.311		
---					
<b>Signif. codes</b>	0 (****)	0.001 (***)	0.01 (**)	0.05 (.)	0.1 ( ' ) 1



**Fig. 8. Number of leaves of *R. racemosa* for 5 weeks after propagule emergence**

### 3.1.5 Effect of treatments on propagule height growth

Emergence of propagules of *R. racemosa* reduced at high salinities of 25%. Growth (first 30 days after emergence) is possible only for low to moderate (5–10%) salinities. Seawater salinities of 25% restrict the emergence and growth of seedlings of *R. racemosa*. However, a salinity of 5% is adequate for nursing because it enhances both emergence and growth. Mud is the best substrate whatever the salinity. He gives best emergence and growth. The best treatment for seedling is the combination of low salinity and mud. If substrate salinity is low or high where the seedlings will be transplanted later, water of low or moderate salinity would be suitable in nurseries to avoid the risk of dieback from osmotic shock after

transplantation. The results may be useful to non-governmental organizations (NGOs), governments and institutions engaged in mangrove restoration through plantations.

### 3.1 Discussion

Plant seedling growth can be influenced by the environment conditions. In this study, we observed that highest emergence rate was achieved on the mud and mud-sand mixture as compared to sand substrate. These results are consistent with Clough (2011) [27], who obtained higher survival rates in the mangrove soil for *Aviceniagerminans* and *Rhizophora mangle* as compared to sand substrate. Similarly, Abrahão (1998) [28] found higher emergence rate (up to 82%) of seedlings of *A. schauerian* grown in nursery using a mixture of sand and compost fertilizer (N, P, K). This suggests that mixture of composite soil may be more suitable for emergence of mangrove plant species seedling in nursery than sandy soil. Furthermore, our results indicated that with the mud substrate, the emergence rate was very high even under high level of salinity. This can be explained by the ability of the mud to retain excess salt in irrigation water thus facilitate the emergence and good growth of seedlings. Similar to the Sinsin *et al.*, (2021) [14], the best emergence rates were obtained with low and moderate salinity levels, confirming that the application of salt nutrient within this range can speed up propagules emergence in *R. racemosa* seedling production. Salinity influences seedling emergence through its effect on water and osmotic potential [29-31]. Previous studies have demonstrated that high salt concentration decreased emergence and growth rates in *Aegicerascorniculatum* (L.) Blanco, *Allophylus cobbe* (L.) Reusch, *Pemphisacidula* J. R. Forst. & G. Forst, and *Sonneratiacaseolaris* (L.) Engl. *Rhizophora* species are considered to be less tolerant to salt than *Avicennia* [27,30]. Previous studies on *Rhizophora* showed that propagule emergence is also inversely correlated with salinity [32,20,33]. Our results showed that emergence is very low and almost inhibited under high salinity level (25%). This could be explained by the very low osmotic potential of propagules compared to that of water [27,34]. The optimal salinity range observed (5-10%) is not identical to that reported for other *Rhizophora* species. Biber (2006) [32], found that *R. apiculata*, *R. stylosa* and *R. mangle* all had optimal growth at 15% and Khan and Aziz (2001) [18] at 17%. Osmoregulation is generally considered the most important adaptation against higher salinity of media [34]. Salt tolerance mechanisms may vary among haplotypes and provenances. This suggests further studies to establish the interaction effect of provenances and salinity levels on the emergence of *R. racemosa* propagule. Differences were also noted between the height and the number of leaves following the water salinity levels. Reduced growth occurred at higher salinities with significantly lower plant height and leaf production, which can be interpreted as the result of an investment in tolerance mechanisms [16,35]. Thus, adequate salinity for the early stages of the seedling establishment in nursery life cycle remains uncertain. For many mangrove species, regardless of salinity tolerance, low salinity is recommended for the first five months. Kodikara *et al.*, (2018) [36] suggested that moderate salinity could improve mangrove plantations after the first 4-5 months of low salinity. Lugo *et al.*, (2007) [37] reported in Puerto Rico, in hypersaline lagoons, that tree height was proportional to salinity, resulting in stunted *R. mangle* trees.

### 4. CONCLUSION

As part of the restoration of mangrove ecosystems, this study investigated the effect of salinity and substrate on the emergence and growth of propagules of the mangrove species *R. racemosa*. The results showed that these environmental factors play an important role in the development of seedlings. In terms of salinity, we found that *R. racemosa* propagules are more likely to emerge under low to moderate salinity (5-10%) conditions. The propagules of *R. racemosa* had a significant early growth in mud compared to sand and mud-sand mixture. The combination of the two factors gave a complex interaction between salinity and

substrate. High salinity levels, associated with different substrates, were found to result in low but remarkable emergence and growth rates for mud.

## REFERENCES

1. Friess DA, Rogers K, Lovelock CE, Krauss KW, Hamilton SE, Lee SY, Lucas R, Primavera J, Rajkaran A, Shi S. The State of the World's Mangrove Forests: Past, Present, and Future. *Annual Review of Environment and Resources*. 2019;44:89–115.
2. Romañach SS, De Angelis DL, Koh HL, Li Y, Teh SY, Raja Barizan RS, Zhai L. Conservation and restoration of mangroves: Global status, perspectives, and prognosis. *Ocean and Coastal Management*. 2018;154:72–82.
3. Alongi DM. Mangrove forests: Resilience, protection from tsunamis, and responses to global climate change. *Estuarine, Coastal and Shelf Science*. 2008;76(1):1–13.
4. Alongi DM. Carbon cycling and storage in mangrove forests. *Annual Review of Marine Science*. 2014;6:195–219.
5. Lee SY, Primavera JH, Dahdouh-guebas F, Mckee K, Bosire JO, Cannicci S, Diele K, Fromard F, Koedam N, Marchand C, Mendelssohn I, Mukherjee N, Record S. Ecological role and services of tropical mangrove ecosystems: A reassessment. *Global Ecology and Biogeography*. 2014;23(7):726–743.
6. . Allassane O, Cecch P. Vulnerability of coastal and estuarine societies and environments in West Africa. In L'harmattan-Senegal. (Ed.), *Status report and conservation of mangroves in Ivory Coast*; 2021. English
7. . Egnankou MW. Rehabilitation of mangroves between Fresco and Grand-Lahou in Ivory Coast: Important areas for fishing. *Nature & Wildlife*. 2009;24:85–93. English
8. RCM (Réseau Camerounais des Mangroves). *Les Mangroves*. Matanda News; 2015 9,13. French
9. Kodikara KAS, Mukherjee N, Jayatissa LP, Dahdouh-guebas F, Koedam N. Have mangrove restoration projects worked? An in-depth study in Sri Lanka. *Restoration Ecology*. 2017;25(5):705–716.
10. . Tazo Fopi RD, Ngankam Tchamba M, Nwutih Ajonina G. Physico-chemical characterization and dendrometry in mangrove regeneration treatments in the Cameroon Estuary. *International Journal of Biological and Chemical Sciences*. 2021;15:2701-2714. English
11. Maxted N, Kell SP. Establishment of a global network for the in-situ conservation of crop wild relatives: status and needs. *FAO Commission on Genetic Resources for Food and Agriculture*. 2009;1–266.
12. Jiang J, Zhou C, An S, Yang H, Guan B, Cai Y. Sediment type, population density and their combined effect greatly change the short-time growth of two common submerged macrophytes. *Ecological Engineering*. 2008;34(2):79–90.

13. Hossain MD, NuruddinAA. Soil and Mangrove: A review. *Journal of Environmental Sciences and Technologies*. 2016;9:198-207.DOI: 10.3923/jest.2016.198.207
14. Sinsin CBL, SalakoKV, Fandohan AB, ZanvoMGS, Kouassi KE, GlèlèKakaï, RL. Pattern of seedling emergence and early growth in *Avicenniagerminans* and *Rhizophora racemosa* along an experimental salinity gradient. *African Journal of Ecology*. 2021;59(4):1013–1022.
15. Ball MC. Interactive effects of salinity and irradiance on growth: implications for mangrove forest structure along salinity gradients. 2002:126–139.
16. NaidooG. Relations and on the accumulation of solutes in three. 1985;22:133–143.
17. Aziz I, Ajmal Khan M. Effect of seawater on the growth, ion content and water potential of *Rhizophora mucronata* Lam. *Journal of Plant Research*. 2001;114(3):369–373.
18. Khan MA, Aziz I. Salinity tolerance in some mangrove species from Pakistan. *Wetlands Ecology and Management*. 2001;9:229–233.
19. Bosire JO, Dahdouh-guebas F, Walton M, CronaBI. Author's personal copy Functionality of restored mangroves: A review. *Aquatic Botany Journal*. 2008;89:251–259.
20. Hoppe-speerSCL, AdamsJB, Rajkaran A, Bailey D. The response of the red mangrove *Rhizophora mucronata* Lam. to salinity and inundation in South Africa. 2011;95: 71–76.
21. Kirui BYK, Huxham M, Kairo J, Skov AEM. Influence of species richness and environmental context on early survival of replanted mangroves at Gazi bay, Kenya. 2008;171–181.
22. Krauss KW, Lovelock CE, Mckee KL, Lo L, Ewe SML, Sousa WP. Environmental drivers in mangrove establishment and early development: A review. 2008;89:105–127.
23. de Almeida JPN, Lessa BF, da T, Pinheiro CL, Gomes FM, Filho SM, Silva CC. Germination and development of *amburanacearensis* seedlings as a function of seedweight, light and temperature. *Acta Scientiarum – Agronomy*. 2017;39(4):525–533. <https://doi.org/10.4025/actasciagron.v39i4.32786>. Portuguese
24. . Kouassi H, N'go A, Zro FGB. Spatio-temporal dynamics of natural habitats of the Ramsar site of the Sassandra-Dagbego complex by remote sensing. *International Journal of Innovation and Scientific Research*. 2020;27:243–249. English
25. . Marie-Claude AKL. Spatio-temporal monitoring of the mangroves of the Sassandra-Dagbego complex. *Ivorian Review of Geography of the Savanes*. 2021;10:1–20. English
26. RavishankarT, Ramasubramanian R. *Manual on Mangrove Nursery Techniques*. M.S. Swaminathan Research Foundation, Chennai; 2004
27. Clough BF. Primary productivity and growth of mangrove forests, in: Robertson A.I. & Alongi D.M. (Eds.), *Coastal and Estuarine Studies*. American Geophysical Union, USA (Washington). 1992.

28. Abrahão RG. A new landscape design in engineering works in coastal regions with native mangrove plants (Via Expressa Sul Ilha de Santa Catarina-Brazil). MSc Dissertation, Federal University of Santa Catarina, Florianópolis; 1998. Portuguese
29. Hadas A. Water uptake and germination of leguminous seeds under changing external water potential in osmotic solutions. *Journal of Experimental Botany*. 1976;27(3):480–489.
30. Méndez-Alonzo R, López-Portillo J, Moctezuma C, Bartlett MK, Sack L. Osmotic and hydraulic adjustment of mangrove saplings to extreme salinity. *Tree physiology*. 2016;36(12):1562–1572.
31. Robertson AI, Alongi DM. *Tropical mangrove ecosystems*. American Geophysical Union, Washington, DC; 1992.
32. Biber PD. Measuring the effects of salinity stress in the red mangrove, *Rhizophora mangle* L. *African Journal of Agricultural Research*. 2006;1(1)
33. Smith SM, Snedaker SC. Salinity Responses in Two Populations of Viviparous *Rhizophora mangle* L. Seedlings. *Biotropica*. 1995;27:435.
34. Jefferies RL. The role of organic solutes in osmoregulation in halophytic higher plants. pp.135-154. In: *Genetic Engineering of Osmoregulation: Impact on Plant Productivity for Food, Chemicals and Energy*. (Eds. A. Hollaender, D.W. Rajjins and R.C. Valentine). Plenum Press, New York; 1980.
35. Suárez N, Medina E. Salinity effect on plant growth and leaf demography of the mangrove, *Avicennia germinans* L. *Trees*. 2005;19:721–727.
36. Kodikara A, Kodikara S, Jayatissa LP, Huxham M, Dahdouh-guebas F, Koedam N. The effects of salinity on growth and survival of mangrove seedlings changes with age. *Acta Botanica Brasilica*. 2018;32:37–46.
37. Lugo A, Medina E, Cuevas E, Cintrón G, Laboy-Nieves EN, Schaffer-Novelli Y. Ecophysiology of a fringe mangrove forest in Jobos Bay, Puerto Rico. *Caribbean Journal of Sciences*. 2007;43:200-219.