

A critical review on Millets (Superfood of all Time): Importance, challenges and opportunities

Abstract:

Climate change, water scarcity, population growth, rising food prices, and other socioeconomic repercussions are anticipated to pose a serious danger to agriculture and global food security in the twenty-first century, especially for the world's poorest residents of arid and subarid countries. We must concentrate on dry lands in order to significantly boost the production of grains because agricultural fields with irrigation systems have been fully utilized. It is difficult to use arid areas to produce sufficient quality grains because of their poor productivity. The most important source of food in the world is cereal grains, which also play an important part in the global diet of people. In the semiarid tropics of Africa and Asia, millet is one of the most significant drought-resistant crops and is a major source of carbohydrates and proteins for the local population. Additionally, millet grain is currently attracting more interest from food scientists, technologists, and nutritionists due to its significant contribution to national food security and potential health advantages. The phytochemicals found in millet grains have a beneficial impact on human health by reducing phytates and cholesterol levels. The quest for substitute grains is necessary to relieve the pressure caused by the frenetic demands on cereals and their uses in numerous industries. The performance improved when pearl and finger millets were substituted for maize in the diets of several animals. In terms of poor growth conditions and high nutritional value, millets outperform other grains like wheat and rice as crops that are climate change compliant. These strategies will aid in the fight against hunger and malnutrition while also providing monetary benefits to the millet growers and other stakeholders.

Introduction:

“The world's population is projected to grow to 9.1 billion people by 2050” (FAO, 2009a). “It is obvious that urgent action is required to ensure increased food supply and food security considering the restricted land resources. Food security has grown more important on both the international and domestic fronts as a result of the global supply, economic growth, and rising population in emerging countries. Two of the largest difficulties facing the modern world are addressing hunger and feeding the planet's population. Conflicts, insufficient production of food, imbalances in supply and demand, and shortages in the supply of micro- and macronutrients are some of the causes of this problem causing global instability in a number of places” [91,92,93]. “The concept of food security encompasses both sustained access to wholesome nutrition as well as consistent access to physical food availability. Food insecurity is first divided into two categories by the Food and Agriculture Organization (FAO): moderate (limited access to food results in poorer dietary quality, eating patterns, health, and wellbeing) and severe (running out of food, experiencing hunger, and having health and wellbeing at severe risk) security (Food and Agriculture Organization, 2019). The threat of climate change and global warming persists even while some of these causes of hunger can be addressed, resulting in a marginal decrease in the number of people suffering from hunger and malnutrition from about one billion in 1990–1992 to 850 million in 2010–2012” (FAO,2012). “Estimates indicate that by 2050, 2-3 billion people may experience hunger and other food and nutritional insecurities due to the decline in food production rates and the additional burden of feeding a population greater than 9 billion” (Wheeler and Braun, 2013). “Crop yields, crop production, and the overall sustainability of our food systems are all said to be directly impacted by climate change and rising world average temperatures. At both the macro and micro levels, ensuring food and nutrition security for India's enormous and diverse population is a challenge, and work has been done on numerous fronts. There is no question that significant efforts have been undertaken recently to achieve food and nutrition security in India, and the advancements made cannot be disputed. However, unless concerted efforts are made, food insecurity and malnutrition appear to be continuing and may get worse in the near future. So, the goal of sustainable improvement in food and nutrition security continues to be the main objective. According to some estimates, a few places may gain from climate change because of higher production and yields, but this would not be enough to feed the growing population around the world” (Downing *et al.*, 2000). “Furthermore, the majority of scientists concur that crop output would be severely decreased by the current rates of global warming and greenhouse gas emissions.

Therefore, ensuring food security depends greatly on reducing greenhouse gas emissions in order to control global temperatures. However, one of the main sources of greenhouse gases like methane in the environment is the agriculture industry. Intensive agricultural practices, which are practised in various parts of the world, typically result in higher emissions” (Olesen *et al.*, 2002). “In addition to being a large source of macronutrients including carbohydrates, lipids, and proteins, cereal crops also have a sizable potential to contribute to global warming. Wheat has the biggest global warming potential of all the major cereal crops, with an estimated 4 tonnes CO₂ eq/ha, followed by rice and maize (an estimated 3.4 tonnes CO₂ eq/ha). Additionally, the carbon equivalent emissions from these crops are considerable, with an average of 1000, 956, and 935 kg C/ha for wheat, rice, and maize, respectively” (Jain *et al.*, 2016). “They are widely cultivated and the main sources of nutrition for the entire world's population despite having greater emission rates. Other minor cereal crops, including millet and sorghum, have significantly lower carbon footprints. This is one of the primary reasons why millets could be one of the crops that reduce the global carbon footprint” (Prasad and Staggenborg, 2009). “Improved nutritional security can also be achieved by diversifying the agricultural system and using drought-resistant crops. A class of coarse grain cereals known as millets can be very helpful in achieving this. Millets are a group of diverse small-grain cereal crops grown belonging to the Poaceae grass family and are considered one of the oldest cultivated crops. They are composed of up to a dozen crop species, mostly from developing nations in Asia and Africa” (Gupta *et al.*, 2017). “Ninety-seven percent of millet is produced in developing nations, making millet significant crops in the semiarid tropical regions of Asia and Africa” (Vinoth and Ravidran, 2017). “Despite significant pressure from competing crops in terms of regulations and production support, India is the world's largest producer of millets, generating approximately 40% of the millets consumed worldwide. In temperate, subtropical, and tropical latitudes, the crop is primarily farmed on marginal lands in arid locations. The crop is preferred because of its yield and brief growth season in hot, dry conditions. Throughout human history, millets have played a significant role as a staple meal, especially in Asia and Africa. Since millets are C₄ plants, they have a much more efficient photosynthetic system than C₃ plants like rice and wheat. The majority of millet grains are now commonly referred to as "Nutri-cereals" because they have higher protein, fibre, calcium, and mineral contents than the widely consumed grains of rice and wheat. For the past 10,000 years, East Asia has been cultivating them. Millets are native to many regions of the world, and pearl

millet is the variety that is grown the most widely. It is a significant crop in India and several regions of Africa” (Lu *et al.*, 2009). “The most significant species of millet in terms of cultivated land and contributions to food security in areas of Africa and Asia is pearl millet. The most significant varieties include foxtail, finger, proso, and pearl millet. Pearl millet accounts for roughly half of the world's millet production. In developed nations and some regions of Asia, proso millet is utilized for food and as bird seed. The crop of foxtail millet is significant in China and Eastern Europe” (Singh *et al.*, 2019). In the colder, higher-altitude regions of Africa and Asia, finger millet is widely grown as a food crop and used as a popular ingredient in traditional beer. The other species—barnyard, kodo, and little millets, as well as folios and teff—are regionally or internationally significant food grains. The physical traits, qualitative characteristics, soil, climate requirements, and growth cycles of the various species vary. The scientific and common names of various millets vary depending on the place in which they are grown, and these millets are grown in various parts of the world and require different growth conditions (Table 1). “Every day, millions of people consume rice, wheat, maize, and, to a lesser extent, millet as their primary sources of nourishment. The growth pattern of these crops is determined by temperature and water availability” (Prasad and Staggenborg, 2009). “While wheat is mostly grown in regions that have restricted water supplies and suitable temperatures, rice and maize are grown where there is a plentiful supply of water. Sorghum and millets are cultivated in regions with limited water supplies. Due to its resilience to biotic and abiotic challenges and its large production on low-quality lands with little input, millets can even be grown in semi-arid and arid areas” (Awika, 2011). Millets could be crucial functional foods for the prevention of non-communicable diseases because they are a rich source of antioxidants, fibre, and essential and non-essential amino acids, vitamins, and minerals. This review aims to provide detailed nutritional composition of millet and its benefits to humans and livestock, need of growing millets, various constraints and perspectives associated with it.

Table 1. Scientific names, common names and growing conditions required for cultivation of different types of Millet around the world.

Type of Millets	Scientific Name	Common Name	Growing Conditions	References
Pearl Millet	<i>Pennisetum typhoides</i>	Bulrush millet	Dry climates, marginal soils, Rainfall 200–500 mm	Guigaz and Del'agronome, 2002
Finger Millet	<i>Eleusine coracona</i>	Birds' food millet or African millet	Resist higher temperatures and salt, with a soil pH range of 5 to 8.2 and a moderate amount of rainfall.	Devi <i>et al.</i> , 2014
Porso Millet	<i>Panicum miliaceum</i>	Common millet, hog millet, broom corn, yellow hog, Hershey and white millet	Less water, Rainfall less than 600 mm, average temperature 17 °C during daytime	Zarnkow <i>et al.</i> , 2009
Foxtail Millet	<i>Setaria italica</i>	Italian millet, German millet, or hay millet	Less water, short duration	Vetriventhan <i>et al.</i> , 2012
Barnyard Millet	<i>Echinochloa crusgalli</i> var. <i>Fruventacea</i>		Drought tolerant, rapid maturation rate	Zhang <i>et al.</i> , 2007
Kodo Millet	<i>Paspalum scorbiculatum</i>		High drought resistance, Good yields, period of 80–135 days	Arendt and Dal Bello, 2011

Nutritional Importance: “There are plenty of chronic diseases and health problems in the world today. According to the 2016 Global Nutrition Report, in 129 nations (countries with data available), 44% of the population suffers from very serious levels of undernutrition, adult overweight, and obesity” (IFPRI, 2016). “The majority of these disorders are caused by nutrient imbalances in the diet. United Nations Food and Agriculture Organization estimates that 7.9% of the world's population, or 795 million people, were undernourished in 2015. However, more than 1.9 billion adults under the age of 18 (or 39% of the world's population) were reported to be overweight, and another 13% to be obese” (FAO, 2015). “The World Health Organization has previously classified obesity-related problems like diabetes and cardiovascular disease as an epidemic” (WHO, 2015). “The majority of undernourished people in the world reside in India. Globally, 469,000 deaths were attributed to protein energy malnutrition (PEM), while 84,000 deaths were attributed to a lack of other essential nutrients as iron, iodine, and vitamin A” (Von Grebmer *et al.*, 2017). With a prevalence incidence of 11% for men and 15% for women, obesity is a significant public health issue in India. Because millets are a great source of several essential elements, they offer an extra benefit in the fight against nutrient deficiencies in third-world countries.

Millets contain high amount of calories, protein, and macronutrients when comparable to regular cereals. Their high calorie content, together with calcium, iron, zinc, fats and high-quality proteins, make them an important part of both animal and human diets. They are also rich sources of nutritional fibre and vitamins. Here, is the nutrient composition of millets:

1. **Carbohydrates:** The carbohydrates in pearl millet grains includes dietary fibre, starch and soluble sugars. The endosperm of pearl millet contains glucose in the form of amylase and amylopectin and is regarded as having a high starch content. The starch content of different pearl millet grain genotypes varies from 62.8 to 70.5 % and approximately 71.82 to 81.02 %. Soluble sugars range from 1.2 to 2.6 %, and amylose from 21.9 to 28.8 % (Cheik *et al.*, 2006) and amylose from 21.9 to 28.8 %. The starch included in pearl millet can be employed as bulking, thickening, and gelling agents for food texture (Hadimani *et al.*, 2001). However, according to Bhatt *et al.*, 2003, the total carbohydrate content of finger millet ranges from 72 to 79.5 %. Additionally, Wankhede *et al.*, 1997 (a) noted that the

detailed profile of the carbohydrates ranged between 59.5 and 61.2 % for starch, 6.2-7.2 % for pentosans, 1.4-1.8 % for cellulose, and 0.04-0.6 % for lignin.

2. **Protein:** Protein is the second main component of millet. Pearl millet has 11.6 % protein, which is more than the 7.2 % in rice, 11.5 % in barley, 11.1 % in maize, and 10.4 % in sorghum (Jha *et al.*, 2013). Additionally, Anitha *et al.* (2019) noted that pearl millet contained 9.79 % protein. The crude protein content of pearl millet grain is thought to be 8–60 % higher by weight compared to maize (Burton *et al.*, 1972). Contrarily, finger millet has about 5-8 % protein (Chethan and Malleshi, 2007) recorded the greatest protein level for finger millet at nearly 11 %, and Anitha *et al.* (2019) documented a proportion of 6.32 % in finger millet. Furthermore, Taylor (2004) revealed that the amino acid profile of pearl millet is equivalent to that of wheat, barley and rice but contains more lysine, threonine, methionine and cysteine than sorghum and maize proteins. Moreover, the distribution of proteins in pearl millet grain is thought to be similar to that of maize, notably true prolamins, which are thought to be soluble in alcohol. Additionally, McIntosh *et al.* (2003) noted a high degree of critical amino acid balance and endorsed pearl millet as a crucial source of protein and energy for people. Additionally, arginine, threonine, valine, isoleucine, and leucine were discovered to have greater digestibilities in pearl millet than in maize. Because finger millet contains more lysine, threonine, and valine than other millet, it is somewhat balanced in terms of its concentration of important amino acids (Ravindran, 1991).
3. **Dietary fibre:** Fibre is thought to be beneficial for gut health, and a moderate diet of foods high in fibre may help to promote gut health (McIntosh *et al.*, 2003). Fibre plays a similar role in the prevention of diabetes, colon cancer, and heart disease (Eshak *et al.*, 2010). Bowel movement is enhanced by the pearl millet's high dietary fibre content, which ranges from 8 % to 9 % (Rooney and Miller, 1982). Additionally, due to its slow digestion, it prolongs the transit time, reducing the absorption of glucose into the blood, which benefits non-insulin-dependent diabetic patients. People who eat millet have a lower incidence of diabetes (Shobana *et al.*, 2009). Additionally, the fibre in millet may help lower bad cholesterol while raising good cholesterol. It inhibits the body from producing bile acids, which results in gallstones (Shweta, 2015). In addition, pearl millet's high fibre content

slows the passage of food from the stomach to the intestines. This results in longer gaps between meals, which helps avoid obesity.

4. **Lipids:** The fat content of pearl millet is thought to be between 5 and 7 percent (Gopalan *et al.*, 2003) as compared to 3.21 to 7.71 percent in maize (Ullah *et al.*, 2010). The percentage of lipids in finger and pearl millet was also reported by (Himanshu *et al.*, 2018) to be 1 % and 5 %, respectively. Pearl millet has a high concentration of fatty acids such as palmitic, stearic, and linoleic acids, while maize has a lower percentage of oleic acid (Adeola and Orban, 1995). The overall lipid content of pearl millet grains is larger than that of other millet species, ranging from 1.5 to 6.8 % (Taylor, 2004). The contents of free and bound lipids in pearl millet are, respectively, 5.6 to 6.1 and 0.6 to 0.9 %. Sridhar and Lakshminarayana (1994) reported that the overall lipid content of finger millet, was 5.2% (2.2 % free lipids, 2.4 % bound lipids, and 0.6 % structural lipids). On the other hand, oleic, palmitic, and linoleic acids were found to be the main fatty acids found in finger millet. Unsaturated fatty acids make for 74.4 % of finger millet's total fatty acid production compared to saturated fatty acids' 25.6 % (Kunyanga *et al.*, 2013).
5. **Macronutrients:** Martnez-Ballesta *et al.* (2010), environmental stresses such as high salt concentrations, limited water accessibility, and extreme temperatures have been shown to affect the mineral content of food. The overall mineral and trace element content of pearl millet depends on the type of soil. Ash concentration in pearl millet and maize, respectively, ranges from 1.6 to 3.6 % and 0.86 to 1.35 %. Pearl millet has higher quantities of minerals than maize, including calcium, phosphorus, magnesium, manganese, zinc, iron, and copper (Adeola and Orban, 1995). Due to its high oil content, pearl millet is also regarded as an excellent source of fat-soluble vitamin E (2 mg/100 g). The grain is regarded as an excellent source of vitamin A (Taylor, 2004). Florence *et al.* (2014), pearl millet contains between 45.6 and 48.6 mg of calcium per 100 g. Additionally, it contains a lot of phosphorus, a crucial component found in the mineral matrix of bone, as well as adenosine triphosphate, or ATP, the body's primary energy source. Also, it promotes bone development and healing. According to certain research (Nambiar *et al.*, 2011) the bioavailability of iron may be hampered by phytates, oxalates, and polyphenols, which are found in high concentrations in pearl millet. Contrarily, finger millet has a high calcium content that varies from 162 mg/100 g to 487 mg/100 g depending on the genotype

(Vadivoo *et al.*, 1998). According to a report by Singh Raghuvanshi (2012), finger millet has 344 mg of calcium per 100 g. Additionally, millets contain a significant amount of magnesium, which is thought to help the body fight diseases like cancer. Bachar *et al.* (2013) stated that magnesium content in finger millet ranged from 84.71 mg/100 g to 567.45 mg/100 g. Additionally, finger millet is regarded as a rich source of natural calcium, which promotes bone growth and lowers the incidence of fractures.

Table 2. Nutrient composition of millets in comparison to fine cereals (per 100 g)

Food grain	Carbohydrates (g)	Protein (g)	Fat (g)	Energy (KCal)	Crude fibre (g)	Mineral matter (g)	Calcium (mg)	Phosphorus (mg)	Iron (mg)
Sorghum	72.6	10.4	1.9	349	1.6	1.6	25	222	4.1
Pearl millet	67.5	11.6	5.0	361	1.2	2.3	42	296	8.0
Finger millet	72.0	7.3	1.3	328	3.6	2.7	344	283	3.9
Little millet	67.0	7.7	4.7	341	7.6	1.5	17	220	9.3
Kodo millet	65.9	8.3	1.4	309	9.0	2.6	27	188	0.5
Foxtail millet	60.9	12.3	4.3	331	8.0	3.3	31	290	2.8
Barnyard millet	65.5	6.2	2.2	307	9.8	4.4	20	280	5.0
Proso millet	70.4	12.5	1.1	341	2.2	1.9	14	206	0.8
Wheat (whole)	71.2	11.8	1.5	346	1.2	1.5	41	306	5.3
Rice (raw, milled)	78.2	6.8	0.5	345	0.2	0.6	10	160	0.7

(Hariprasanna *et al.*, 2014)

Polyphenols: Primary polyphenols, which are plentiful in millet and comprise phenolic acids and tannins, are thought to fortify the body's immune system (Chandrasekara *et al.*, 2010). Millet phenolics can partially block the enzymatic degradation of complex carbohydrates. They also block the actions of pancreatic amylase, malt amylase, and -glucosidase, all of which reduce postprandial hyperglycemia. Similarly, it is believed that the presence of ferulic and p-coumaric acids in whole pearl millet can inhibit the growth of cancer cells (Chandrasekara and Shahidi, 2011). The phenols in millets have been demonstrated by Devi *et al.* (2014) to have a number of

health advantages, such as antiviral, anti-inflammatory, and antioxidant properties. Furthermore, celiac disease is an inherited condition brought on by consuming gluten.

As millets are gluten-free, they reduce the discomfort associated with eating normal cereal grains that contain gluten, which helps to prevent celiac disease (Saleh *et al.*, 2013). Sixty percent of the phenolic acids found in millets are found both in bound and free forms. Hydroxycinnamic acids are the most common phenolics in millets, and they are mostly found in the insoluble-bound portions of phenolic acids (Liang and Liang, 2019). Hydroxycinnamic acid is most commonly found in the form of the antioxidant ferulic acid. Antioxidants are well-known compounds with anti-inflammatory and anti-free radical damage effects on the body (Lobo *et al.*, 2010). Furthermore, it has been found that ferulate dimers, which have significant antioxidant activity, are present in millet grains (Emiola and Rosa, 1981).

Additionally, anthocyanidins, chalcones, amino phenolics, flavanols, flavones, and flavanones are among the flavonoids included in millet grains. Additionally, Dykes and Rooney (2006) state that some millet cultivars may contain proanthocyanidins, also referred to as condensed tannins. The majority of coloured millet varieties contain significant levels of tannin (McDonough *et al.*, 2000). Condensed tannins were found to be primarily responsible for the grain's colour, which was linked to this finding. However, the bioavailability of minerals and proteins may be adversely affected by a high concentration of condensed tannins (Chavan *et al.*, 2001).

Other health benefits: Sireesha *et al.* (2011) reported that the aqueous extract of foxtail millet (*Setaria italica*) has been shown to have anti-hyperglycaemic and anti-lipidemic effects in streptozotocin-induced diabetic rats. The study found that when diabetic rats were given 300 mg of *Setaria italica* seed aqueous extract per kilogramme (kg) of body weight, their blood glucose levels significantly decreased (by 70%) after six hours. They also found that in diabetic treated rats as opposed to diabetic untreated rats, there were higher levels of HDL (high-density lipoproteins) cholesterol and lower levels of triglycerides, total LDL (low-density lipoproteins), and VLDL (very low-density lipoproteins) cholesterol, which is indicative of the hypolipidemic effect of the aqueous extract. Dietary protein from Korean foxtail millet has been shown to improve insulin sensitivity and lipid levels (Choi *et al.*, 2005). This experiment demonstrated a significant drop in insulin levels in rats given foxtail millet. When Lee *et al.* (2010) looked into the impact of

consuming millet on lipid levels and C-reactive protein concentration, they discovered that, contrary to their earlier studies (Choi *et al.*, 2005), hyperlipidemic rats fed foxtail millet had lower levels of triglycerides. Reduced levels of C reactive protein, an indicator of inflammation, were also observed in rats given foxtail millet. There is evidence that the aqueous and ethanolic extracts of Kodo millet reduce fasting blood glucose in a dose-dependent manner (Hegde *et al.*, 2002).

Dual use of millets as food and feed: Millet grains are considered unique crops because they are rich in calcium, dietary fibre, polyphenols, and protein, among other vital elements (Devi *et al.*, 2014). They serve as the main source of food for many Asian and African countries. The majority of the millet produced is consumed by humans, with the remainder going towards making beer, animal feed, and bird feed. Millet is prepared as a porridge with a thick or thin consistency in some parts of Africa, and as couscous in other parts (Obiana, 2003). Research using whole grain or crushed grain fed to chickens verified the usefulness of pearl millet as a feed item for poultry production (Cisse *et al.*, 2016). The primary goal of pearl millet production in Africa and Asia is thought to be grain, with forage serving as a key secondary product utilised for animal feed, construction, and fire (Andrews and Kumar, 1992). While season-specific crops like wheat and rice guarantee food security, all-season millets ensure fodder, nutrition, health, and a sustainable livelihood (Shivananjappa, 2018). Pearl millet grains hold considerable potential as a human food source due to their gluten-free nature and higher nutritious fibre content than rice. Furthermore, compared to traditional cereals like wheat and rye, they include more essential amino acids like leucine, isoleucine, and lysine and the same amount of fat as maize cereal (ICRISAT, 2016). In India, millet is transformed into the dense bread known as dosa, where it is commonly used, by combining millet with other grains. Furthermore, according to Dias-Martins *et al.* (2018), it's used to make couscous, sushi, roti, biscuits, and pizza made without yeast. Another product made from millet is madua, an Indian beverage made with finger millet. Also, Oshikundu, a customary alcoholic or non-alcoholic beverage in Namibia, is made using millet (Kumar *et al.*, 2018).

Monetary benefits of using millet

The cost of millet is believed to be 40% cheaper than that of maize, making it a cheap and gluten-free cereal (Gomes *et al.*, 2008). Silva *et al.* (2014) estimated that trade value of pearl millet was less than or equal to 77.78 % of the price of maize grain. Pearl millet grain has more protein per grain than maize, it might allow diets to be formulated without protein supplements, lowering the

price of food and feed in the process. Aside from that, millet is less expensive to produce than other cereals like sorghum and maize. For instance, the water use of Brazilian pearl millet cultivars is more efficient than that of sorghum and maize grown in semi-arid regions of the country (56 ± 2.8 kg DM/ha/mm water vs. 45 ± 1.9 kg DM/ha/mm water for sorghum; (Silva *et al.*, 2011) and 21 ± 2.4 kg DM/ha/mm water for the Brazilian maize cultivars (Dos Santos *et al.*, 2010). For instance, the water use of Brazilian pearl millet cultivars is more efficient than that of sorghum and maize grown in semi-arid regions of the country (56 ± 2.8 kg DM/ha/mm water vs. 45 ± 1.9 kg DM/ha/mm water for sorghum (Silva *et al.*, 2011); and 21 ± 2.4 kg DM/ha/mm water for the Brazilian maize cultivars; (Dos Santos *et al.*, 2010). In an investigation by (Gomes *et al.*, 2008), it was discovered that completely substituting pearl millet for maize in the diet of feedlot cattle was the most cost-effective option. The price of lean and fat cattle, initial weight, final weight, cost of concentrate, cost of roughage, consumption of concentrate, and consumption of roughage were listed as the factors that influenced the financial indicator (Silva *et al.*, 2020). Furthermore, it is reasonable to believe that millet's positioning as a grain that competes with maize will shift the supply's balance and release pressure on maize consumption, thus, lowering its prices. A different study by Rao *et al.* (2002) found that the cost of feed needed to create one kilogram of live weight gain in chickens fed on maize was higher than the cost for the same amount of gain in chickens fed on pearl millet, finger millet, and sorghum. According to Medugu *et al.* (2010), millet grain feed has the lowest cost per kg of feed and the lowest cost of feed per unit weight gain, making broiler chicken production more affordable and cost-effective. Wilson *et al.* (2007) found that the annual net profit from using pearl millet as the only feedstock was \$25,175,000 in comparison to \$23,758,000 for maize feedstocks, a difference of almost \$1.4 million.

Constraints with food security in developing nations: A condition where people have physical and financial access to wholesome food that satisfies their dietary needs is referred to as food security (World Food Summit, 1996). The scope of food security problems is frequently reduced to the availability of agricultural products like cattle (Hatab *et al.*, 2019). The difficulties are thought to be more intricate than simply increasing the supplies. Several elements, including urbanization and accessibility, are among the restrictions Hatab *et al.* (2019) listed. The structure and procedures that govern economies and societies, as well as institutional failings, are frequently mentioned as contributing reasons to food insecurity (Pangaribowo *et al.*, 2013). Fraval *et al.*

(2020) reported that the casual intervention to prevent food insecurity is not always simple, therefore proxies of actions and a deeper comprehension of the various potential pathways are crucial for successful interventions. To overcome these intricate obstacles, multisectoral strategies and planning are needed. Pangaribowo *et al.* (2013) stated that the best strategy to address the problems of food and nutrition security is to combine the indicators of food insecurity with the relevant socioeconomic and environmental variables of a specific entity. Despite the difficulties facing food security in developing nations, the importance of locally grown and underutilized grains like millet cannot be understated. Some of these problems cannot be solved by standard grain crops (Muthamilarasan and Prasad, 2021). The expansion of small grains like millet could worsen poverty among the underprivileged population, especially in light of recent events like the extraordinary COVID-19 epidemic (Muthamilarasan and Prasad, 2021). Millets are crops that, in the context of climatic change, have the potential to endure challenging circumstances and contribute to the stability of food security. Padulosi *et al.* (2009) reported that minor millets like finger, kodo, foxtail, tiny, porso, and barnyard have the capacity to flourish in a variety of soils, different climatic conditions, distinct photoperiods, and in due to their genetic adaptability. These qualities make millets suitable for replacing staples like wheat and rice in difficult climate zones, eventually resulting in food security in these regions. However, millet is regarded as a neglected agro-biodiversity, despite the fact that it has the ability to improve the food security of underprivileged people in developing countries (Garí,2020).

Effect of processing on the antioxidant activity of millet grains:

It was also found that millet grains' antioxidant content and activity were affected by several processing methods, including boiling, soaking, decortication, and malting. According to Rao and Muralikrishna (2002), after 96 hours of malting finger millet, the antioxidant capacity of the fraction containing free phenolic acids increased twofold, whereas the fraction containing bound phenolic acids showed a drop in antioxidant capacity. In a different study, a growth-promoting media was developed to boost the germination of millet and raise the production of a water-soluble protein that inhibits hydroxyl radicals. The single-factor test revealed that H₂O₂ is necessary for inhibitory function, according to Li *et al.* (2007). To further increase the yield of the water-soluble protein that inhibits hydroxyl radicals from stress-germinated millet, the effects of the sprouting circumstances (temperature, duration, and pH of stress medium) on the hydroxyl radical inhibition were also investigated. It was discovered that the stress medium's pH should be 7.5, the culture

should last for 54 hours, and the temperature should be 28 °C. Under optimal conditions, the maximum inhibition of 60.38 percent was reached (Li *et al.*, 2008). Furthermore, researchers were able to create flour with a significant DPPH-scavenging activity after 3 days of foxtail millet germination (Coulibaly and Chen 2011). Roasting or boiling kodo or finger millet reduced its antioxidant activity. Separating the husk and endosperm of kodo millet decreased its DPPH quenching activity as well, and the phytochemicals seemed to complement one another (Hegde and Chandra, 2005). It was also investigated how little millet's antioxidant properties were affected by germination, steaming, and roasting. The results showed that compared to the original sample, the total phenolic, flavonoid, and tannin contents of small millet increased by 21.2, 25.5, and 18.9 mg/100 g, respectively (Pradeep and Guha, 2011).

However, it was found that the phenolic content and antioxidant potential of pearl millet grains were affected by dehulling and hydrothermal treatments (Chandrasekara and Shahidi, 2012). Oxidation and chemical degradation during heat treatments such as boiling, roasting, and frying have resulted in a decrease in antioxidant activity and levels. The reduction that came about as a result of dehulling, however, can be attributed to the removal of the grains' pericarp layer, which is known to be rich in polyphenol and antioxidant compounds. Consequently, in order to maintain their quality and potential health benefits, millet grains, their fractions, and food items need to be processed under optimal conditions. Additionally, the endogenous enzymes that are converted to simpler molecules with enhanced antioxidant activity during germination are responsible for the rise in antioxidant levels and their activities.

Challenges and future perspectives: The information discussed above demonstrates that millet grains were similar to other grains like wheat, rice, and maize despite their high nutritional content and possible health benefits. Additionally, although millet grain has been shown to improve its nutritional value and edibility through processing methods such soaking, fermentation, malting, and fortification/supplementation, its household usage remains mostly limited to rural populations. This is due to the lack of state-of-the-art millet processing methods that can provide large quantities of safe, manageable, ready-to-cook, or ready-to-eat commodities and meals that can be used to feed a large number of people in urban settings (Ushakumari *et al.*, 2004). However, the need for food, fuel, and other requirements rises along with the global population. Therefore, there will be pressure on society to either alter its current crop-consumption patterns or boost agricultural output (Licker *et al.*, 2010). Crop diversification needs to be encouraged at the national and household

levels in addition to higher yields. Offering more conventional and wholesome whole-grain and multigrain substitutes for refined carbohydrates is a vital component of therapeutic dietary modification and promoting the consumption of minor-grain meals (Singh and Raghuvanshi 2012). It is commonly known that gluten protein plays a crucial part in producing excellent yet manageable baked goods and other grain-based foods that require elastic and extensible dough. However, considering that millet grains are gluten-free and based on the outcomes of several laboratory testing, it appears that they are not ideal for making pure-millet bread and other solid food products that are simple to handle. In order to encourage the use of millet grains, it appears that using it as a substitute for wheat in composite flours, complementary foods, and food blends is the best way to create nutrient-dense, "healthy," and secure food products that are of high caliber and can be stored for a long time. Additionally, there is a need for cutting-edge processing technology for decortication, milling, and other preparation procedures of millet grain food in order to make high-quality products on a commercial scale for urban consumers. In return, it is necessary to produce millet cultivars with high essential amino acid content and maintain a steady supply of premium millet grains for industrial needs. Future research studies should assess the nutritional value and potential health advantages of millet grains and their fractions in animal and human models to support efforts to promote their use as food.

Conclusions: There is a lot of potential for millets to increase food and nutritional security. It follows those millets, along with wheat and rice, should be included in the list of staple foods. Even in harsh environments like drought, millets grow well; certain wild types can even flourish in wetlands and inhabited regions. They are rich in minerals (calcium, iron, copper, magnesium, and other elements), B vitamins, antioxidants, and low-glycaemic index. They are also free of gluten. Millets are nutrient-dense and climate change-resistant crops because of these remarkable characteristics. These not only give farmers an extra source of income, but they can also improve the general health of the community. Initiatives for research and development as well as the establishment of policies are therefore necessary; some of them have already been carried out internationally, notably in India, whereas others must be done. Minor millets are highly nutritious and have a simple cultivation technique, but the scientific community needs to pay more attention to them. Scientists refer to them as "orphan cereals" since they are often overlooked. However, novel processing and preparation methods are needed to improve the millet diet's quality and enhance the bioavailability of the micronutrients. More research is needed on the metabolism,

bioavailability, and health advantages of millet grains and all of its constituent parts in people. We'll examine the existing dietary inadequacies in underdeveloped countries in terms of protein, calcium, and iron by the inclusion of millet-based foods in international, national, and state-level feeding programmes.

References:

1. Adeola, O. and Orban, J.I. 1995. Chemical composition and nutrient digestibility of pearl millet (*Pennisetum glaucum*) fed to growing pigs. *Journal of Cereal Science*, **22**(2): 177-184.
2. Andrews, D.J. and Kumar, K.A. 1992. Pearl millet for food, feed, and forage. *Advances in agronomy*, **48**: 89-139.
3. Anitha, S., Govindaraj, M. and Kane-Potaka, J. 2020. Balanced amino acid and higher micronutrients in millets complements legumes for improved human dietary nutrition. *Cereal Chemistry*, **97**(1): 74-84.
4. Arendt, E. and Dal Bello, F. eds., 2011. *Gluten-free cereal products and beverages*. Elsevier.
5. Awika, J.M. 2011 Major cereal grains production and use around the world. In *Advances in Cereal Science: Implications to Food Processing and Health Promotion*; American Chemical Society: Washington, DC, USA, pp. 1–13.
6. Bachar, K., Mansour, E., Khaled, A.B., Abid, M., Haddad, M., Yahya, L.B., El Jarray, N. and Ferchichi, A. 2013. Fiber content and mineral composition of the finger millet of the Oasis of Gabes Tunisia. *Journal of Agricultural Science*, **5**(2): 219.
7. Bhatt, A., Singh, V., Shrotria, P.K. and Baskheti, D.C., 2003. Coarse grains of Uttaranchal: ensuring sustainable food and nutritional security. *Indian farmer's Digest*, **7**(1): 34-38.
8. Burton, G.W., Wallance, A.T. and Radice, K.O., 1972. Chemical composition during maturation and nutritive value of pearl millet. *Crop Science*, **12**(2): 187-188.
9. Chandrasekara, A. and Shahidi, F. 2012. Bioaccessibility and antioxidant potential of millet grain phenolics as affected by simulated in vitro digestion and microbial fermentation. *Journal of functional foods*, **4**(1): 226-237.

10. Chandrasekara, A. and Shahidi, F. 2011. Determination of antioxidant activity in free and hydrolyzed fractions of millet grains and characterization of their phenolic profiles by HPLC-DAD-ESI-MSn. *Journal of Functional Foods*, **3**(3): 144-158.
11. Chavan, U.D., Shahidi, F. and Naczki, M. 2001. Extraction of condensed tannins from beach pea (*Lathyrus maritimus* L.) as affected by different solvents. *Food Chemistry*, **75**(4): 509-512.
12. Chethan, S. and Malleshi, N.G. 2007. Finger millet polyphenols: Optimization of extraction and the effect of pH on their stability. *Food Chemistry*, **105**(2): 862-870.
13. Choi, Y.Y., Osada, K., Ito, Y., Nagasawa, T., Choi, M.R. and Nishizawa, N., 2005. Effects of dietary protein of Korean foxtail millet on plasma adiponectin, HDL-cholesterol, and insulin levels in genetically type 2 diabetic mice. *Bioscience, biotechnology, and biochemistry*, **69**(1): 31-37.
14. Cisse, R.S., Hamburg, J.D., Freeman, M.E. and Davis, A.J. 2017. Using locally produced millet as a feed ingredient for poultry production in Sub-Saharan Africa. *Journal of Applied Poultry Research*, **26**(1): 9-22.
15. Coulibaly, A. and Chen, J. 2011. Evolution of energetic compounds, antioxidant capacity, some vitamins and minerals, phytase and amylase activity during the germination of foxtail millet. *American Journal of Food Technology*, **6**(1): 40-51.
16. Devi, P.B., Vijayabharathi, R., Sathyabama, S., Malleshi, N.G. and Priyadarisini, V.B., 2014. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. *Journal of food science and technology*, **51**: 1021-1040.
17. Dias-Martins, A.M., Pessanha, K.L.F., Pacheco, S., Rodrigues, J.A.S. and Carvalho, C.W.P., 2018. Potential use of pearl millet (*Pennisetum glaucum* (L.) R. Br.) in Brazil: Food security, processing, health benefits and nutritional products. *Food research international*, **109**: 175-186.
18. Dos Santos, R.D., Pereira, L.G.R., Neves, A.L.A., Azevê, J.A.G., de Moraes, S.A. and Costa, C.T.F., 2010. Agronomic characteristics of maize varieties for silage production in the submédio São Francisco river valley. *Acta Scientiarum-Animal Sciences*, **32**(4): 367-373.
19. Downing, T.E., Barrow, E.M., Brooks, R.J., Butterfield, R.E., Carter, T.R., Harrison, P.A., Hulme, M., Olesen, J.E., Porter, J.R., Schellberg, J. 2000. Quantification of uncertainty in

- climate change impact assessment. In *Climate Change, Climatic Variability and Agriculture in Europe*; Environmental Change Institute: Oxford, UK, **pp.** 415–434.
20. Dykes, L. and Rooney, L.W. 2006. Sorghum and millet phenols and antioxidants. *Journal of cereal science*, **44**(3): 236-251.
 21. Emiola, L.O. and De la Rosa, L.C. 1981. Characterization of pearl millet nonstarchy polysaccharides. *Journal of Food Science*, **46**(3): 781-785.
 22. Eshak, E.S., Iso, H., Date, C., Kikuchi, S., Watanabe, Y., Wada, Y., Wakai, K., Tamakoshi. A. Dietary fiber intake is associated with reduced risk of mortality from cardiovascular disease among Japanese men and women. *The Journal of Nutrition*, **140**(8): 1445-1453.
 23. FAO. The State of Food Insecurity in the World 2015. Food and Agricultural Organization of the United Nations; 2015. <http://www.fao.org/world-food-situation/csdb/en/>.
 24. Florence, S.P., Urooj, A., Asha, M.R. and Rajiv, J., 2014. Sensory, physical and nutritional qualities of cookies prepared from pearl millet (*Pennisetum typhoideum*). *Journal of Food Processing & Technology*, **5**(10): 377
 25. Food and Agriculture Organization 2019. The state of food security and nutrition in the world: Safeguarding against economic slowdowns and downturns. Rome, Italy.
 26. Food and Agriculture Organization of the United Nations. Technical Note: FAO Methodology to Estimate the Prevalence of Undernourishment; FAO: Rome, Italy, 2012.
 27. Fraval, S., Yameogo, V., Ayantunde, A., Hammond, J., De Boer, I.J., Oosting, S.J. and Van Wijk, M.T., 2020. Food security in rural Burkina Faso: the importance of consumption of own-farm sourced food versus purchased food. *Agriculture & Food Security*, **9**(1): 1-17.
 28. Garí, J.A., 2002. Review of the African millet diversity. In *International workshop on fonio, food security and livelihood among the rural poor in West Africa*, **pp.** 19-22.
 29. Gomes, P.C., Rodrigues, M.P., Albino, L.F.T., Rostagno, H.S., Gomes, M.F.M., Mello, H.H.D.C. and Brumano, G., 2008. Determination of chemical composition and energy value of millet and their use in rations of broilers from 1 to 21 days of age. *Revista Brasileira de Zootecnia*, **37**: 1617-1621.
 30. Gopalan, C., Rama Sastri, B.V, Balasubramanian, S.C. 2003. Nutritive value of Indian foods. Hyderabad: National Institute of Nutrition.

31. Guigaz, M. Memento Del'agronome; CIRAD-GRET and Ministère des Affaires Étrangères: Montpellier, France, 2002.
32. Hadimani, N.A., Muralikrishna, G., Tharanathan, R.N. and Malleshi, N.G. 2001. Nature of carbohydrates and proteins in three pearl millet varieties varying in processing characteristics and kernel texture. *Journal of Cereal Science*, **33**(1): 17-25.
33. Hariprasanna, K., Gomashe, S., Ganapathy, K.N. and Patil, J.V. 2014. Millets for ensuring nutritional security.
34. Hatab, A., Cavinato, M.E.R. and Lagerkvist, C.J. 2019. Urbanization, livestock systems and food security in developing countries: A systematic review of the literature. *Food Security*, **11**(2): 279-299.
35. Hegde PS, Chandra TS. 2005. ESR spectroscopic study reveals higher free radical quenching potential in kodo millet (*Paspalum scrobiculatum*) compared to other millets. *Food Chemistry*, **92**:177–82.
36. Hegde, P.S. and Chandra, T.S. 2005. ESR spectroscopic study reveals higher free radical quenching potential in kodo millet (*Paspalum scrobiculatum*) compared to other millets. *Food Chemistry*, **92**(1): 177-182.
37. Himanshu, K., Sonawane, S.K. and Arya, S.S. 2018. Nutritional and nutraceutical properties of millets: a review. *Clin. J. Nutr. Diet*, **1**(1): 1-10.
38. ICRISAT. Smart food millet recipes 2016. <http://www.icrisat.org/PDF/Food-Booklet-Millet.pdf>. Accessed 16 May 2020.
39. IFPRI. Global Nutrition Report: Malnutrition Becoming the “New Normal” Across the Globe; 2016.
40. Jain, N., Arora, P., Tomer, R., Mishra, S.V., Bhatia, A., Pathak, H., Chakraborty, D., Kumar, V., Dubey, D.S., Harit, R.C. and Singh, J.P. 2016. Greenhouse gases emission from soils under major crops in Northwest India. *Science of the Total Environment*, **542**: 551-561.
41. Jha, A., Tripathi, A.D., Alam, T. and Yadav, R. 2013. Process optimization for manufacture of pearl millet-based dairy dessert by using response surface methodology (RSM). *Journal of food science and technology*, **50**: 367-373.
42. Kumar, A., Tomer, V., Kaur, A., Kumar, V. and Gupta, K. 2018. Millets: a solution to agrarian and nutritional challenges. *Agriculture & food security*, **7**(1): 1-15.

43. Kunyanga, C.N., Imungi, J.K. and Vellingiri, V. 2013. Nutritional evaluation of indigenous foods with potential food-based solution to alleviate hunger and malnutrition in Kenya. *Journal of Applied Biosciences*, **67**: 5277-5288.
44. Lee, S.H., Chung, I.M., Cha, Y.S. and Park, Y. 2010. Millet consumption decreased serum concentration of triglyceride and C-reactive protein but not oxidative status in hyperlipidemic rats. *Nutrition Research*, **30**(4): 290-296.
45. Li, J., Chen, Z., Yao, H. and Xu, Y. 2007. Optimization of stress medium enhance hydroxyl radical inhibition by water-soluble protein from germinated millet. *LWT-Food Science and Technology*, **40**(9): 1630-1636.
46. Liang, S. and Liang, K. 2019. Millet grain as a candidate antioxidant food resource: a review. *International Journal of Food Properties*, **22**(1): 1652-1661.
47. Licker, R., Johnston, M., Foley, J.A., Barford, C., Kucharik, C.J., Monfreda, C. and Ramankutty, N. 2010. Mind the gap: how do climate and agricultural management explain the 'yield gap' of croplands around the world?. *Global ecology and biogeography*, **19**(6): 769-782.
48. Lobo, V., Patil, A., Phatak, A. and Chandra, N. 2010. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy reviews*, **4**(8): 118-126
49. Lu, H., Zhang, J., Liu, K.B., Wu, N., Li, Y., Zhou, K., Ye, M., Zhang, T., Zhang, H., Yang, X. and Shen, L., 2009. Earliest domestication of common millet (*Panicum miliaceum*) in East Asia extended to 10,000 years ago. *Proceedings of the National Academy of Sciences*, **106**(18): 7367-7372.
50. Martínez-Ballesta, M.C., Dominguez-Perles, R., Moreno, D.A., Muries, B., Alcaraz-López, C., Bastías, E., García-Viguera, C. and Carvajal, M.J.A.F.S.D. 2010. Minerals in plant food: effect of agricultural practices and role in human health. A review. *Agronomy for sustainable development*, **30**(2): 295-309.
51. McDonough, C.M., Rooney, L.W., Serna-Saldivar, S.O. In: Kurl K, Ponte JG Jr, editors. *The millets in handbook of cereal science and Technology, second edition, revised and expanded*, chapter 4. New York: Marcel Dekker Inc; 2000.
52. McIntosh, G.H., Noakes, M., Royle, P.J. and Foster, P.R. 2003. Whole-grain rye and wheat foods and markers of bowel health in overweight middle-aged men. *The American journal of clinical nutrition*, **77**(4): 967-974.

53. Medugu, C.I., Kwari, I.D., Igwebuikwe, J., Nkama, I., Mohammed, I.D. and Hamaker, B. 2010. Performance and economics of production of broiler chickens fed sorghum or millet as a replacement for maize in the semi-arid zone of Nigeria. *Agriculture and Biology Journal of North America*, **1**(3): 321-325.
54. Muthamilarasan, M. and Prasad, M. 2021. Small millets for enduring food security amidst pandemics. *Trends in Plant Science*, **26**(1): 33-40.
55. Nambiar, V.S., Dhaduk, J.J., Sareen, N., Shahu, T. and Desai, R. 2011. Potential functional implications of pearl millet (*Pennisetum glaucum*) in health and disease. *Journal of Applied Pharmaceutical Science*, **10**: 62-67.
56. Obilana, A.B. 2003. Overview: importance of millets in Africa. *World (all cultivated millet species)*, **38**: 28-34
57. Olesen, J.E. and Bindi, M. 2002. Consequences of climate change for European agricultural productivity, land use and policy. *European journal of agronomy*, **16**(4): 239-262.
58. Cheik, A.O., Aly, S., Yaya, B., Alfred, T.S. and et de la Terre, V., 2006. A comparative study on nutritional and technological quality of fourteen (14) cultivars of pearl millets [*Pennisetum glaucum* (L) Leeke] in Burkina Faso. *Pakistan Journal of Nutrition*, **5**(6): 512-521.
59. Padulosi, S., Mal, B., Ravi, S.B., Gowda, J., Gowda, K.T.K., Shanthakumar, G., Yenagi, N. and Dutta, M. 2009. Food security and climate change: role of plant genetic resources of minor millets. *Indian Journal of Plant Genetic Resources*, **22**(1): 1-16.
60. Pangaribowo, E.H., Gerber, N. and Torero, M. 2013. Food and nutrition security indicators: a review.
61. Pradeep, S.R. and Guha, M., 2011. Effect of processing methods on the nutraceutical and antioxidant properties of little millet (*Panicum sumatrense*) extracts. *Food chemistry*, **126**(4): 1643-1647.
62. Prasad, P.V. and Staggenborg, S.A. 2009. Growth and production of sorghum and millets. *Soils, plant growth and crop production*, **2**(9): 186:193
63. Rao, S.V., Sunder, G.S., Panda, A.K., Reddy, M.R., Raju, M.V.L.N. and Praharaj, N.K. 2002. Utilization of different millets replacing maize in colored broiler chicken diet. *Indian Journal of Animal Nutrition*, **19**(4): 353-358.

64. Rao, M.V.S.S.T. and Muralikrishna, G. 2002. Evaluation of the antioxidant properties of free and bound phenolic acids from native and malted finger millet (Ragi, *Eleusinecoracana* Indaf-15). *Journal of Agricultural and Food Chemistry*, **50**(4): 889-892.
65. Ravindran, G.J.F.C. 1991. Studies on millets: Proximate composition, mineral composition, and phytate and oxalate contents. *Food Chemistry*, **39**(1): 99-107.
66. Rooney, L.W., Miller, F.R. and Mertin, J.V. 1981. Variation in the structure and kernel characteristics of sorghum. In *Proceedings of the international symposium on sorghum grain quality*, **28**: 143-162.
67. Saleh, A.S., Zhang, Q., Chen, J. and Shen, Q., 2013. Millet grains: nutritional quality, processing, and potential health benefits. *Comprehensive reviews in food science and food safety*, **12**(3): 281-295.
68. Shivananjappa, M., 2018. Meats are malady and millets are magical. *Journal Nutrition and Food Technology*, **1**(1): 14-15.
69. Shobana, S., Sreerama, Y.N. and Malleshi, N.G., 2009. Composition and enzyme inhibitory properties of finger millet (*Eleusine coracana* L.) seed coat phenolics: Mode of inhibition of α -glucosidase and pancreatic amylase. *Food Chemistry*, **115**(4): 1268-1273.
70. Shweta, M. 2015, Pearl millet nutritional value and medicinal uses. *International Journal of Advance Research and Innovative Ideas in Education*, **1**(3): 2395–2398.
71. Silva, G.A.H., Restle, J., Missio, R.L., Bilego, U.O., Fernandes, J.J.R., Rezende, P.L.P., Medeiros Da Silva, R., Pereira, M.L.R. and Lino, F.A. 2020. Milhetoem substituiçãoo milionadieta de novilhos confnados. *Semina Ciências Agrárias*, **35**(20): 77–94.
72. Silva, R.M.D., Taveira, R.Z., Restle, J., Fabricio, E.D.A., Camera, A., Maysonave, G.S., Bilego, U.O., Pacheco, P.S. and Vaz, F.N., 2020. Economic analysis of the risk of replacing corn grains (*Zea mays*) with pearl millet grains (*Pennisetum glaucum*) in the diet of feedlot cattle. *Ciência Rural*, **13**: 50.
73. Singh, P. and Raghuvanshi, R.S. 2012. Finger millet for food and nutritional security. *African Journal of Food Science*, **6**(4): 77-84.
74. Singh, R.B., Khan, S., Chauhan, A.K., Singh, M., Jaglan, P., Yadav, P., Takahashi, T. and Juneja, L.R. 2019. Millets as functional food, a gift from Asia to Western World. In *the Role of Functional Food Security in Global Health*, pp. 457-468.

75. Sireesha, Y., Kasetti, R.B., Nabi, S.A., Swapna, S. and Apparao, C. 2011. Antihyperglycemic and hypolipidemic activities of *Setaria italica* seeds in STZ diabetic rats. *Pathophysiology*, **18**(2): 159-164.
76. Sridhar, R. and Lakshminarayana, G. 1994 Contents of total lipids and lipid classes and composition of fatty acids in small millets: foxtail (*Setaria italica*), proso (*Panicum miliaceum*), and finger (*Eleusine coracana*). *Cereal Chemistry*, **71**: 355-359.
77. Taylor, J.R.N. 2004. In: Wrigley C, Corke H, Walker CE, 2004. Millet, *encyclopaedia in grain science*, **2**: 253–256.
78. Ullah, I., Ali, M. and Farooqi, A., 2010. Chemical and nutritional properties of some maize (*Zea mays* L.) varieties grown in NWFP, Pakistan. *Pakistan Journal of Nutrition*, **9**(11): 1113-1117.
79. Ushakumari, S.R., Latha, S. and Malleshi, N.G., 2004. The functional properties of popped, flaked, extruded and roller-dried foxtail millet (*Setaria italica*). *International journal of food science & technology*, **39**(9): 907-915.
80. Vadivoo, A.S., Joseph, R. and Meenakshi Ganesan, N., 1998. Genetic variability and diversity for protein and calcium contents in finger millet (*Eleusine coracana* (L.) Gaertn) in relation to grain color. *Plant foods for human nutrition*, **52**: 353-364.
81. Vetriventhan, M., Upadhyaya, H.D., Anandakumar, C.R., Senthilvel, S., Parzies, H.K., Bharathi, A., Varshney, R.K. and Gowda, C.L.L., 2012. Assessing genetic diversity, allelic richness and genetic relationship among races in ICRISAT foxtail millet core collection. *Plant Genetic Resources*, **10**(3): 214-223.
82. Vinoth, A. and Ravindhran, R. 2017. Biofortification in millets: a sustainable approach for nutritional security. *Frontiers in Plant Science*, **8**: 29-31.
83. Von Grebmer, K., Bernstein, J., Hossain, N., Brown, T., Prasai, N. and Yohannes, Y. 2017. Global hunger index: The inequalities of hunger. *International Food Policy Research Institute*.
84. Wankhede, D.B., Shehnaj, A. and Raghavendra Rao, M.R., 1979. Carbohydrate composition of finger millet (*Eleusine coracana*) and foxtail millet (*Setaria italica*). *Qualitas Plantarum*, **28**: 293-303.
85. Wheeler, T. and Von Braun, J. 2013. Climate change impacts on global food security. *Science*, **341**(45): 508-513.

86. WHO. Obesity and overweight Fact sheet N°311"; 2015.<http://www.who.int/mediacentre/factsheets/fs311/en/>. Accessed 23 Jan 2018.
87. Wilson, J.P., McAloon, A.J., Yee, W., McKinney, J., Wang, D. and Bean, S.R. 2007. Biological and economic feasibility of pearl millet as a feedstock for ethanol production. *Issues in new crops and new uses*, pp.56-9.
88. World Food Summit. *The world free from hunger*. Rome; 1996
89. Zarnkow, M., Mauch, A., Burberg, F., Back, W., Arendt, E.K., Kreis, S. and Gastl, M. 2009. Proso millet (*Panicum miliaceum* L.) a sustainable raw material for the malting and brewing process: A review. *Brewing Science*, **62**(8): 119-140.
90. Zhang, J.P., Zhang, J.P., Liu, T.S., Zhang, J.P., Liu, T.S., Zheng, J., Zhang, J.P., Liu, T.S., Zheng, J., Jin, Z. and Zhang, J.P., 2007. Cloning and characterization of a putative 12-oxophytodienoic acid reductase cDNA induced by osmotic stress in roots of foxtail Millet: Full-length research paper. *DNA Sequence*, **18**(2): 138-144.
91. M. SC, V. VP, M. J, I. J. IP kumar. Millets-A Powerhouse for Food Security and Combating Malnutrition. *J. Exp. Agric. Int.* [Internet]. 2023 Aug. 31 [cited 2024 May 25];45(10):44-50. Available from: <https://journaljeai.com/index.php/JEAI/article/view/2199>
92. Kanojia S, Gomadhi G, Kamalasundari S, Malathi G, Krishnaveni SA, Ramasamy M, Dhanalakshmi K, Chitra K, Debnath A. Exploring Millet Grains: Nutritional Benefits, Processing Advancements, and Future Directions for Food Security and Health Improvement. *Eur. J. Nutr. Food. Saf.* [Internet]. 2024 May 3 [cited 2024 May 25];16(5):101-8. Available from: <https://journalejnfs.com/index.php/EJNFS/article/view/1426>
93. Kumar A, Tomer V, Kaur A, Kumar V, Gupta K. Millets: a solution to agrarian and nutritional challenges. *Agriculture & food security*. 2018 Dec;7(1):1-5.