

Effect of Integrated Nutrient Management Practices on Physiological Growth Parameters of Perennial Fodder Crops in custard apple based horti-pastoral system

Commented [U1]: Better to add Nitrogen

ABSTRACT

A field experiment was conducted to assess the impact of integrated nutrient management practices on perennial fodders within a custard apple (*Annona squamosa*)-based horti-pastoral system during the Rabi season in-of 2021 at the Agroforestry Research Block of Professor Jayashankar Telangana State Agricultural University, located in Rajendranagar, Hyderabad. The study featured two main plots dedicated to the fodder species *Cenchrus ciliaris* (Anjan grass) and *Megathyrsus maximus* (Guinea grass), while seven-six nutrient management practices, along with a control, were implemented as sub-plots using-in a split-plot design. The experiment was replicated thrice among the four year old custard apple tree species. Results revealed that *M. maximus* has recorded significantly higher Physiological growth parameters viz., CGR, RGR at different growth stages in first cut. Among the integrated nutrient management practices, higher CGR values were recorded at 45 DAI with the application of T3-75% RDN + 25% N through PGLM (13.47 g m⁻² day⁻¹) which was on par with T6-the application of 75% RDN + 25% N Through PM (12.78 g m⁻² day⁻¹) and RGR found non-significant. The experiment concludes that *M. maximus* found best for horti-pastoral system with PGLM or PM supplemental nutrient management.

Commented [U2]: Give full form of PGLM and PM

Commented [U3]: Correct it

Commented [U4]: Correct it

Commented [U5]: Properly mention with dose

Keywords: Fodder species, Nutrient management practice, CGR, RGR.

Commented [U6]: Add Anjan and guinea grass in place of fodder sp.

INTRODUCTION

Cenchrus ciliaris, commonly known as Anjan grass in India, stands out-as a predominant pasture grass (Dabadghao and Shankarnarayan, 1973). The *Cenchrus* species demonstrates adaptability to a wide range of soils and environments, owing to its tolerance to drought and high temperatures. It is Hhighly palatable with substantial nutritional value (protein content of 8-10% and digestibility ranging from 60-70%),-it servesand serves as a valuable resource for various grazing animals (Sawal et al., 2009). *Megathyrsus maximus*, also known as guinea grass, is a perennial, tufted grass with a short, creeping rhizome. This robust grass can attain a height of up to 2 m and is compatible with horticultural crops. It thrives in wastewater and marshy lands, offering a crude protein content ranging from 8-13% (Prakash et al., 2019). It is Rrecommended for regions with an annual rainfall between 800 to 1800 mm and well-drained soil of medium to high fertility (Muir and Jank, 2004).

The country faces deficiencies in the availability of green and dry fodder, posing a challenge to

the future development and growth of livestock (Hembram et al., 2016; Jhonsonraju et al., 2023). Low fertility often results in reduced forage yield, and nutrient management plays a pivotal role in enhancing forage productivity. Integrated nutrient management, utilizing locally available resources like organic manures and green leaf manures, proves beneficial by reducing production costs and enhancing farm system resilience through improved soil fertility in rainfed areas (Ravi et al., 2009). A judicious combination of organic manures and chemical fertilizers, tailored to the soil and crop characteristics, not only maximizes crop production and quality but also sustains soil fertility (Madhavi Lata et al., 2014).

Organic amendments, such as farmyard manure, are recognized for enhancing soil physical properties. Poultry manure, rich in nitrogen and various micro-nutrients, serves as a readily available nutrient source for crops (Pratap et al., 2008). Pongamia green leaf, with its high content of easily mineralizable nitrogen, proves crucial for improving soil physical properties and serves as a nutrient source for low-fertility soils in dryland agriculture (Ranveer Singh et al., 2013). Therefore, this study was aimed to assess the impact of integrated nutrient management practices on the yield and economic aspects of fodder species in a horti-pastoral system.

Commented [U7]: But this study mainly focused on physiological growth parameters

MATERIALS AND METHODS

A field experiment titled Influence of Integrated Nutrient Management Practices on Perennial Fodder Species in a Custard Apple-Based Horti-Pastoral System was conducted during the *Rabi* season of 2020-2021 at the Agroforestry Research Block, Professor Jayashankar Telangana State Agricultural University, situated in Rajendranagar, Hyderabad, Telangana. The experimental site, located at 17°19' N latitude, 78°28' E longitude and at an altitude of 555 m above mean sea level, falls under the Southern Telangana agro-climatic zone. Throughout the crop growth period, which extended from October 2020 to February 2021, weekly mean maximum temperature ranged from 27.14 °C to 31.64 °C with an average of 29.29 °C. whileWhile the weekly mean minimum temperature ranged from 11.07 °C to 18.57 °C with an average of 14.85°C. The total rainfall received during this period was 18.08 mm, occurring overwith 4 rainy days.

Commented [U8]: Do same as Title

The soil at the experimental site was identified as sandy loam with a pH of(6.70), EC of(0.125 dS m⁻¹) and organic carbon of(0.66%). Soil was medium in available nitrogen (247.2 kg ha⁻¹) and phosphorus (30.65 kg ha⁻¹) while low in available potassium (179.1 kg ha⁻¹). The experiment was designed using a split-plot layout with three replications. The main plot treatments included two fodder species viz., Anjan grass and Guinea grass, and seven integrated nutrient management treatments *viz.*, T₁-Control (no fertilizer and manure), T₂-100% RDF (60:60:40 NPK kg ha⁻¹), T₃-75% RDN + 25% N through PGLM, T₄-75% RDN + 25% N through NGLM, T₅- 75% RD-N + 25% N through SGLM, T₆- 75% RDN + 25% N through PM, T₇ 75% RDN + 25% N through

Commented [U9]: No description about PGLM, NGLM, SGLM, PM in introduction nor in materials and methods.

FYM [were accommodated in sub plots](#). Phosphorus and potassium were common for all treatments except the control, and organic manure was applied as ~~a~~-basal, ~~with-while~~ nitrogen supplied in three splits after each cutting. Various physiological parameters were computed from the data obtained on the dry weight of different plant parts. Crop growth rate ($\text{g m}^{-2} \text{day}^{-1}$) is the rate of dry matter production per unit ground area per unit time and was calculated by using the formula suggested by Watson (1952).

$$\text{CGR} = \frac{W_2 - W_1}{t_2 - t_1} \times \frac{1}{A}$$

W_1 = Total dry weight of plant (g) at time t_1 , W_2 = Total dry weight of plant (g) at time t_2
 $t_2 - t_1$ = time interval (days), A = Land area (m^2)

Relative growth rate ($\text{g g}^{-1} \text{day}^{-1}$) is the rate of increase in dry weight per unit time and was calculated by using the formula suggested by Radford (1967).

$$\text{RGR} = \frac{\text{Log}_e W_2 - \text{Log}_e W_1}{t_2 - t_1}$$

W_1 = Total dry weight of plant (g) at time t_1 , W_2 = Total dry weight of plant (g) at time t_2
 $t_2 - t_1$ = time interval (days)

RESULTS AND DISCUSSION

1. Crop Growth Rate ($\text{g m}^{-2} \text{day}^{-1}$)

The data pertaining to the Crop Growth Rate (CGR) of different fodder species under the ~~influence of~~ various nutrient management practices is presented in Table 1 and visually represented in Figure 1. Upon examination of the data, significant differences in CGR at 15, 30, and 45 days after initiation (DAI) were observed between the two fodder species. Guinea grass exhibited the highest CGR at 15 [DAI](#) ($9.02 \text{ g m}^{-2} \text{day}^{-1}$), 30 [DAI](#) ($9.46 \text{ g m}^{-2} \text{day}^{-1}$), and 45 DAI ($12.26 \text{ g m}^{-2} \text{day}^{-1}$) compared to Anjan grass, which recorded CGR values of 5.39, 7.31, and $9.25 \text{ g m}^{-2} \text{day}^{-1}$ ~~at 15, 30 and 45 DAI~~, respectively.

Among the nutrient management practices, at 15 DAI maximum CGR was observed in T_3 -75% RDN + 25% N through PGLM ($8.43 \text{ g m}^{-2} \text{day}^{-1}$) followed by T_6 -75% RDN + 25% N through PM ($8.41 \text{ g m}^{-2} \text{day}^{-1}$), T_2 -100% RDF (60:60:40) NPK kg ha^{-1} ($7.63 \text{ g m}^{-2} \text{day}^{-1}$) and T_7 -75% RDN + 25% N through FYM ($7.49 \text{ g m}^{-2} \text{day}^{-1}$) which ~~was significantly~~ CGR values than other nutrient treatments and control ($5.06 \text{ g m}^{-2} \text{day}^{-1}$). ~~Similar trend was also followed at~~ 30 DAI ~~and also follows similar~~

Commented [U10]: Revise sentence

end with maximum CGR was recorded in T₃ -75% RDN + 25% N through PGLM (10.0 g m⁻¹ day⁻¹) followed by T₆ -75% RDN + 25% N through PM (9.98 g m⁻¹ day⁻¹), T₇ -75% RDN + 25% N through FYM (9.40 g m⁻¹ day⁻¹) and T₂ -100% RDF (60:60:40) NPK kg ha⁻¹ (9.35 g m⁻¹ day⁻¹). At 45 DAI, significantly higher CGR was observed in T₃ -75% RDN + 25% N through PGLM (13.47 g m⁻¹ day⁻¹) which was on par with T₆ -75% RDN + 25% N through PM (12.78 g m⁻¹ day⁻¹) and significantly higher with each other nutrient treatments and control.

The integration of organic and inorganic nutrition, specifically through PGLM, PM, and FYM sources, resulted in favorable crop growth rates comparable to those recorded with 100% RDF.

Increased plant density was found to significantly enhance CGR (Ahmadi et al., 2014), with some researchers attributing the impact of crop growth rate to the plants' photosynthetic area (HabibZadeh et al., 2006). The interaction effect between the type of fodder species and nutrient management practices on crop growth rate was found to be non-significant at all stages.


Commented [U11]: Write it like INM enhanced plant density and ultimately increased CGR

Commented [U12]: Very old ref.??discussion missing for fodder species.

Table 1. Crop Growth Rate (g m⁻² day⁻¹) of different fodder species as influenced by integrated nutrient management in horti-pastoral system

Particulars	15 DAI	30 DAI	45 DAI (1 st cut)
Main-plots (Fodder species)			
1. <i>Cenchrus ciliaris</i> (Anjan grass)	5.39+	7.31	9.25
2. <i>Megathyrsus maximus</i> (Guinea grass)	9.03±7	9.46	12.26
S.Em +	0.13	0.09	0.09
CD(P=0.05)	0.78	0.52	0.59
Sub-plots (Nutrient management)			
T ₁ -Control (No fertiliser)	5.06	6.32	8.50
T ₂ -100% RDF (60:60:40) N-P-K kg ha ⁻¹	7.63	9.35	11.61
T ₃ -75% RDN + 25% N Through PGLM	8.43	10.00	13.47
T ₄ -75% RDN + 25% N Through NGLM	6.59	6.56	9.06
T ₅ -75% RDN + 25% N Through SGLM	6.86	7.11	9.85
T ₆ -75% RDN + 25% N Through PM	8.41	9.98	12.78
T ₇ -75% RDN + 25% N Through FYM	7.49	9.40	10.01
Mean	7.20	8.38	10.75
S.Em +	0.43	0.43	0.41
CD(P=0.05)	1.27	1.26	1.20
Interaction (Main × Sub)			
S.Em +	0.61	0.61	0.58
CD(P=0.05)	NS	NS	NS
Interaction (Sub × Main)			
S.Em +	0.58	0.57	0.54
CD(P=0.05)	NS	NS	NS

Commented [U13]: Add plus minus symbol after SEm everywhere



Commented [U14]: Why gm-1, write main and sub plot just below the treatments. Add DAI after 15, 30 and 45

Figure 1. Crop Growth Rate ($\text{g m}^{-2} \text{day}^{-1}$) of different fodder species as influenced by integrated nutrient management in horti-pastoral system

2. Relative Growth rate ($\text{g g}^{-1} \text{day}^{-1}$)

Data pertaining to relative growth rate (RGR) of different fodder species as influenced by the different nutrient management practices is illustrated in Table 2. ~~Among the Perusal of the data revealed that RGR for two different types of fodder species, has been shown that~~ significantly higher RGR ($\text{g g}^{-1} \text{day}^{-1}$) recorded in guinea grass at 15 DAI (0.142), 30 DAI (0.022) and 45 DAI (0.017) ~~0.142, 0.022 and 0.0173 $\text{g g}^{-1} \text{day}^{-1}$, respectively~~ over anjan grass. The ecological advantage of a high RGR is evident as it allows the plant to rapidly increase in size, enabling it to occupy a large space both below and above the ground. Additionally, a high RGR may facilitate the rapid completion of the plant's life cycle (Ahmadi et al., 2014).

Regarding nutrient management practices, the RGR at 15, 30 and 45 DAI was found non-significant at all three stages. This might be due to similar growth rate of two forages under specific nutrient management treatments. All the sources are better in influencing RGR. The interaction effect between the type of fodder and nutrient management practices was also found to be non-significant.

Table 2. Relative Growth Rate ($\text{g g}^{-1} \text{day}^{-1}$) of different fodder species as influenced by integrated nutrient management in horti- pastoral system

Particulars	15 DAI	30 DAI	45 DAI (1 st cut)
Main plots (Fodder species)			
1. <i>Cenchrus ciliaris</i> (Anjan grass)	0.127	0.017	0.017 9
2. <i>Megathyrsus maximus</i> (Guinea grass)	0.142	0.022	0.017 3
S.Em \pm	0.001	0.001	0.001

CD(P=0.05)	0.006	0.003	0.003
Sub-plots (Nutrient management)			
T ₁ -Control (No fertiliser)	0.124	0.016	0.020
T ₂ -100% RDF (60:60:40) N-P-K kg ha ⁻¹	0.136	0.020	0.018
T ₃ -75% RDN + 25% N Through PGLM	0.139	0.020	0.017
T ₄ -75% RDN + 25% N Through NGLM	0.132	0.020	0.018
T ₅ -75% RDN + 25% N Through SGLM	0.133	0.021	0.017
T ₆ -75% RDN + 25% N Through PM	0.139	0.021	0.016
T ₇ -75% RDN + 25% N Through FYM	0.136	0.020	0.015
Mean	0.134	0.0199	0.0172
S.Em ±	0.004	0.001	0.001
CD(P=0.05)	NS	NS	NS
Interaction (Main × Sub)			
S.Em ±	0.006	0.002	0.001
CD(P=0.05)	NS	NS	NS
Interaction (Sub × Main)			
S.Em ±	0.005	0.001	0.001
CD(P=0.05)	NS	NS	NS

CONCLUSION

Based on the findings, it can be concluded that guinea grass with the application of 75% RDN + 25% N through PGLM recorded higher CGR and RGR which was on par with 100% RDF (60:60:40) N-P-K kg ha⁻¹, 75% RDN + 25% N through PM and 75% RDN + 25% N Through FYM. This resulted in increased yields, including total green and dry fodder, as well as favorable economic returns. The study underscores the significance of incorporating organic or green leaf manures in conjunction with chemical fertilizers for sustainable development.

Commented [U15]: Redo. Make it simple and easy to understand

REFERENCES

Commented [U16]: Lots of mistakes in this section. Follow Journal style, >85% Ref. is more than 5 Year old. Add new Ref. to improve your work

Dabadghao, P.M. and Shakarnarayan, K.A. 1973. *The grass cover of India*. (Indian council of Agricultural Research: New Delhi). P.714.

Ahmadi, B., Shirani Rad, A.M. and Delkhosh, B., 2014. Evaluation of plant densities on analysis of growth indices in two canola forage (*Brassica napus* L.). *European J Exp Biol*, 4(2): 286-294.

Habibzadeh, Y, R. Mamghani, A. Kashani, Iranian J. Crop. Sci., 2006, 8, 66-78

Commented [U17]: Incomplete

Hembram, J.A.D.U.N.A.T.H. and Kundu, C.K., 2016. Effect of integrated nutrient management on growth, yield and quality of forage cropping sequence. *Journal of Agricultural Science and Research*, 3(1), 17-22.

Jhonsonraju, S., Krishna, A., Madhavalata, A., Chaitanya, T. and Sekhar, C.C., 2023. Influence of integrated nutrient management practices on yield and economics of fodders in custard apple based horti-pastoral system. *Range Management and Agroforestry*, 44(1), pp.198-203.

Karforma, J., Ghosh, M., Ghosh, D. C., Mandal, S. and Ghosh P.K. 2016. Effect of integrated management on soil fertility and productivity in maize. *Bulletin of environment, Pharmacology and life sciences*. 2(8):61-67.

Commented [U18]: Not cited in text

Madhavi Lata, A., Srinivasa Raju, M., Joseph, B., Chandrasekhara Rao, P and Siva Sankar, A. 2014. Integrated nutrient management in Ashwagandha (*Withaniasomnifera* L.) under tree based cropping systems in drylands. *Indian Journal of Agroforestry*. 16 (1): 30-36.

Madhavi Lata, A., Srinivasa Raju, M., Joseph, B., Chandrasekhara Rao, P and Siva Sankar, A. 2014. Integrated nutrient management in Ashwagandha (*Withania somnifera* L.) under tree based cropping systems in drylands. *Indian Journal of Agroforestry*. 16 (1): 30-36.

Commented [U19]: Repeattion

Muir P James and Liana jank. 2004. Guinea grass. *Warm season (C4) grasses*:589-621

Patel, P.R., Patel, B.J., Vyas, K.G. and Yadav, B.L., 2014. Effect of integrated nitrogen management and bio-fertilizer in kharif pearl millet (*Pennisetum glaucum* L.). *Adv. Res. J. Crop Improv*, 5(2), 122-125.

Commented [U20]: Not cited in text

Pallavi, Ch., Joseph, B., Aariff Khan, M.A and Hemalatha, S. 2015. Yield, nutrient content, uptake, and available nutrient status of finger millet as influenced by nutrient management in agri-silvi sytem. *International journal of current research*. 7(11): 22311-22314.

Commented [U21]: Not cited in text

Prakash kumar Rathod and Sreenath Dixit. 2019. Green fodder production: A manual for field functionaries. Patancheru 502 324, Telangana, India: International Crops Research Institute for the Semi-Arid Tropics. 56

Pratap, R., Sharma, O.P and Yadav, G.L. 2008. Effect of integrated nutrient management under varying levels of zinc on pearl millet yield. *Annals of Arid Zone*. 47 (2): 197-199

Radford, P.J., 1967. Growth analysis formulae-their use and abuse 1. *Crop science*, 7(3): 171-175.

Ranveer Singh., Ram, T., Choudhary, G.L and Gupta, A.K. 2013. Effect of integrated nitrogen management on nutrient uptake, quality, economics and soil fertility of pearl millet under rainfed conditions. *Elixir Agriculture*. 54:12373-12375.

Rasool, S., Kanth, R.H., Shaban Hamid, Raja, W., Alie B. A. and Dar Z.A.2015. Influence of integrated nutrient management on growth and yield of sweet corn under temperate conditions of Kashmir valley. *American journal of experimental agriculture*. 7(5): 315-325.

Commented [U22]: Not cited in text

Ravi. R., Divya, M.P., Ratha Krishnan. 2009. Evaluation of fodder crops under *Ailanthus excelsa roxb.* Based agroforestry system. *Indian journal of agroforestry*. 11(2), 90-93.

Sawal R K, Ratan R, Chander S. 2009. Nutritive evaluation of *Lasiurus sindicus* and *Cenchrus ciliaris* hays in sheep. *Indian Journal of Small Ruminants* 15(2): 277-80.

Watson, D.J., 1952. The physiological basis of variation in yield. In *Advances in agronomy* (4): 101-145.

UNDER PEER REVIEW

