

Evaluation of bio-stimulants for active growth, yield and shelf life of papaya cv. Red Lady

ABSTRACT

Bio-stimulants, which are natural substances aiding plant growth and resilience, are comprised of various organic and mineral compounds, serving as eco-friendly tools for sustainable farming. The research study was conducted at the Horticulture Farm of Agriculture at Lovely Professional University in Phagwara, Punjab, during the years 2023-24. The study employed a randomized block design with seven treatments replicated three times. Bio-stimulants were sprayed at 60, 90, 120, and 150 days after planting (DAP) on papaya plants cv. Red Lady to evaluate its effect on various growth, quality and shelf-life parameters. Various growth parameters including plant height, stem girth, canopy spread, petiole length of leaves, as well as floral characteristics such as days taken for flowering, height at initial flowering, number of flowers per node, fruit set percentage and yield-related parameters were evaluated. The study found significant effects of different bio-stimulant applications on papaya growth, flower and fruit production, fruit quality and shelf-life. Treatment T₄ (RDF + Diatomaceous earth) showed notable improvements in growth parameters, while treatment T₅ (RDF + Panchgavya) resulted in enhanced productivity, fruit quality and shelf-life. Overall, the study suggests that the application of Diatomaceous earth alongside fertilizers promotes vegetative growth, while Panchgavya application enhances flowering, fruiting and quality attributes, offering insights for optimizing papaya cultivation practices for improved yield and quality.

Key words: Papaya, Red lady, bio-stimulants, panchgavya, growth, yield, quality and shelf-life

1. INTRODUCTION

Papaya (*Carica papaya* L.) characterized as an herbaceous plant [1], exhibits a variable growth pattern depending on the specific variety, with some reaching heights of up to ten meters [2]. The leaves of papaya plants are palmately-lobed [3], featuring lengthy [4], hollow petioles and blades divided into 5-9 segments [5]. Flower buds develop at the axils of the leaves [6] and the fertilized fruit can contain up to 1000 seeds [5]. The fruit's skin transitions from green in its unripe stage to a vibrant yellow orange when it

ripens [7]. In general, papaya plants have a lifespan ranging from five to ten years [2]. Papaya differs in its nutrient requirements as compared to other crops as flowering, fruit initiation and formation and fruit development is a continuous phenomenon. The source-sink interaction is also very prominent for this crop and such it responds very well to nutrient application whether basal or foliar [8].

Bio-stimulants refer to the formulations that reduce the dependency of plants on fertilizers [9]. These provide an impetus to the plant growth and provide a resistance mechanism to the plants against various abiotic stresses [10]. Although these are not comparable to the plant growth regulators, yet when applied in small concentrations, tend to streamline the plant physiological processes resulting in high yields and fairly good quality of fruits [11]. Application of plant bio-stimulants enhances the nutritional efficiency of the plant [12], plant morphological traits [13] resistance to abiotic stresses irrespective of the source of nutrients used [14]. Bio-stimulant compositions have varied ingredients such as humic acids, algal extracts, hormones and growth promoting bacteria for plants [15]. While they are not the same as fertilizers, they are seen as eco-friendly tools for sustainable farming, promoting healthier plant growth without harming the environment [11]. Bio-stimulant like Panchgavya is a form of organic liquid made purely from 5 organic ingredients like cowdung, cow urine, cow milk, curd and ghee [16]. All the five constituents are known to have medicinal properties and were used singly or in combination against many diseases. It is well documented that the organically produced fruits, vegetables and grains are more beneficial for human health [17] and contain higher hydrolysable polyphenols and higher soluble and hydrolysable antioxidant capacities [18]. Several studies indicated the positive effect of foliar application of panchagavya food crops etc. [19,20]. Various other bio-stimulants like vermiwash, jeevamrutham and brassinosteroids have shown positive effect on the growth and development of fruit plants. This study was planned to evaluate the effect of foliar application of bio-stimulants on growth, yield and shelf life of papaya cv. Red Lady.

2. MATERIAL AND METHODS

The current study was conducted at the Horticultural Farm and within the laboratories of the School of Agriculture, Lovely Professional University, located in Phagwara, Punjab, throughout the duration of 2023. Experimental site is located within the Doaba region of Punjab, situated at coordinates 31.25° N latitude and 75.70° E longitude, with an elevation of 249 meters above sea level. For this experiment, various bio-stimulants were employed across seven distinct treatments, alongside a control group. These bio-stimulants were administered through both foliar and soil applications. The experimental design adopted was a randomized block design, with three replications. The research was conducted in randomized block designs (RBD). Total number of treatments was 7 (T₁: Recommended dose of fertilizer (Conventional method), T₂: RDF + seaweed extract spray, T₃: RDF + Brassinosteroids, T₄: RDF + Diatomaceous earth, T₅: RDF + Panchagavya, T₆: RDF + Vermiwash, T₇: RDF + Jeevamrutham) having 3 replications (each replication had 2 plants per treatment), each, with total number of plants 42 for papaya cv. Red Lady.

2.1 OBSERVATIONS RECORDED

Every 30 days after spraying till the harvesting of fruits, vegetative parameters were recorded. Plant height was recorded using a graduated stick. Stem girth was recorded using a Vernier caliper. Number of leaves of selected plants from each replication were numerically counted, Petiole length was measured at 150 and 240 days of transplanting by taking three tagged leaf petioles from the top. It was measured with the help of measuring tape. Plant Spread was recorded using a measuring tape. Number of days taken from transplanting till the opening of first female/hermaphrodite flower was recorded to access days to flowering. Height of the plant from ground level to the node where first flower appeared was recorded. Number of flowers produced per node in each plant was recorded for three months, starting from first flowering. Number of fruits per plant, Fruit physical characters, Fruit yield per plant, Days taken for maturity were recorded as per standard methods. Fruit Acidity (%) and Ascorbic acid (mg/100g) was calculated as per the method given by Ranganna (1977), Shelf-life parameters were recorded at an interval of 4 days and were analyzed using standard methods of observations.

The data generated was statistically analyzed using OPSTAT software in a One way Anova model.

3. RESULTS AND DISCUSSION

3.1 VEGETATIVE PARAMETERS

The maximum plant height on 150th DAP was observed under T₄ (RDF+ Diatomaceous earth) having the value of 152.33cm followed by T₆ (RDF + Vermiwash) with 146.83 cm of plant height. Treatment T₄ was found significant for stem girth which recorded 8.11 cm of stem girth at 150th DAP followed by T₇ (RDF+ Jeevamrutham) with 8.01cm of stem girth. There was a notable difference in the treatment T₄ (RDF + Diatomaceous earth) which had the highest number of leaves (31) at 150th DAP followed closely by treatment T₅ (RDF + Panchagavya) with 30.67 leaves per plant. Treatment T₄ (RDF + Diatomaceous earth) resulted in maximum canopy spread (197.83 N-S; 203.33 E-W) followed by the treatment T₂ (RDF + Seaweed extract) with 195.33 cm N-S; 198.83 cm E-W at 150th DAP. The findings at 240 DAP revealed a substantial difference: the treatment T₄ had the longest petiole length of leaves (67.06 cm) followed by the treatment T₃ (RDF + Brassinosteroids) resulted a petiole length of 66.50 cm. The maximum chlorophyll index value of 35.90 was reported in the treatment T₄ (RDF + Diatomaceous earth) at 150th DAP followed by the treatment T₅ (RDF + Panchagavya). It is clear from the obtained data in Table (1) that spraying papaya cv. Red lady with diatomaceous earth along with recommended dose of fertilizers at 30, 60, 90, 120 DAP significantly accompanied with enhancing plant height, stem girth, Leaf count per plant, Canopy spread, Petiole length and Chlorophyll index. These findings were similar with that of [21] in Hindi Khalsa mango, [22] in orange, [23] in Keitte mango. According to [24], plants sprayed with silicon showed significantly greater absorption of nitrogen, potassium, phosphorus and zinc. The

application of Diatomaceous Earth as a silicon source led to an increase in leaf chlorophyll content, which consequently reduced chlorophyll degradation, reported by [25] in Pomegranate var. Kesar.

Table 1. Effect of bio-stimulant administrations on Vegetative Parameters of Papaya cv. Red Lady at 150 Days after planting

Treatments	Plant height (cm)	Stem girth (cm)	No. of Leaves	Canopy spread (N-S)	Canopy spread (E-W)	Petiole length of leaves (cm) [150, 240 days]		Chlorophyll index of Leaves
T ₁	138.5	7.34	28.67	179.67	170.33	56.33	63.72	32.95
T ₂	139.5	7.8	29.5	195.33	198.83	60	64.78	33.27
T ₃	144.67	7.95	30.17	185.67	192.5	65.39	66.5	34.27
T ₄	152.33	8.11	30.5	197.83	203.33	65.72	67.06	35.9
T ₅	139.33	7.89	31	188.83	188.67	59.33	64.67	35.9
T ₆	146.83	7.91	30.67	194.5	174.5	62.67	64.39	35.07
T ₇	142	8.01	29.17	194.17	173.5	61.17	64.5	34.72
S.Em (±)	4.5	0.31	0.8	6.23	8.87	1.73	1.36	0.74
C.D. at 5%	9.08	0.63	1.61	12.5	17.8	3.49	2.73	1.49

T₁= Recommended dose of fertilizer, T₂= RDF + Seaweed extract, T₃= RDF + Brassinosteroids, T₄= RDF + Diatomaceous earth, T₅ = RDF + Panchagavya, T₆ = RDF + Vermiwash, T₇ = RDF + Jeevamrutham.

3.2 FLOWER AND FRUIT PARAMETERS

The treatment T₅ combining RDF with Panchagavya resulted in the increased number of flowers per node (6.33) at 150th DAP followed by the treatment T₇ (RDF + Jeevamrutham) which produced 5.33 flowers per node. The minimum days to flower opening was observed for the treatment T₅ (RDF + Panchagavya) with minimum of 99.67 days to open flowers followed by the treatment T₇ (RDF + Jeevamrutham) with 103.67 days. The maximum fruit set percentage of 70.35 was observed under the treatment T₅ followed by the treatment T₇ of about 69.85 % of fruit set. Treatment T₅ demonstrated the superior performance with 36.83 fruits per plant, followed by the treatment T₃ (RDF + Brassinosteroids) with 34.83 fruits per plant. Statistical examination of data regarding fruit yield per plant and fruit yield per hectare as depicted in Table (2) reveals that the various treatments exerted a significant influence on fruit yield. The treatment T₅ recorded the higher fruit yield of 56.78 kg/plant and 83.83 t/hectare followed by 44.83 kg/plant and 83.10 t/hectare with treatment T₇. The treatment T₅ (RDF + Panchagavya) resulted in the substantially highest length and width of fruits (24.2 cm and 21.0 cm), followed by the treatment T₇ (RDF+ Jeevamrutham) with the fruit length and width of about 21.26 cm and 20.23 cm. The minimum days taken for fruit maturity (197.6) was observed with the treatment T₅ (RDF + Panchagavya) followed by the treatment T₇ (RDF + Jeevamrutham) with minimum days of 198 for fruit maturity. Highest fruit weight was observed under the treatment T₅ (0.97 kg) followed by the treatment T₇ of about 0.96 kg.

These readings were similar with that of [26] in Bitter gourd, [27] in Capsicum.

Table 2. Effect of bio-stimulant administrations on flower and fruit parameters of Papaya cv. Red Lady

Treatments	No.of flowers/node	Days to flower opening	Fruit set percentage (%)	No. of fruits/plant	Fruit yield per plant (kg)
T ₁	4.5	125.17	63.44	31.83	19.57
T ₂	4.33	122.83	66.43	33.67	26.68
T ₃	4.67	106.5	68.88	34.83	43.35
T ₄	5	108.83	65.55	33.83	25.01
T ₅	6.33	99.67	70.35	36.83	56.78
T ₆	4.67	107.33	66.63	34.5	35.29
T ₇	5.33	103.67	69.85	32.5	44.83
S.Em (±)	0.37	8.78	3.87	1.72	2.255
C.D. at 5%	0.73	17.7	7.79	3.46	7.02

T₁= Recommended dose of fertilizer, *T₂*= RDF + Seaweed extract, *T₃*= RDF + Brassinosteroids, *T₄*= RDF + Diatomaceous earth, *T₅* = RDF + Panchagavya, *T₆* = RDF + Vermiwash, *T₇* = RDF + Jeevamrutham.

Treatments	Fruit yield per hectare (t)	Days taken for maturity	Fruit length (cm)	Fruit width (cm)	Fruit weight (Kg)
T ₁	50.46	207.17	13	12.03	0.63
T ₂	66.7	202.88	16.73	14.83	0.8
T ₃	79.86	200.50	19.6	18.93	0.92

T ₄	62.53	203.83	14.86	13.46	0.74
T ₅	83.83	197.67	24.2	21	0.97
T ₆	79.33	202.17	18.33	16.53	0.9
T ₇	83.1	198	21.26	20.23	0.96
S.Em (±)	4.12	4.42	0.3	0.27	0.02
C.D. at 5%	12.83	1.42	0.94	0.85	0.063

T₁= Recommended dose of fertilizer, T₂= RDF + Seaweed extract, T₃= RDF + Brassinosteroids, T₄= RDF + Diatomaceous earth, T₅ = RDF + Panchagavya, T₆ = RDF + Vermiwash, T₇ = RDF + Jeevamrutham.

3.3 QUALITY PARAMETERS

The maximum pulp weight of 0.8 kg was noticed under the treatment T₅ (RDF + Panchagavya) followed by the treatment T₇ (RDF + Jeevamrutham) having the pulp weight of 0.79 kg. Treatment T₅ (RDF + Panchagavya) recorded the maximum peel weight and seed weight of 92.41 g and 86.06 g respectively followed by the treatment T₇ (RDF + Jeevamrutham) having the peel and seed weight of about 89.75 g and 84.16 g respectively. The finding of results shows that minimum recorded acidity was found in the treatment T₅ having the value of 0.019 % followed by the treatment T₇ with the acidity value of 0.02 %. TSS concentration was notably highest in the treatment T₅ (RDF + Panchagavya) with a reading of 8.93 °Brix, followed by treatment T₇ (RDF + Jeevamrutham) with 8.6 °Brix. The maximum concentration of ascorbic acid was detected for the treatment T₅ (RDF + Panchagavya) of 41.91 mg/100g, followed by treatment T₇ (RDF + Jeevamrutham) of about 40.06 mg/100g. The highest sugars % (reducing, Non-reducing and Total sugars) was observed under the treatment T₅ (RDF + Panchagavya) with the readings of 8.56%, 1.25% and 10.26% respectively followed by the treatment T₇ (RDF + Jeevamrutham) with the readings of 8.30%, 1.24% and 9.98%. Research by [28] indicates that the improved performance of plants treated with panchagavya can be attributed to its properties as a growth promoter and immunity booster, as well as its ability to mitigate common diseases. Similarly, [29] and [30] support these findings.

Table 3. Effect of bio-stimulant treatments on quality parameters of Papaya cv. Red Lady

Treatments	Pulp weight (kg)	Peel weight (g)	Seed weight (g)	Titrateable acidity (%)	Total soluble solids
T ₁	0.48	74.11	69.76	0.027	6.6

T ₂	0.63	83.56	77.46	0.025	6.77
T ₃	0.76	88.06	81.98	0.021	7.96
T ₄	0.58	78.86	74.3	0.026	6.78
T ₅	0.8	92.41	86.06	0.019	8.93
T ₆	0.72	86.11	78.56	0.024	7.43
T ₇	0.79	89.75	84.16	0.02	8.6
S.Em (±)	0.02	0.55	2.55	0.001	0.41
C.D. at 5%	0.064	1.71	0.82	0.002	1.29

T₁= Recommended dose of fertilizer, T₂= RDF + Seaweed extract, T₃= RDF + Brassinosteroids, T₄= RDF + Diatomaceous earth, T₅ = RDF + Panchagavya, T₆ = RDF + Vermiwash, T₇ = RDF + Jeevamrutham.

Treatments	Ascorbic acid (mg/100g)	Reducing sugars (%)	Non-reducing sugars (%)	Total sugars (%)
T ₁	34.31	5.841	1.135	7.283
T ₂	37.3	7.177	1.163	8.718
T ₃	39.5	8.186	1.227	9.843
T ₄	36.28	6.829	1.155	8.343
T ₅	41.91	8.56	1.253	10.263
T ₆	38.03	7.293	1.192	8.868
T ₇	40.06	8.308	1.243	9.988
S.Em (±)	0.48	0.16	0.01	0.17
C.D. at 5%	1.51	0.5	0.04	0.53

T₁= Recommended dose of fertilizer, T₂= RDF + Seaweed extract, T₃= RDF + Brassinosteroids, T₄= RDF + Diatomaceous earth, T₅ = RDF + Panchagavya, T₆ = RDF + Vermiwash, T₇ = RDF + Jeevamrutham.

3.4 SHELF LIFE PARAMETERS

Physiological loss in weight was minimum with the treatment T₅ (RDF + Panchagavya) having the value 11.7 %, followed by the treatment T₇ (RDF + Jeevamrutham) with the value 12.9%. The data of findings shows that maximum shelf life was noted in the treatment T₅ (RDF + Panchagavya) with maximum days of about 6.83 days followed by the treatment T₇ (RDF + Jeevamrutham) remain

marketable for about 6.16 days. Similar findings were observed by [31] in bitter gourd and [32] in Tomato. Panchagavya has been found to contain growth regulators such as IAA, GA and cytokinin, as demonstrated by [33]. Panchagavya harbours the highest populations of total bacteria, actinomycetes, phosphate solubilizers, fluorescent pseudomonads and nitrifiers among the microorganisms [32]. Dehydrogenase activity and microbial biomass were greater in Panchagavya treated plants [34]. The outcomes align with the findings presented by [35] in their research involving the strawberry cultivar Camarosa.

Table 4. Effect of bio-stimulant treatments on shelf- life parameters of Papaya cv. Red Lady

Treatments	Physiological loss in weight (%)	Shelf life (days)
T ₁	18.418	3.167
T ₂	16.355	4.5
T ₃	13.673	5.833
T ₄	17.778	4.167
T ₅	11.72	6.833
T ₆	15.172	5.167
T ₇	12.917	6.167
S.Em (±)	0.23	0.25
C.D. at 5%	0.73	0.79

T₁= Recommended dose of fertilizer, *T₂*= RDF + Seaweed extract, *T₃*= RDF + Brassinosteroids, *T₄*= RDF + Diatomaceous earth, *T₅* = RDF + Panchagavya, *T₆* = RDF + Vermiwash, *T₇* = RDF + Jeevamrutham.

4. CONCLUSION

Based on the findings of this study, it is recommended that papaya trees should be treated with Diatomaceous earth in conjunction with the recommended fertilizer dosage to enhance vegetative growth. Additionally, the application of Panchagavya alongside the recommended fertilizer dosage is advised to promote earlier flowering, increase flower production, improve fruit set percentage, boost fruit yield, enhance fruit quality, and prolong shelf life.

REFERENCES

1. Godínez, L. J. G., Escobar, J. G. R., Rodríguez-Falconi, R., Arteaga-Garibay, R. I., & Avendaño-Arrazate, C. H. (2024). Isolation and Characterization of fungal pathogens

associated with *Carica papaya* L. and their biocontrol with *Trichoderma* sp.: Aislamiento y caracterización de patógenos fungicos asociados a *Carica papaya* L., y su biocontrol con *Trichoderma* sp. *Agro Productividad*.

2. Ávila-Hernández, J. G., del Rosario Cárdenas-Aquino, M., Camas-Reyes, A., & Martínez-Antonio, A. (2023). REVIEW Sex Determination in Papaya: Current Status and Perspectives. *Plant Science*, 111814.
3. Dutta, S., Das, M., & Mitra, A. (2024). Macro-microscopic and phytochemical studies of Papaya (*Carica papaya* L.) with special reference to quality assurance profiles. *Journal of Drug Research in Ayurvedic Sciences*, 9(2), 99-110.
4. Encina, C. L., Granero, M. L., & Regalado, J. J. (2023). In Vitro Long-Term Cultures of Papaya (*Carica papaya* L. cv. Solo). *Horticulturae*, 9(6), 671.
5. Bunu, S. J., Okei, J. O., Miediegha, O., Ebeshi, B. U., & Chukwuemerie, O. L. (2023). Assessment of Secondary Metabolites and Thin-Layer Chromatographic Analysis of *Carica papaya* (Caricaceae) Leaves Ethanolic Extract. *Journal of Pharmaceutical Research International*, 35(36), 21-28.
6. Sahu, M. K., Sharma, G. L., Panigrahi, H. K., & Singh, P. (2023). Studies on influence of plant growth regulators on flowering, fruiting, fruit setting, and maturity of papaya (*Carica papaya* L.) cv. Red Lady under net house condition.
7. Godfried, a., adade, d., obeng, m., & ayikpah, a. y. an investigation into fruit supply chains in ghana: the case of pineapple and pawpaw.
8. Auxilia, J., Sujatha, K.B and Prakash Patil (2022). Precision Farming in Papaya for the Enhancement of Fruit Yield and Quality. *Biological Forum – An International Journal*, 14(3): 484-489.
9. Miranda, A. M., Hernandez-Tenorio, F., Villalta, F., Vargas, G. J., & Sáez, A. A. (2024). Advances in the Development of Biofertilizers and Biostimulants from Microalgae. *Biology*, 13(3), 199.
10. Trivedi, K., Vijay Anand, K. G., Kubavat, D., Kumar, R., Vaghela, P., & Ghosh, A. (2017). Crop stage selection is vital to elicit optimal response of maize to seaweed bio-stimulant application. *Journal of applied phycology*, 29, 2135-2144.
11. Paradikovic, N., Teklic, T., Zeljkovic, S., Lisjak, M., & Špoljarevic, M. (2018). Biostimulants research in some horticultural plant species-a review. *Food Energy Secur.* 8: e00162.

12. Reddy, K. S., Wakchaure, G., Khapte, P., & Changan, S. (2023). Plant Bio-stimulants for Mitigating Abiotic Stresses in Agriculture. *Indian J. Fertil*, 19, 788-800.
13. Senousy, H. H., Hamoud, Y. A., Abu-Elsaoud, A. M., Mahmoud Al zoubi, O., Abdelbaky, N. F., Zia-Ur-Rehman, M., ... & Soliman, M. H. (2023). Algal Bio-Stimulants Enhance Salt Tolerance in Common Bean: Dissecting Morphological, Physiological, and Genetic Mechanisms for Stress Adaptation. *Plants*, 12(21), 3714.
14. HONGAL, G., SIDDARAJU, R., BAI, S. K., VISHWANATH, K., RAMANAPPA, T., & PAVAN, A. (2023). Bio-Stimulants in Oil Seed Crops-Its Influence on Growth, Yield and Quality. *Mysore Journal of Agricultural Sciences*, 57(4).
15. Dubey, S. C., & Sharma, K. (2023). Biostimulant: an innovative approach for sustainable crop production. *Current Science* (00113891), 125(4).
16. Jnanasha, A. C., Venugopal, S., Kumar, S. R., Kumar, A., Bisht, D., Chanotiya, C. S., & Lal, R. K. (2024). Optimization of a new organic approach to natural biostimulant (Jeevamrutha) for yield and quality management in Senna (*Cassia angustifolia* Vahl.): an agriculturally highly export-oriented crop. *Technology in Horticulture*, 4(1).
17. Rahman, A., Baharlouei, P., Koh, E. H. Y., Pirvu, D. G., Rehmani, R., Arcos, M., & Puri, S. (2024). A Comprehensive Analysis of Organic Food: Evaluating Nutritional Value and Impact on Human Health. *Foods*, 13(2), 208.
18. Mercado-Mercado, G., Blancas-Benitez, F. J., Velderrain-Rodríguez, G. R., Montalvo-González, E., González-Aguilar, G. A., Alvarez-Parrilla, E., & Sáyago-Ayerdi, S. G. (2015). Bioaccessibility of polyphenols released and associated to dietary fibre in calyces and decoction residues of Roselle (*Hibiscus sabdariffa* L.). *Journal of functional foods*, 18, 171-181.
19. Behera, S. R., Pandey, R., Golui, K., Sahoo, S., Jakhwal, R., & Pal, R. (2024). Application of Panchagavya, a Cow-based Liquid Formulation, as a Lever for Sustainable and Enhanced Vegetable Crop Production: A Review. *International Journal of Environment and Climate Change*, 14(5), 214-232.
20. Brar, P. S., Bhardwaj, G., & Kaushal, R. (2024). Influence of organic amendments, PGPR and fermented folk liquid formulations on bell pepper in mid-hill conditions of Himachal Pradesh state of India. *Journal of Plant Nutrition*, 47(1), 97-109.

21. Abd El-Gawad, N. H. G., A. I. Abu El-Azm and M. S. Hikal (2017). Effect of potassium silicate on tuber yield and biochemical constituents of potato plants grown under drought stress conditions. *Middle East J. Agri. Res.*, 6 (3):718-731.
22. El-Giousy, S. F. (2016). Productivity, fruit quality and nutritional status of 'Washington Navel' Orange trees as influenced by foliar application with salicylic acid and potassium silicate combinations. *J. Horti. Sci. & Ornamental Plants*. 8 (2): 98- 107.
23. Abd El-Rahman, M. M. A. (2015). Relation of spraying silicon with fruiting of Keitte mango trees growing under Upper Egypt conditions. *Stem Cell*, 6 (2):1-5
24. Satisha, G. C., A. K. Saxena, and A. N. Ganeshamurthy (2017). Effect of silicon and micronutrients on plant growth, yield and disease incidence in chili (*Capsicum annum L.*). proceedings of abstracts, 7th International conference on silicon in agriculture. pp. 144.
25. Kalatippi, Anand. (2017). Effect of Diatomaceous Earth on Chlorophyll Content of Leaves, Yield and Disease Occurrence in Pomegranate var. Kesar. *International Journal of Pure & Applied Bioscience*. 5. 1298-1303. 10.18782/2320-7051.2982.
26. Rana, D. (2023). Influence of Panchgavya on Growth, Yield and Quality parameters of Bitter Gourd (*Momordica charantia L.*). *International Journal of Plant & Soil Science*, 35(18), 1408-1415.
27. Mishra, Nityamanjari & Sahu, G & Mishra, P & Ray, Monika. (2015). I J T A Effect of Panchagavya on Growth and Yield of Capsicum. 33. 2613-2616.
28. V. Tiwari, S. Maji, S. Kumar, G. Prajapati and R. Yadav (2016). Use of kitchen waste-based bio-organics for strawberry (*Fragaria x ananassa Duch*) production. *Afr. J. Agric. Res.*, 11: 259-265.
29. J. Vallimayil and R. Sekar (2012). Investigation on the effect of panchagavya on Southern Sunnhemp mosaic virus (SSMV) infected plant systems. *Global J. Env. Res.*, 6: 75-79.
30. P. U. Amareswari and P. Sujathamma (2014). Jeevamritha as an alternative of chemical Fertilizers in rice production. *Agric. Sci. Dig.*, 34: 240-242.

31. Somashekar Gajjela, Ranjit Chatterjee, Sushmita Subba and Eggadi Ramesh. 2018. Studies on Storage Behaviour of Bitter Gourd as Influenced by Liquid Organic Manures. *Int.J.Curr.Microbiol.App.Sci.* 7(10): 1427-1434.
32. Muthukumar, M., Jeyakumar, P., Sriharan, N., Somasundaram, E., Ganesan, K., Effect of Organic and Inorganic Fertilizers on Postharvest Physiological Characters of Tomato, *Int. J. Pure App. Biosci.* 7(3): 454-460 (2019)
33. S. Sreethu and S. Singh (2020). Effect of nitrogen and panchagavya on growth and yield of baby corn yield of baby corn (*Zea mays L.*). *The Bioscan*, 15: 243-246.
34. E. L. D. Amalraj, G. P. Kumar, S. K. M. H. Ahmed, R. Abdul and N. Kishore (2013). Microbiological analysis of panchagavya, vermicompost and FYM and their effect on plant growth promotion of pigeon pea (*Cajanus cajan L.*) in India. *Org. Agric.*, 3: 23-29.
35. Senthilkumar, S., Kondi, R. K. R., Kumar, K. P., Kumar, S. M., Reddy, P. R., & Sravani, S. (2023). A study of certain organic inputs on phenological, yield and quality related traits of strawberry (cv. Camarosa) in pot culture. *Emergent Life Sciences Research*, 9, 77-82.