

Effect of PGR and Micronutrients on Growth, Yield and Quality of Strawberry (*Fragaria × ananassa*) cv. Winter Dawn

ABSTRACT :-

The experiment was laid out in Randomized Block Design which had 8+1(control) Treatments replicated thrice. The vegetative growth and fruit yield is concerned, application of different treatment of Plant Growth Regulator and Micronutrient significantly enhanced the Plant Height(cm), Number of leaves per plant, Plant Spread(cm) and Yield. Maximum (Plant Height (19.00cm), No. of leaves (20.67), leaf area (130.33cm²), Plant spread (25.50cm), (No. of fruit per plant(20.44), No. of flower per plant (32.00), fruit yield (9.21q/ha) were recorded best in treatment T2 (75ppm Gibberellic acid) in terms of Vegetative Parameter.

1. INTRODUCTION:-

The modern cultivation strawberry (*Fragaria x ananassa* Duch) is a hybrid between two largely dioecious octaploid species, *Fragaria chiloensis* and *Fragaria virginiana*. *Fragaria* species belongs to the family Rosaceae, with basic chromosome number of $x=7$. However the cultivated strawberry, (*Fragaria x ananassa*), is an octaploid having chromosome number of $2n=56$. In addition to *Fragaria x ananassa* the genus *fragaria* includes atleast 17 other species including diploids, tetraploids, octaploids and a hexaploids. Initially strawberry was growing in temperate zone of the country. However, it can also be cultivated under subtropical climate, even at higher altitudes of tropical climate and it is propagated through one year old runners.

Nutritionally strawberry is a low calorie carbohydrate fruit but a rich source of vitamin A (60IU/100g of edible portion), vitamin C (30-120mg/100g of edible portion) and fiber and also has high pectin (0.55%), available in the form of calcium pectate. Water is a major constituent (90%) of strawberry fruit. The strawberry fruit contains 5 percent of total sugar and 0.90 to 1.85% acids the prominent being malic acid, citric acid and succinic acid, with traces of quinic, glyceric, glycolic and oxaloacetic acid. Ripe strawberries attain attractive red colour on sudden rise in temperature after commencement of Fruiting in February. The fruits are delicious and attractive, having pleasant aroma and a delicate flavour.

Commented [DD1]: Bracket is missing

Commented [DD2]: Need to improve as grammatical mistakes are there

In strawberry Ellagic acid is a naturally occurring plant phenol , it has been found to inhibit the cancer disease. It also controls asthma by the regular consumption of his fruits.

The total world area and production of strawberry was 2.6 lakh and 3.61 MT respectively. Europe producing the about 1/3rd of total production strawberries of the world .

Commented [DD3]: Whatever data is mentioned for yield or nutrition composition source need to be mentioned

The quality and yield of fruit depends on different attributes which are closely associated with nutrient uptake by the plant. The supply of nutrients to the plants should be balanced, ensuring not to over or underfertilize. In addition to NPK , micronutrients have a great bearing in influencing the yield attributes and fruit production. Micronutrients are essentially as important as macronutrients to have better growth, yield and quality in plants. In the past, there was no need of micronutrients because these trace elements were naturally supplied by the soil.

The plant growth regulator are being successfully employed for improving germination, rooting, flowering, seed setting, tuberization, fruit setting, inducing parthenocarpy, fruit thinning and various plant growth hormones have been reported to improve fruit quality and of various horticultural and agricultural crops such as Ethrel, cycocel, and NAA, etc. the ripe fruit is ovate in shape and aggregate of achenes or one seeded fruits.

Commented [DD4]: It should be regulators

Thus, this study was undertaken to evaluate the effect of PGR and Micronutrients on Growth, yield and Quality of strawberry.

2. MATERIALS AND METHODS:-

Commented [DD5]: Details of observations/traits to be recorded is missing

2.1 Geographical location of the experimental site

The experimental site is located at a latitude of 25.41° North and longitude of 81.84 °East, with an altitude of 98 meters above the mean sea level(MSL).

2.2 Experimental details

Treatment Combinations

T₀ : Control, T₁ : 50 ppm Gibberellic acid, T₂ : 75 ppm Gibberellic acid, T₃ : 0.4% Zn, T₄ : 0.6% Zn, T₅ : 50 ppm Gibberellic acid+ 0.4% Zn, T₆ : 75 ppm Gibberellic acid + 0.4% Zn, T₇ : 50 ppm Gibberellic acid + 0.6% Zn, T₈ : 75 ppm Gibberellic acid + 0.6% Zn

3.RESULTS AND DISCUSSION

3.1Effect of PGR and micronutrient on plant height (cm), Number of leaves per plant,Leaf area (cm²), Plant Spread of Strawberry cv. Winter Dawn

3.1.1 Plant Height(cm)

At 120 DAP, The maximum plant height (19.00 cm) was observed with treatment T₂ (75 ppm Gibberellic acid) followed by T₈(75 ppm Gibberellic acid + 0.6% Zn) with (15.89cm). Minimum plant height of (10.00 cm) was observed T₀ (control) .

Among the different treatments of application of plant growth regulators and micronutrients the maximum plant height was observed under T₂ (75 ppm Gibberellic acid), while the minimum plant height was recorded under the treatment RDF + Control (T₀).

The increased plant height of strawberry may be due to the increased nitrogen fixation, organic nitrogen utilization, development of root system (**Beer et al., 2017**). Similar results were also reported by **Tripathi et al. (2010)** in strawberry

3.1.2Number of leaves per plant

At 120 DAP, The maximum number of leaves per plant (20.67) was observed with treatment T₂ (75 ppm Gibberellic acid), followed by T₈ (75 ppm Gibberellic acid + 0.6% Zn) with (19.22). Minimum number of leaves per plant (12.56) was observed T₀ (control).

Among the different treatments of soil application of plant growth regulators and micronutrients the maximum number of leaves per plant was observed under T₂ (75 ppm Gibberellic acid), while the minimum number of leaves per plant was observed under the treatment Control (T₀).

Higher number of leaves & leaf area might be due to higher cell division caused by cytokinins & also due to higher supply of assimilate mediated by biofertilizers application. Increased number of leaves might have increased the photosynthetic activity resulting in higher accumulation of carbohydrates. Relatively higher amount of carbohydrate could have promoted the growth rate & in turn increased the berry weight. The findings are in close agreement with the findings of **Dwivedi et al. (1999)** in strawberry.

3.1.3 Leaf area (cm²)

At 120 DAP, the superiority of treatment T₂ (75 ppm Gibberellic acid), registered maximum leaf area (130.33 cm²), which was found at par with T₈ having the leaf area of 128.89 cm². However the minimum leaf area (115.78 cm²) was recorded under the treatment Control (T₀).

Among the different treatments of application of plant growth regulators and micronutrients the maximum leaf area was observed under T2 (75 ppm Gibberellic acid), while the minimum leaf area was recorded under the treatment Control (T₀).

The increase in leaf area due to application of plant growth regulators and micro nutrients has also been reported by Jain *et al.* (2016). Applications in strawberries can increase beneficial microbial populations, which enhance production of plant growth hormones (auxin, gibberellins and cytokinins) and humic acids. Several experiments in strawberry have indicated that these hormones and acids may improve plant growth (leaf area, shoot biomass, number of flowers and runners).

3.1.4. Plant Spread:

At 120 DAP, the superiority of treatment T2 (75 ppm Gibberellic acid), registered maximum plant spread (25.50 cm), which was found at par with T8 having the plant spread of 24.56 cm. However the minimum plant spread (18.22 cm) was recorded under the treatment Control (T₀).

Among the different treatments of application of plant growth regulators and micronutrients the maximum leaf area was observed under T2 (75 ppm Gibberellic acid), while the minimum leaf area was recorded under the treatment Control (T₀).

Beer *et al.*, 2017, reported increase in plant spread might be due to increased growth of plant in the form of height, which accumulated more photosynthesis & thereby increases leaf area per plant. These findings are corroborates with the finding of Umar *et al.* (2009).

Table 1. Effect of PGR and micronutrient on Plant height (cm), Number of leaves per plant, Leaf area (cm²) , Plant Spread of Strawberry cv. Winter Dawn

Treatment	Plant height (cm)	Number of leaves per plant	Leaf area (cm ²)	Plant Spread (cm)
	120 DAS	120DAS	120 DAS	120 DAS
T ₀	10.00	12.56	115.78	18.22
T ₁	14.44	14.11	128.00	19.00
T ₂	19.00	20.67	130.33	25.50
T ₃	15.78	18.11	123.44	19.00

Commented [DD6]: In text it is quoted as 120 DAP and in table it is 120 DAS? Kindly clarify

T ₄	13.00	19.00	119.56	18.78
T ₅	14.00	18.11	128.33	21.44
T ₆	13.56	15.78	122.67	19.22
T ₇	13.44	15.78	124.22	23.89
T ₈	15.89	19.22	128.89	24.56
F test	S	S	S	S
SEd±	0.82	0.88	1.6	0.95
CD at 5%	0.37	0.40	0.72	0.43
CV	7.14	5.51	3.86	4.56

3.2 Effect of PGR and micronutrient on first flower Initiation(days), Number of flowers per plant, Number of fruit per plant, Polar diameter, Radial diameter (mm), Fruit yield (q per ha) of Strawberry cv. Winter Dawn

3.2.1 First flower Initiation(days)

Minimum days taken first flowering and fruiting was recorded with T₂ (75 ppm Gibberellic acid), (59.17). Closely followed by T₈ (60.44). Maximum days taken for first flowering and fruiting was recorded in T₆ (65.44).

These outcomes have got the support of the findings of Kumar et al., (2015) who also recorded earliest flowering and fruiting with the application of cent per cent PGR and organic fertilizers, Shukla *et al.*, (2009), who recorded earliest flowering. Tripathi *et al.* (2015) also recorded advancement in flowering. Number of days required minimum to produce first flower might be due to balanced supply of vermi-compost & bio-fertilizers, which provides all the necessary elements to plant to become early flowering and fruiting.

3.2.2 Number of flowers per plant

The Maximum number of flowers per plant was recorded with T₂ (75 ppm Gibberellic acid) 32.00. Closely followed by T₈ (25.67). Minimum number of flowers per plant was recorded in T₀ control (17.11).

This might be due to the effect of plant growth regulators and micronutrients obtained from PGR the cell elongation and division. Soluble PGR increases photosynthetic activities of leaves, which leading to development of primary flowers, production of viable flowers. Similar results were also obtained by **Neetu & Sharma *et al.* (2018)** **Tripathi *et al.* (2015)** and **Singh *et al.* (2015)** in strawberry.

Commented [DD7]: This needs to be rewritten

Commented [DD8]: Wrongly cited

3.2.3 Number of fruit per plant

The Maximum number of fruits per plant and was recorded with T2 (75 ppm Gibberellic acid) with 20.44. Closely followed by T8 (18.67). Minimum number of fruits per plant was recorded in T3 control (14.11).

The results are consistent with those of Singh *et al.* (2006) in strawberry, and they are similar to those of Al-Madhagiet *al.* (2012). Nor *et al.* (2014) also reported increased fruit volume with the application of PGR and micronutrients in strawberry.

3.2.4 Polar diameter

The Maximum polar diameter of fruit was recorded with T2(75 ppm Gibberellic acid) 48.33 mm closely followed by T5 (48.33 mm). Minimum polar diameter of fruit was recorded in T0 control (29.44 mm).

3.2.5 Radial diameter (mm)

The Maximum radial diameter of fruit was recorded with T2 (75 ppm Gibberellic acid) 29.83 mm closely followed by T8 (28.33 mm). Minimum polar diameter of fruit was recorded in T0 control (7.44 mm).

3.2.6 Fruit yield (q per ha)

The Maximum fruit yield (q per ha) was recorded with T2 (75 ppm Gibberellic acid) 9.21 closely followed by T8 (9.1). Minimum fruit yield (q per h a) was recorded in T0 control (6.10).

The results are in line with **Nowsheen *et al.*, 2006** and **Umar *et al.*, 2009** in the strawberry who described maximum yield in the strawberry. Increment in the number of runners per plant may be because of enhanced growth of vegetative and leaves number that gathered more photo- synthates and hence increased runners and the leaf area per plant. PGR and micronutrients might have an effect on runner production because of its ability to

increase the nutrient availability in the soil as well as provide better environment for better and healthy root development.

Table 2. Effect of PGR and micronutrient on first flower Initiation(days), Number of flowers per plant, Number of fruit per plant, Polar diameter, Radial diameter (mm), Fruit yield (q per ha) of Strawberry cv. Winter Dawn

Treatment	first flower Initiation(days)	Number of flowers per plant	Number of fruit per plant	Polar diameter	Radial diameter (mm)	Fruit yield (q per ha)
T ₀	64.55	17.11	15.00	29.44	7.44	6.1
T ₁	63.44	22.44	18.44	43.33	15.33	7.21
T ₂	59.17	32	20.44	48.33	29.83	9.21
T ₃	61.33	21.78	14.11	35.00	15.56	6.66
T ₄	62.55	22.11	17.56	43.33	17.00	7.32
T ₅	60.66	23.11	16.56	48.33	27.78	8.62
T ₆	65.44	19.78	16.78	32.78	16.67	7.91
T ₇	62.33	17.33	17.11	31.67	17.00	8.22
T ₈	60.44	25.67	18.67	47.22	28.33	9.1
F test	S	S	S	S	S	S
SEd±	0.68	1.51	0.64	2.56	2.51	0.36
CD at 5%	0.31	0.68	0.29	1.15	1.13	0.16
CV	3.26	5.64	6.14	19.66	8.67	3.80

3.3. CONCLUSION

On the basis of our experiment findings it is concluded that the treatment T₂(Gibberellic acid - 75ppm) was found to be best in the term of vegetative parameter, flowering, fruiting, yield and quality of strawberry.

References:-

Agafonov, N.V., Blinovski, I.K. and Solovei, E.P.(1978). The preparation jury as strawberry. Sodobodstvo (Russian).

Ahire, Dipakkumar. (2017). Effect of PGR on growth, yield and quality of strawberry (Fragaria x ananassa Duch) cv. Sweet Charlie. 10. 8817-8819.

Asrey, R.,Jain, O.K. and Singh, R.(2000-01).Effect of plant growth regulators on growth and survival of strawberry runners under semi arid region of Punjab. Indian J. Plant Physiology

Asrey , Ram., R.K. Jain and Singh, R. (2003). Effect of plant growth regulators on growth and survival of strawberry runners under semi arid region of Punjab, Indian J. Plant Physiology.

Beer, Karma &Syamal, Alok. (2017). Effect of Organic, Inorganic and Bio-Fertilizer on Growth, Flowering, Yield and Quality of Strawberry (Fragaria × Ananassa Duch.) cv. Chandler. International Journal of Current Microbiology and Applied Sciences. 6. 2932-2939. 10.20546/ijemas.2017.605.332.

Commented [DD9]: Wrongly cited reference

Brahamachari, V.S. and Rani, R.(2001) Effect of growth substances on productivity, cracking, repining and quality of fruits in litchi.

Brahamachari, V.S., Mandal, A.K., Kumar, R. and Rani, R.(1995). Effect of growth substances of fruit set and physico- chemical characteristics of sardar guava.

Commented [DD10]: Guava?

Dwivedi, M.P.(1987). Effect of photoperiod and growth regulators on vegetative growth, flowering and yield of strawberry. Ph.D. thesis, Dr. Y.S Parmar University of Horticulture and forestry, Solan , India

Dwivedi, M.P., Negi , K.S. and Rana, H.S.(1994). Influence of photoperiod and bioregulators on vegetative growth of strawberry under controlled condition.

Dwivedi, M.P., Negi, K.S., Jindal, K.K. and Rana, H.S.(2000). Effect of bioregulators on vegetative growth of strawberry.

Foda, S.A., Nassar, H.H. and Mansoor, S.A.(1979). Effect of some growth regulators on runner production and yield of strawberry.

Gujar, S. & Pathade, A. & Pathade, G. (2024). Impact of Plant Growth Regulators on Strawberry Plant – A Review. *Ecology, Environment and Conservation*. 30. 252-257. [10.53550/EEC.2024.v30i02s.053](https://doi.org/10.53550/EEC.2024.v30i02s.053).

Jain, N., Mani, A., Kumari, S., Kaseera, S., & Bahadur, V. (2017). Influence of INM on yield, quality, shelf life and economics of cultivation of strawberry (*Fragaria × ananassa* Duch.) cv. Sweet Charlie. *Journal of Pharmacognosy and Phytochemistry*, 6(5), 1178-1181

Commented [DD11]: Year cited in text and in reference are different

Katel, S., Mandal, H., Kattel, S., Yadav, S. P., & Sharma L. B. (2022). Impacts of plant growth regulators in strawberry plant: A review. *Heliyon*. 8. e11959. [10.1016/j.heliyon.2022.e11959](https://doi.org/10.1016/j.heliyon.2022.e11959).

Khokhar, U.U., Prasad, J. and Sharma, M.K. (2004).Influence of growth regulator on yield and quality of strawberry cv. Chandler Haryana, *J. Horti. Science*

Kumar, R., Bakshi, M. and Singh, D.B. (2012). Influence of plant growth regulators on growth, yield and quality of strawberry (*Fragaria x Ananassa Duch.*) under U.P. Subtropics, *Asian J. Hort.*, 7 (2) : 434-436.

Kumar, R., Bakshi, M., & Singh, D. B. (2012). Influence of plant growth regulators on growth, yield and quality of strawberry (*Fragaria x ananassa* Duch.) under UP sub tropics. *Asian J. Hortic*, 7, 434-436.

Kumar, N., Singh, H. K., & Mishra, P. K. (2015). Impact of organic manures and biofertilizers on growth and quality parameters of Strawberry cv. Chandler. *Indian Journal of Science and Technology*, 8(15), 1-6.

Kumar, R., Bakshi, M., & Singh, D. B. (2012). Influence of plant growth regulators on growth, yield and quality of strawberry (*Fragaria x ananassa* Duch.) under UP sub tropics. *Asian J. Hortic*, 7, 434-436.

Commented [DD12]: This same reference repeated thrice and no where cited in text

Kushwah, M. S., Singh, D., Singh, S., & Bairwa, M. (2018). Effect of integrated nutrient management on growth, yield and quality traits of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler. *Journal of Pharmacognosy and Phytochemistry* 2018; SP1: 712-715

Massetani, F. and Neri, D. (2016). Strawberry plant architecture in different cultivation systems. *Acta Hort.* 1117, 291-296.

Marwaha, Himansh & Singh, Lakhwinder &Kachawaya, Dilip. (2023). Effect of Bio-fertilizers and Plant Growth Regulators on Growth, Yield, Quality and Economics of Strawberry (*Fragaria × ananassa* Duch.) Cv. Camarosa. *Biological Trace Element Research.* 15 (15a). 345.

Mehraj, H. & Reza, Mamunur& Islam, Mohidul& Hoque, Md. Azizul & Sikder, Ripon & Uddin, Dr. (2015). Influence of different GA3 concentrations on growth and yield of broccoli. 10. 332-335. 10.5829/idosi.aejsr.2015.10.5.12663.

Palei, Suvalaxmi& Kumar, Arun & Sahoo, Ajit & Dash, Kumar & Swain, Saudamini. (2016). INFLUENCE OF PLANT GROWTH REGULATORS ON STRAWBERRY (*FRAGARIA × ANANASSA*) Cv. CHANDLER UNDER ODISHA CONDITION. 7. 9945-9948.

Rathod, K. D., Ahlawat, T. R., Kumar, S., Sarkar, M., & Chakraborty, B. (2021). Effect of Plant Growth Regulators on Growth, Yield and Quality of Strawberry (*Fragaria × ananassa* Duch.) cv. Winter Dawn under Open Field Conditions of South Gujarat. *Agricultural Science Digest*, 41(2), 329-333.

Reddy, G.C., Goyal, R.K., &Godara, A.K. (2023). Use of organic manures in combination with Azotobacter in strawberry: effect on growth, yield and quality attributes. *Journal of Plant Nutrition*, 46, 4188 - 4198.

Rustam, Chovatia, R.S. and Makhmale, S.J. (2017). Effect of GA3, urea and ZnSO4 on growth and yield parameters of strawberry (*Fragaria x ananassa*Duch.) cv. SWEET CHARLIE under protected condition. *Adv. Res. J. Crop Improv.*, 8 (1) : 70-74,

Sharma, R. and Singh, R. (2009). Gibberellic acid influences the production of malformed and button berries, and fruit yield and quality in strawberry (*Fragaria×ananassa* Duch.). *Scientia Horticulturae.* 119. 430-433. 10.1016/j.scienta.2008.11.002.

Ughareja, A. and Pandya, Y. (2023) Studies on the Effect of Plant Growth Regulators on Growth and Yield of Strawberry cv. (Winter Dawn) Under Protected

Cultivation International Journal of Agriculture and Animal Production ISSN 2799-0907
Vol : 03 , No. 02.

Verma, Neelendra Singh &Kv, Ramana & Rao, & Rajwade, Yogesh & Yadav, Deepika & Trivedi, Ayushi & Rao, Ramana. (2023). Growth and yield of strawberry (Fragaria x ananassa Duch) under different mulches in vertisols of Madhya Pradesh. The Pharma Innovation. 12. 1324-1327.

Yadav, A., Pratap, B., Shivam, A. K., & Patro, A. (2018). Assess the effect of micronutrients and plant growth regulators on quality parameters of strawberry cv. Chandler. The Pharma Innovation J, 7(1), 303-305.

Commented [DD13]: Out of all the references only two are cited in the text. And reference cited in text are not in references section?

UNDER PEER REVIEW