

## Destabilization of Abnormal Methylation Enzymes ~~as~~ The Only Viable Option for the Elimination of Cancer Stem Cells to Save Cancer Patients.

### ABSTRACT

Cytotoxic chemotherapy was a tragic by-product of World War II. It was the major therapeutic modality employed ~~during 1971-1976 declared by~~ ~~but~~ President Nixon ~~but~~ failed to win the War on Cancer. ~~during 1971-1976 declared by President Nixon~~. Despite the failure, killing of cancer cells (CCs) remained the commanding principle of cancer ~~institutions establishments~~ to guide development of cancer therapies throughout. Apparently, this commanding principle is incorrect, since cancer mortality continues to increase.

Cancer evolves due to wound unhealing. Wound healing requires the proliferation and the terminal differentiation of progenitor stem cells (PSCs). PSCs are cells with abnormal methylation enzymes (MEs), which allow cells to gain a great advantage on cell growth. The nature creates chemo-surveillance made up by wound healing metabolites, differentiation inducers (DIs) and differentiation helper inducers (DHIs), as a protection mechanism to keep cells with abnormal MEs under control. Wound unhealing is due to the collapse of chemo-surveillance resulting in the failure to induce terminal differentiation of PSCs. ~~The nature~~ Nature does not have a mechanism to detect the collapse of chemo-surveillance to make the correction. PSCs are forced to proliferate, which is limited by contact inhibition, another protection mechanism ~~the nature~~ nature created to limit the buildup of normal stem cells. PSCs are then forced to evolve into cancer stem cells to escape contact inhibition. It takes a single hit to silence ~~TET-1~~ enzyme, an enzyme responsible to direct lineage transitions, to turn PSCs into cancer stem cells (CSCs). The evolution of CSCs from PSCs is quite easy, since PSCs are equipped with very active abnormal MEs. The propagation of CSCs still cannot heal the wound, because the problem is the collapse of chemo-surveillance. The CSCs will be forced to progress to faster growing CCs by chromosomal abnormalities such as translocations to activate oncogenes, or deletions to inactivate suppressor genes. So, the correct solution ~~ef~~to cancer is to restore chemo-surveillance by providing DIs and DHIs to destabilize abnormal MEs to achieve terminal differentiation of PSCs and CSCs which are closely linked to wound unhealing. Killing of CSCs and PSCs cannot heal the wound. Therefore, destabilization of abnormal MEs to achieve terminal differentiation of CSCs is the only viable option to take out CSCs by healing the wound to save cancer patients. The elimination of CSCs is essential to the success of cancer therapy. We have ~~carried out~~ conducted extensive studies of natural and unnatural DIs and DHIs for the manufacturing of ~~CDA~~ formulations to save cancer patients.

**Keywords:** Cancer therapies; CCs; CSCs; DIs; DHIs; MEs; PSCs.

**Comment [T1]:** Please organize the abstract with Aim of the Study, Methods, Results and Conclusion.

**Formatted:** Font: Bold

**Formatted:** Font: Bold

**Comment [T2]:** Please spell out the acronym first time it is mentioned in the manuscript.

**Comment [T3]:** Please spell out the acronym first time it is mentioned in the manuscript

**Formatted:** Font: Bold, Italic

**Formatted:** Font: Bold, Italic

## 1. INTRODUCTION

**Comment [T4]:** Please use spell check throughout the manuscript.

**Formatted:** Font: Bold

Cancer therapy got to a bad start ~~when it relied to rely~~ on cytotoxic chemicals to kill CCs. It was a tragic by-product of World War II. During the war, toxic sulfur mustard gas bombs were used. Victims of toxic gas all displayed depletion of leukocytes in their blood specimens, which inspired oncologists to use toxic chemicals to treat leukemia patients. Cytotoxic drugs became the standard care of cancer therapy, and the disappearance of cancer cells or tumor became the standard criteria for the evaluation of the efficacy of cancer therapy. Cytotoxic drugs and radiation were the major therapeutic modalities employed during the War on Cancer declared by President ~~Nixon during in~~ 1971-1976, which was not successful [1]. When therapeutic modalities were drilled through as a presidential project that received unlimited support from national resources but failed to achieve the goal, it was fair to conclude that the therapeutic modalities employed were not good for cancer therapy. Apparently, cancer ~~institutions establishments~~ agreed to this conclusion and tried to find replacements. They turned to gene and targeted therapies ~~during from~~ 1976 to 1996, and then to anti-angiogenesis during 1996-2016 and now to immunotherapy from 2016 onwards [2]. They wasted ~~dtwenty 20~~ years to learn the difficulties of gene ~~therapy, and therapy and~~ wasted ~~d~~ another ~~twenty 20~~ years on ~~the~~ unsuccessful attempts of developing anti-angiogenesis therapy. Can they succeed on immunotherapy? Immunotherapy definitely is a better version than cytotoxic chemotherapy and radiotherapy. The therapeutic target of cell surface antigens after all is a specific cancer target to reduce adverse effects. But immunotherapy has the same ~~draw back drawback~~ as the cytotoxic chemotherapy to show ineffectiveness against CSCs and to cause damage to chemo-surveillance, which are the reason to contribute to the failure of winning the war on cancer. The effect of immunotherapy to cause the damage of chemo-surveillance is much worse than cytotoxic ~~chemotherapy, because chemotherapy because~~ immunological response triggering the production of tumor necrosis factor (TNF) is the primary reason to cause the damage of chemo-surveillance. It is doubtful that immunotherapy can save cancer patients better than cytotoxic chemotherapy. Since cancer establishments have not found acceptable replacements, they stayed on to use failed drugs to treat cancer patients. The outcome is expectable that cancer mortality keeps on increasing. It has reached 10 million annual deaths in 2019 worldwide with an annual increment of 5% as predicted by the NCI experts [3]. The mistake of cytotoxic chemotherapy was made at a time when we did not have complete information of cancer. But now we have better knowledge of cancer, we should not make the same mistake in selecting drugs that can only kill CCs. A perfect cancer drug must be able to take out both CCs and CSCs, and to restore the functionality of chemo-surveillance [4]. Cancer evolves from wound unhealing due to the collapse of chemo-surveillance [5-11]. Healthy people tend to produce enough metabolites active as DIs and DHIs to ensure destabilization of abnormal MEs of PSCs to achieve terminal differentiation [12, 13]. DIs are chemicals capable of eliminating telomerase from abnormal MEs and DHIs are inhibitors of MEs capable of potentiating the activity of DIs. If chemo-surveillance is damaged due to pathological conditions that create cachexia symptoms, then wound cannot be healed to result in the evolution of CSCs from PSCs in order to escape contact inhibition which limits the extent of the proliferation of PSCs. It takes a single hit to silence TET-1 enzyme [14, 15], which is the enzyme responsible to direct lineage transitions, for

the conversion of PSCs to become CSCs, a task that can be easily accomplished by PSCs since these cells are equipped with exceptionally active MEs. The propagation of CSCs is still unable to heal the wound, because the problem is the collapse of chemo-surveillance which ~~the nature~~ nature does not have a mechanism to correct. Chromosomal abnormalities then set in to activate oncogenes by translocations or to inactivate suppressor genes by deletions to turn CSCs to faster growing CCs. Evidently, ~~damages~~ damage created by the wound are responsible for the evolution of cancer. Killing of CCs creates more wounds to aggravate the already ~~bad situation~~ dire situation and to promote the proliferation of CSCs to work on the damages cytotoxic drugs created [16]. Eventually, the proportion of CSCs will increase from ~~less~~ less than 2% in the primary cancer which respond well to cytotoxic chemotherapy to reach more than 10 % like that of the primary brain cancer which is unresponsive to cytotoxic chemotherapy [17, 18]. CSCs are protected by drug resistance and anti-apoptosis mechanisms, and, therefore, are unresponsive to cytotoxic chemotherapy and radiotherapy [19-22]. Solution of CSCs is very difficult with cytotoxic approaches. Yet, the solution of CSCs is essential to save cancer patients [23]. Since CSCs are critically linked to wound unhealing, destabilization of abnormal MEs is the only option to solve CSCs [24, 25]. The biological mission of CSCs is wound healing just like that of PSCs. Wound healing metabolites are the partners of their biological mission, and, therefore, can easily access CSCs to achieve induction of terminal differentiation. Cancer establishments are focused on the development of drugs to kill CCs, none of which can affect CSCs. Cancer establishments are not very nice to put up toxic drugs that cause so many cancer deaths and to block the development of CDA formulations that can solve CSCs to reduce cancer deaths [26]. They should be removed to save cancer patients [27, 28]. We count on supreme authorities of King Charles of England and President Biden of USA to rectify cancer therapies to save cancer patients, King Charles to save himself and President Biden of to achieve his cancer moonshot initiative.

## 2. COMMENTARIES AND DISCUSSION

### 2-1. Cancer Evolves as A Consequence of Wound Unhealing

The concept of cancer evolve as a consequence of wound unhealing was first introduced by the great German scientist Virchow in the 19<sup>th</sup> century [5]. It was again brought up by Dvorak in 1986 [6]. The close relationship between cancer and wound healing was noticed by MacCarthy-Morrrough and Martin [7]. We provided the most important details on this subject that included abnormal MEs to promote perpetual proliferation of CCs [29-31]; chemo-surveillance as the nature's creation of allosteric regulation on abnormal MEs to ensure perfection of wound healing to avoid disastrous consequence of wound unhealing [12, 13, 32-34]; DIs and DHIs as wound healing metabolites and also as active players of chemo-surveillance [12, 13, 32-34]; hypomethylation of nucleic acids as a critical mechanism on the induction of terminal differentiation of cells with abnormal MEs [35]; mechanism of wound healing to involve the proliferation and the terminal differentiation of PSCs [8-11]; and the evolution of CSCs from PSCs through a single hit to silence TET-1 enzyme [14, 15, 36]. These studies strongly support

Formatted: Font: 12 pt, Bold

Formatted: Font: Bold

Formatted: Font: 12 pt, Italic

Formatted: Font: Italic

the concept that cancer evolves as a consequence of wound unhealing. Our carcinogenesis studies also confirm the validity of this concept. During the challenges with hepatocarcinogens, we noticed the appearance of numerous tiny preneoplastic hyperplastic nodules before the appearance of ~~large size~~enormous size carcinomas, which displayed abnormal MEs [37]. These tiny hyperplastic nodules must represent the proliferation of PSCs in the process of active wound healing. Most of these tiny nodules disappeared shortly afterward, indicating completion of wound healing, and only those which did not disappear later developed to become ~~large size~~enormous size carcinomas. During the challenges with hepatocarcinogens, if the animals were provided Antineoplaston 10, which ~~is was~~ phenylacetylglutamine ~~that is~~ effective as anti-cachexia chemical [12], ~~then~~ the development of ~~hepatocarcinomas~~hepatocarcinoma could be prevented [38]. Therefore, cancer evolves as a consequence of wound unhealing due to the collapse of chemo-surveillance is a valid concept. Cancer therapy based on pro-wound healing approach we advocate is, therefore, the most appropriate modality of cancer therapy [8, 13, 16, 24, 25, 34, 39, 40]. We have attributed the success of extremely difficult moon shot project of President Kennedy to the assumption of right approach. ~~A right~~The right approach is the magic code to ~~the success~~success [41]. The commanding principle of killing CCs preferred by cancer establishments is an anti-wound healing approach, clearly a wrong approach for cancer therapy. No wonder it has not been successful,~~or even~~ supported as a presidential project.

2-2. CDA-2*(Cell Differentiation Agent-2)* as A Persuasive Good Cancer Drug for the Elimination of Cancer Stem Cells

~~Myelodyaplasi~~Myelodysplastic syndromes (MDS) are a classic case of cancer with a clear connection to wound unhealing, and CDA-2 is a preparation of wound healing metabolites as the drug of choice for the therapy of MDS. MDS often ~~start~~starts with a display of an immunological disorder [42], which prompts the production of inflammatory cytokines. Among cytokines produced TNF is a critical factor related to the development of MDS [43]. It causes excessive apoptosis of bone marrow stem cells, thus severely ~~affect~~affecting the ability of the patient to produce hematopoietic cells such as erythrocytes, platelets or neutrophils. TNF is also responsible for the collapse of chemo-surveillance due to its effect to cause blood vessel hyperpermeability [44, 45], resulting in the excessive urinary excretion of low molecular weight metabolites. Wound healing metabolites are among low molecular weight metabolite excreted to result in the collapse of chemo-surveillance, which ~~the~~ nature does not have a mechanism to rectify. The evolution of cancer starts from the collapse of chemo-surveillance. The ~~high~~high ~~level~~elevated level of telomerase expression in the peripheral and bone marrow leukocytes in MDS patients is an indication of the widespread multiplication of CSCs evolving from PSCs [46, 47]. The expansion of PSCs is limited by contact inhibition. MDS are diseases at the stage of CSCs evolving from PSCs. The propagating pathological cells of MDS have been identified as human CSCs [48]. So, MDS are diseases attributable entirely to the propagation of CSCs. These diseases are ideal for the development of drugs effective against CSCs. So far, Vidaza, Decitabine and CDA-2 are the three drugs approved for MDS therapy by the Chinese FDA. Vidaza and Decitabine are also approved for the therapy of MDS by the US FDA. Professor Jun Ma, the Director of Harbin Institute of Hematology and Oncology, is instrumental in conducting clinical

**Comment [T5]:** Please spell out the acronym.

**Formatted:** Font: 12 pt, Italic

**Formatted:** Font: Italic

**Formatted:** Font: 12 pt, Italic

**Formatted:** Font: Italic

trials of these three MDS drugs for the approval by the Chinese FDA. According to his assessments based on two cycles of treatment protocols, CDA-2 had a noticeable better therapeutic efficacy, although However, it is slower to achieve complete remission, based on cytological evaluation, and evaluation and marked better therapeutic efficacy based on hematological improvement evaluation. This which was based on the dependence of blood transfusion to stay healthy as shown in Figure 1, which is reproduced from the reference [49]. All three drugs achieve MDS therapy by the inactivation of abnormal MEs.




Figure 1

**Fig. 1. Relative effectiveness of MDS drugs**

Formatted: Font: 12 pt, Bold

Formatted: Font: Bold

Vidaza and Decitabine by the covalent bond formation of methyltransferase and the 5'-azacytosine base incorporated into DNA [50], whereas CDA-2 by the elimination of telomerase of abnormal MEs [51]. The action of CDA-2 is selective on the tumor factor of abnormal MEs, whereas the action of Vidaza and Decitabine is non-selective to affect MEs of all stem cells. Thus, CDA-2 is without adverse effects, whereas Vidaza and Decitabine are known carcinogens [52, 53], and very toxic to DNA [54-56]. Obviously, CDA-2 is the drug of choice for the therapy of MDS with a better therapeutic efficacy and devoid of adverse effects. Induction of terminal differentiation of CSCs like the induction of terminal differentiation of PSCs in wound healing is obviously the only option to solve CSCs. Killing of CSCs cannot heal MDS. So that destabilization of abnormal MEs is the only viable option for the elimination of CSCs to save cancer patients. The elimination of CSCs is essential to save cancer patients [23]. Cancer establishments blocked Antineoplastons, which were preparation similar to CDA-2. Apparently, these preparations are the only viable option for the elimination of CSCs to save cancer patients. Killing of CSCs and CCs favored by cancer establishments cannot solve the issue related to CSCs.

### 2-3. Abnormal MEs as the Most Critical Issue of Cancer

Formatted: Font: 12 pt, Italic

Formatted: Font: Italic

Perpetual cell proliferation is the most outstanding feature of cancer. Obviously, cancer is basically a problem of growth regulation going awry. MEs play a pivotal role in the regulation of cell replication and differentiation, and, therefore, are closely related to the issue of

cancer[50, 56, 57]. Since MEs play such ~~an important role~~ ~~a significant role~~ ~~in~~ the regulation of cell growth, they are subjected to exceptional double allosteric regulations: one on the individual enzymes and the other on the enzyme complex [32]. Enzymes ~~involve~~ ~~involved~~ in important biological regulation are often subjected to delicate regulations. Allosteric regulation is the most pervasive regulation to maintain biological equilibrium to avoid extreme often to result in creating clinical symptoms. MEs are a ternary enzyme complex consisting of methionine adenosyltransferase (MAT)-methyltransferase (MT)-S-adenosylhomocysteine hydrolase (SAHH) [58]. In steroid hormone target organs, SAHH is the steroid hormone receptor. SAHH requires a steroid hormone to assume a stable configuration in order to form a dimeric complex with MT. SAHH-MT dimer has a molecular size similar to MAT. These three ~~enzymes~~ ~~are enzymes~~ then form a ternary enzyme complex, which is the stable and active functional unit of MEs. In the absence of steroid ~~hormone~~ ~~hormones~~, the ternary enzyme complex dissociates into individual enzymes to lose activity. MT in the monomeric form ~~has a tendency to~~ ~~tends to~~ be modified to become nuclease which can cause damages to trigger apoptosis, resulting in the involution of the steroid hormone target organs. In telomerase expressing cells such as embryonic stem cells (ESCs) and PSCs which are also ESCs, MEs are associated with telomerase. The association changes kinetic properties of MEs and the regulation greatly in favor of cell growth.  $K_m$  values of telomerase associated MAT-SAHH isozyme pair are 7-fold higher than those of normal isozymes [28-30]. The increased  $K_m$  values indicate that abnormal MEs hold a higher level of S-adenosylmethionine (AdoMet) and S-adenosylhomocysteine (AdoHcy), which are important for the stability of MEs to promote growth of cells with abnormal MEs as Prudova et al. found AdoMet could protect protein from protease digestion [59], and Chiba et al. found when HL-60 cells were induced to undergo terminal differentiation the pool sizes of AdoMet and AdoHcy shrunk greatly [60]. Obviously, abnormal MEs play ~~an important role~~ ~~a vital role~~ ~~to promote~~ ~~in promoting~~ the growth of cells with abnormal MEs. Normal stem cells with abnormal MEs do not seem to cause problems, because normal stem cells are protected by safety mechanisms such as contact inhibition, TET-1 enzyme to direct lineage transitions and chemo-surveillance as a brake to prevent the buildup of cells with abnormal MEs. Evidently, abnormal MEs are important for the normal functions of cells with abnormal MEs. Premature interruption of the normal functioning of abnormal MEs, e.g. the application of thalidomide, is detrimental, resulting in malformation of body ~~parts~~ ~~parts~~, notably limbs. It is a horror ~~balance,~~ ~~balance~~; any misstep will result in disastrous consequences. There is no room for ~~mistake~~ ~~mistakes~~. We have to solve cancer very precisely!

Both abnormal MEs and chromosomal abnormalities for the activation of oncogenes or inactivation of suppressor genes are very critical issues of cancer. Abnormal MEs are responsible for the blockade of differentiation for the proliferation to ~~carry on~~ ~~continue~~, and chromosomal abnormalities are responsible to promote proliferation. Which is more important, the blockade of differentiation or the activation of proliferation? Most people bet on the activation of proliferation, including cancer establishments. They are wrong. Activation of proliferation happens quite late in the carcinogenesis process. There are multiple ways to activate proliferation. One way is solved. There may soon pop up another way to negate the previous effort. It is an endless struggle to put out activation of proliferation. The cancer establishments tried to put out activation of proliferation during 1976-1996. They learned the

difficult difficulty of gene therapy, and gave up. Destabilization of abnormal MEs is the best solution. Abnormal MEs happen on PSCs, the precursors of CSCs, and pass on to CSCs and CCs. Abnormal MEs are universal to all human cancers [36]. A stroke to solve abnormal MEs, MEs can also put to rest chromosomal abnormalities to activate proliferation. After all, oncogenes and suppressor genes are cell cycle regulatory genes. They have important roles to play when cells are in cell cycle replicating. But if replicating cells exit cell cycle to undergo terminal differentiation, they have no roles to play. So, obviously, abnormal MEs are the bullseye of cancer therapy. Destabilization of abnormal MEs is the only option to solve CSCs, and the solution of CSCs is essential to save cancer patients. Cancer establishments must approve CDA formulations for the sake of saving cancer patients, which is also their goal.

#### 2-4. Development of CDA Formulations for the Solution of CSCs to Save Cancer Patients

We have carried out conducted extensive studies of natural and unnatural DIs and DHIs for the manufacture of CDA formulations [2, 8, 23-25, 34, 36, 39, 51, 46, 58, 63-72]. Our findings of effective DIs and DHIs are summarized in Table 1 and 2. ATRA is the standard care of acute

**Table 1. Effective DIs**

DIs	ED <sub>25</sub> (μM)	ED <sub>50</sub> (μM)	ED <sub>75</sub> (μM)	
	ATRA	0.180.36	0.75	
	PGJ27.9	13.8	20.5	
PGE2	20.6	32.0	46.5	
	DicycloPGE221.0		43.5-	
AA	21.0	42.0	-	
BIBR1532	32.3	43.7	55.1	
	Boline	60.1	78.8	94.2

promyelocytic leukemia [73]. It requires the expression of the receptor of ATRA to activate oligoadenylate synthetase to achieve the therapeutic effect. The product of this enzyme, oligoadenylate, is the actual DI [74]. The therapeutic effect of ATRA on promyelocytic leukemia is excellent, achieving 90% complete remission by a single agent, almost reaching the goal of 100% complete remission by a single agent demanded by cancer establishments. DI alone cannot achieve 100% remission. DI alone creates damages damage, e.g. nuclease attack on the replicating DNA, to interrupt the completion of differentiation. The damages damage can be repaired to resume malignant growth. Therefore, the complete remission achieved by ATRA alone cannot last very long. But if a DHI is applied with DI, that can prevent the dissociation of MT-SAHH dimer, then the possibility of MT monomer to become nuclease to cause the problem of recurrence can be greatly reduced. Therefore, it is a good policy to use DI and DHI to achieve good therapy for cancer. Multiple agents are good for the completion of cancer therapy. Arsenic acid is a DHI, which can prevent recurrence of ATRA alone. Arsenic acid is very toxic. We have found excellent DHIs presented in Table 2 which are nontoxic to function as

Formatted: Font: 12 pt, Italic

Formatted: Font: Italic

Comment [T6]: Please spell out the acronym first time it is mentioned in the manuscript

Formatted: Font: 12 pt, Bold

Formatted: Font: Bold

DHI to prevent the recurrence of ATRA alone. The rest of DHIs presented in Table 1 work directly on abnormal MEs. AA and its metabolites PG derivatives are natural DHIs involved in chemo-surveillance. PG derivatives are approved drugs for the delivery. BIBR1532 and boldine are approved cancer drugs as telomerase inhibitors. Application of change of indication of the approved drugs should not take as long as the application for the approval of new indication.

**Table 2. Effective DHIs**

Signal Transduction			
SAHH Inhibitors	RI <sub>0.5</sub> (μM)	Inhibitors	RI <sub>0.5</sub> (μM)
Pyrivinium Pamoate	0.012	Sutent	0.28
Vitamin D <sub>3</sub>	0.61	Berberine	1.62
Dexamethasone	0.75	Vorient	10.1
Beta-Sitosterol	1.72	Gleevec	11.9
Dihydroepiandrosterone	1.79	Selenite	19.7
Prenisolone	2.22		
Hydrocortisone	4.59	Polyphenols	RI <sub>0.5</sub> (μM)
Pregnenolone	7.16		
		Tannic Acid	0.37
MT Inhibitors	RI <sub>0.5</sub> (μM)	EGCG	0.62
		Resveratrol	1.16
Uroerythrin	1.9	Curcumin	1.24
Hycanthone	2.1	Kuromanine	1.43
Riboflavin	2.9	Coumestrol	1.95
		Genisteine	2.19
MAT Inhibitors	RI <sub>0.5</sub> (μM)	Pyrogallol	3.18
		Silibinin	3.80
Indol Acetic Acid	220	Caffeic Acid	3.87
Phenylacetylvaline	500	Ellagic Acid	4.45
Phenylacetyllecucine	780	Gallic Acid	5.35
Butyric Acid	850	Ferulic Acid	7.41
Phenylbutyric Acid	970	Phloroglucinol	38.82

As shown in Table 2, SAHH and MT inhibitors are much better DHIs than MAT inhibitors. MAT is the most stable enzyme of the three MEs. The association with telomerase further increases its stability. Therefore, it is not easy to shake loose of this enzyme. Pregnenolone is a major DHI of CDA-2. Apparently, pregnenolone is an important player of chemo-surveillance. It is the masterprimary substrate of steroid metabolites to have a great influence a profound influence on growth regulation. The production of pregnenolone is bell shape in relation to age with a peak daily production of around 50 mg at 20-25 years old [75]. The youngest and the oldest people produce relatively smaller amounts, and these are the two age groups most vulnerable to develop to cancer. Pregnenolone is a single metabolite to have a great influence a major influence on the evolution of cancer. It is our top choice of natural DHI. The finding of

Formatted: Font: 12 pt, Bold

Formatted: Font: Bold

inhibitors of signal transduction as excellent DHIs is not a ~~surprise, since~~ surprise since signal transductions produce factors to promote cell growth. Gleevec is the standard care of chronic myeloid leukemia and GIST [76]. Thus, DIs and DHIs can be very effective cancer drugs. The finding of polyphenols as excellent DHIs is a ~~surprise, but~~ surprise but is a good surprise a pleasant surprise since polyphenols are considered good for health, and greatly promoted as health food. The finding of polyphenols as excellent DHIs increases their credibility as ~~health~~ healthy food.

Effective CDA formulations can be made by DIs and DHIs with different formulations as  $ED_{25}$  of a DI +  $3xRI_{0.5}$  of a DHI, or  $ED_{50}$  of a DI +  $2xRI_{0.5}$  of a DHI, or  $ED_{75}$  of a DI +  $RI_{0.5}$  of a DHI [69]. We have provided these data in Table 1 and 2.  $RI_{0.5}$  of a DHI is equivalent to  $ED_{25}$  of a DI, which can be determined by the procedure previously described [67]. In the design of CDA formulations, we must take into ~~considerations of~~ consideration non-cancer issues such as blood brain barrier of brain cancer, collagen envelop of pancreatic cancer and hypoxia of melanoma to select DIs and DHIs to overcome non-cancer issues. It is very important too that we ~~have to come up~~ come up with easy procedures for the evaluation of therapeutic endpoint of CDA formulations. A lot of work ~~still~~ remains to be done.

### 3. CONCLUSION

Cancer evolves as a consequence of wound unhealing. Wound healing process is the most appropriate modality ~~off~~ for cancer therapy. The commanding principle of killing CCs opposing wound healing process, which the cancer establishments assumed for cancer therapy in the past is incorrect. That is why that approach failed to win the war on cancer declared by President Nixon in ~~1971-1976, and~~ 1971-1976 and continued to cause the increase of cancer mortality. Destabilization of abnormal MEs is a critical mechanism of wound healing, which is also the only viable option for the elimination of CSCs. The elimination of CSCs by induction of terminal differentiation is critically linked to the completion of wound healing, and the elimination of CSCs is essential for the success of cancer therapy to save cancer patients. CDA formulations are the best drugs for the induction of terminal differentiation of CSCs.

### REFERENCES

1. Liao MC, Fruehauf JP. It has been half a century since President Nixon declared war on cancer: Destabilization of abnormal methylation enzymes has the blessing of the nature to win the war on cancer. *Adv Complement Alt Med.* 2020; 6(1): 538-539.
2. Liao MC, Craig CL. Wound healing metabolites to heal cancer and unhealed wounds. *Intl Res J Oncol.* 2022; 6(3): 8-20.
3. Google search on cancer statistics-NCI.
4. Liao ~~MC~~ MC. A perfect cancer drug must be able to take out both cancer cells and cancer stem cells, and to restore the functionality of chemo-surveillance. 3<sup>rd</sup> International Conference on Medicinal Chemistry and Drug Design. 2021; page 13.

Formatted: Font: 12 pt, Bold

Formatted: Font: Bold

Formatted: Font: 12 pt, Bold

Formatted: Font: Bold

5. Virchow R. Die Cellular Pathologie in Ihrer Begründung auf Physiologische und Pathologische Gewebelehre. Hirschwald. 1858; 16: 440.
6. Dvorak HF. Tumors: Wounds that do not heal. N Engl J Med. 1986; 315(26): 1650-1659.
7. MacCarthy-Morrow L, Martin IP. The hallmark of cancer **are** also the hallmarks of wound healing. Science Signaling. 2020; 13: 648.
8. Liao MC, Baker LL. Wound healing, evolution of cancer and war on cancer. Intl Res J Oncol. 2021; 4(3): 13-20.
9. Liao MC, Craig CL. On the mechanism of wound healing and the impact of wound on cancer evolution and cancer therapy. Intl Res J Oncol. 2021; 5(3): 25-31.
10. Liao MC, Craig CL. No scar as an indication of perfect wound healing, ugly scar as imperfect wound healing and cancer as failed wound healing. J Cancer Tumor Intl. 2022; 12(1): 29-34.
11. Liao MC, Craig CL. Wound unhealing as a grave issue of cancer. Intl Res J Oncol. 2023; 6(1): 97-103.
12. Liao MC, Szopa M, Burzynski B, Burzynski SR. Chemo-surveillance: A novel concept of the natural defense mechanism against cancer. Drug Exptl Clin Res. 1989; 13(Suppl. 1): 72-82.
13. Liao MC, Craig CL. Chemo-surveillance as a natural mechanism to ensure perfection of wound healing to avoid cancer evolution and to cure cancer. In: New Horizons in Medicine and Medical Research. 2022; Vol. 6: 21-28.
14. Kudo Y, Tateishi K, Yamamoto K, Yamamoto S, Asaoka Y, Ijichi H, et al. Loss of 5-hydroxymethylcytosine is accompanied **with** malignant cellular transformation. Cancer Sci. 2012; 103(4): 670-676.
15. Ficiz GM, Gibben JG. Loss of 5-hydroxymethylcytosine in cancer: Cause or consequence? Genomics. 2014; 104(5): 352-357.
16. Liao MC, Baker LL. Destruction promotes the proliferation of progenitor stem cells and cancer stem cells. Therefore, non-destruction strategy is a better choice of cancer therapy. J Pharmacol Pharmaceu Pharmacovig. 2020; 4:029. DOI: 10.24966/PPP-5459/100029.
17. Monger-Sacca M, Dimrino S, De Maria R. Biological and clinical implication of cancer stem cells in primary brain tumors. Front Oncol. 2013; 3:6.
18. Liu YC, Lee IC, Chen PY. Biomimetic brain tumor niche regulates glioblastoma cells toward a cancer stem cell phenotype. J Neurooncol. 2018; 137(3): 511-522.
19. Zhou S, Schuetz JD, Bunting KD, Colapietro AM, et al. The ABC transporter Bcrp1/ABCG2 is expressed in a wide variety of stem cells and is a molecular determinant of the side-population phenotype. Nat Med. 2001; 7(9): 1028-1034.
20. Zhang M, Atkinson RL, Rosen JM. Selective targeting of radiation resistant tumor initiating cells. Proc Natl Acad Sci USA. 2010; 107(8): 3522-3527.
21. Moitra K, Lou H, Dean M. Multidrug efflux pumps and cancer stem cells: Insight into therapy resistance and therapeutic development. Clin Pharmacol Ther. 2011; 89(4): 491-502.
22. Frame FM, Maitland NU. Cancer stem cells, model of study and implication of therapy resistance mechanisms. Adv Exp Med Biol. 2011; 720(2): 105-118.
23. Liao MC, Craig CL, Baker LL. Elimination of cancer stem cells is essential to save cancer patients. Intl J Res Oncol. 2024; 3(1): 1-9.
24. Liao MC, Craig CL, Baker LL. Modifications of imperfect cancer drugs to become perfect cancer drugs to save cancer patients. Br J Healthcare Med Res. 2024; 11(2): 1-17.
25. Liao MC, Craig DL, Baker LL. CDA formulations as persuasive good cancer drugs to save

- cancer patients. *Intl J Clin Oncol Cancer Res.* 2024; 9(1): 15-24.
26. Liao MC, Baker LL. Cancer ~~patients'~~patients' lives matter. *Adv Complement Alt Med.* 2023; 6(5): 638-640.
  27. Liao MC, Craig CL, Baker LL. A drastic change of cancer leaderships to save cancer patients. In: *New Advance in Medicine and Medical Science.* 2023; Vol. 7: 61-69.
  28. Liao MC, Craig CL, Baker LL. Royal family's cancer diagnoses offer a supreme authority to rectify cancer therapies to save cancer patients. *London J Med Health Res.* 2024; in press.
  29. Liao MC, Lin GW, Hurlbert RB. Partial purification and characterization of tumor and liver S-adenosylmethionine synthetases. *Cancer Res.* 1977; 37(2): 427-435.
  30. Liao MC, Chang CF, Giovanella BC. Demonstration of an altered S-adenosylmethionine synthetase in human malignant tumors xenografted into athymic nude mice. *J Natl Cancer Inst.* 1980; 64(5): 1071-1075.
  31. Liao MC, Zhuang P, Chiou GCY. Identification of the tumor factor of abnormal methylation enzymes as the catalytic subunit of telomerase. *Clin Oncol Cancer Res.* 2010; 7(2): 86-96.
  32. Liao MC, Baker LL. The functionality of chemo-surveillance dictates the success of wound healing as well as cancer therapy. *Novel Res Sci.* 2021; 7(2): 1-3.
  33. Liao MC, Craig CL, Baker LL. Exceptional allosteric regulation of methylation enzymes. In: Saraydin SU (ed), *Novel Research Aspect in Medicine and Medical Research.* 2023; Vol. 4: 38-56.
  34. Liao MC, Craig CL, Baker LL. Restoration of chemo-surveillance as a top priority to save cancer patients. *Intl Res J Oncol.* 2023; 6: 227-237.
  35. Liao MC, Lee SS, Burzynski SR. Hypomethylation of nucleic acids: A key to the induction of terminal differentiation. *Intl J Exptl Clin Chemother.* 1989; 2: 187-199.
  36. Liao MC, Kim JH, Fruehauf JP. Destabilization of abnormal methylation enzymes to combat cancer: The nature's choice to win the war on cancer. Lambert Academic Publishing. 2020; 978-620-2-66889-7.
  37. Liao MC, Chang CF, Becker FF. Alteration of S-adenosylmethionine synthetases during chemical hepatocarcinogenesis and in resulting carcinomas. *Cancer Res.* 1979; 39: 2113-2119.
  38. Kamparath BN, Liao MC, Burzynski B, Burzynski SR. Protective effect of Antineoplaston A10 in hepatocarcinogenesis induced by aflatoxin B<sub>1</sub>. *Intl J Tiss React.* 1990; 12(Suppl.): 43-50.
  39. Liao MC, Craig CL, Baker LL. Wound healing process as the most appropriate modality of cancer therapy. *Eur J Applied Sci.* 2023; 11(1): 463-471.
  40. Liao MC, Craig CL, Baker LL. Wound healing process as the best strategy to save cancer patients. *London J Med Health Res.* 2023; 23(13): 1-11.
  41. Liao MC, Fruehauf JP. Cancer moonshot : Moonshot as a magic code to guide successful solution of tough challenges such as cancer. *Intl J Res Oncol.* 2023; 2(1): 1-5.
  42. Williamson PJ, Krugger AR, Reynolds PJ, Hamlin TJ, Oscier DG. Establishing the incidence of myelodysplastic syndromes. *Br J Haematol.* 1994; 87(4): 743-745.
  43. Boula A, Vouglaris M, Giannouli S, Katrinakis G, Psyllaki M, Pontikoglou C, et al. Effect of CA2 of antitumor necrosis factor- $\alpha$  antibody therapy on hematopoiesis of patients with myelodysplastic syndromes. *Clin Cancer Res.* 2006; 12(10): 3099-3108.
  44. Itkin T, Rafii S. Leukemia cells "gas up" leaky bone marrow blood vessels. *Cancer Cell.* 2017; 32(3): 276-278.

45. Passaro D, Di Tullio A, Abarrategi A, Rousault-Pierre K, Foster K, Ariza-McNaughton L, et al. Increased vascular permeability in the bone marrow microenvironment contribute to disease progression and drug response in acute myeloid leukemia. *Cancer Cell*. 2017; 32(3): 324-341.
46. Counter CM, Gupta J, Harley CB, Lever B. Telomerase activity in normal leukocytes and hematological malignancies. *Blood*. 1995; 89(9): 2315-2320.
47. Fu C, Chen Z. Telomerase activities in myelodysplastic syndromes. *Chin Med J (Engl)*. 2002; 115(10): 1475-1478.
48. Woll PS, Kjallquist U, Chowdhury O, Doolittle H, Wedge DC, Thougjuea S, et al. Myelodysplastic syndromes are propagated by rare and distinct human cancer stem cells in vivo. *Cancer Cell*. 2014; 25(6): 794-808.
49. Ma J. Differentiation therapy of malignant tumor and leukemia. *CSCO Treaties on the Education of Clinical Oncology*. 2007; 480-486.
50. Santi DV, Norment A, Garret CE. Covalent bond formation between a DNA-cytosine methyltransferase of DNA containing 5-azacytosine. *Proc Natl Acad Sci USA*. 1984; 81(22): 6993-6997.
51. Liao MC. Pharmaceutical composition inducing cancer cell differentiation and the use for treatment and prevention of cancer thereof. *US Patent*. 2007; 7232578 B2.
52. Prassana P, Shack S, Wilson VL, Samid D. Phenylacetate in chemoprevention of 5-aza-2'-deoxycytidine-induced carcinogenesis. *Clin Cancer Res*. 1995; 1(18): 865-871.
53. Gaudet F, Hodgson JG, Eden A, Jackson-Grudby L, Gausman J, Gray JW, et al. Induction of tumor in mice by genomic hypomethylation. *Science*. 2003; 300(3618): 489-492.
54. Pali SS, van Emburgh BO, Sankpal UT, Brown KD, Robertson KD. DNA methylation inhibitor 5-aza-2'-deoxycytidine induces reversible DNA damage that is distinctly influenced by DNA-methyltransferase 1 and 3B. *Mol Cell Biol*. 2008; 28(20): 752-771.
55. Kezietepe T, Hedeshima T, Catley L, Raje N, Yasui H, Shiraishi N, et al. 5-Azacytidine, a methyltransferase inhibitor, induces ATR-mediated DNA-double strand break responses, apoptosis, and synergistic cytotoxicity with doxorubicin and bortezomib against multiple myeloma cells. *Mol Cancer Ther*. 2007; 6(6): 1718-1727.
56. Yang Q, Wu F, Wang F, Cai K, Zhang Y, Sun Q, et al. Impact of DNA methyltransferase inhibitor 5-azacytidine on cardiac development of zebrafish in vivo and cardiomyocyte proliferation, apoptosis, and the homeostasis of gene expression in vitro. *J Cell Biochem*. 2019; 120(10): 17549-17471.
57. Liao MC. Differentiation therapy of cancer. *Bull Chin Cancer*. 2002; 11: 104-107.
58. Liao MC. Abnormal methylation enzymes: A selective molecular target for differentiation therapy of cancer. *Chin Pharm J*. 2004; 56: 57-67.
59. Liao MC, Chang CF, Saunders GF, Tsai YH. S-Adenosylhomocysteine hydrolases as the primary target enzymes in androgen regulation of methylation complexes. *Arch Biochem Biophys*. 1981; 208(1): 261-272.
60. Prudova A, Bauman Z, Braun A, Vitvitsky V, Lu SC, Banerjee R. S-Adenosylmethionine stabilizes cystathionine beta-synthase and modulate redox capacity. *Proc Natl Acad Sci USA*. 2006; 103(17): 6489-6494.
61. Chiba P, Wallner C, Kaizer E. S-Adenosylmethionine metabolism in HL-60 cells: Effect of cell cycle and differentiation. *Biochim Biophys Acta*. 1988; 971(1): 38-45.

62. Liao MC, Baker LL. Abnormal methylation enzymes as the bullseye of targeted cancer therapies. *Nov Res Sci.* 2021; 7(4): 1-3.
63. Liao MC, Lee SS, Burzynski SR. Differentiation inducing components of Antineoplaston A5. *Adv Exptl Clin Chemother.* 1988; 6/88: 9-26.
64. Liao MC, Burzynski SR. Separation of active anticancer components of Antineoplaston A2, A3 and A5. *Intl J Tiss React.* 1990; 12(Suppl.): 1-18.
65. Liao MC, Ashraf A, Lee SS, Henry LB, Burzynski SR. Riboflavin as a minor active anticancer component of Antineoplaston A2 and A5. *Intl J Tiss React.* 1990; 12(Suppl.): 19-26.
66. Liao MC, Liao CP, Burzynski SR. Potentiation of retinoid acid-induced terminal differentiation by phenylacetic acid and related chemicals. *Intl J Exptl Clin Chemother.* 1992; 5: 9-17.
67. Liao MC, Huang LJ, Lee JH, Chen SC, Kuo SC. Development of differentiation helper inducers for the differentiation therapy of cancer. *Chin Pharm J.* 1998; 50: 289-303.
68. Liao MC, Liao CP. Methyltransferase inhibitors as excellent differentiation helper inducers for differentiation therapy of cancer. *Bull Chin Cancer.* 2002; 11: 166-168.
69. Liao MC, Fruehauf PA, Zheng ZH, Fruehauf JP. Development of synthetic cell differentiation agent formulations for the prevention and therapy of cancer via targeting of cancer stem cells. *Cancer Stu Ther J.* 2019; 4: 1-15.
70. Liao MC, Kim JH, Fruehauf JP. Potentiation of ATRA activity in HL-60 cells by targeting methylation enzymes. *J Pharmacol Pharmaceu Pharmacovigi.* 2019; 3: 009. DOI: 10.24966/PPP-5649/100009.
71. Liao MC, Kim JH, Fruehauf JP. In pursuance of differentiation inducers to combat cancer via targeting of abnormal methylation enzymes. *J Cancer Tumor Intl.* 2020; 10(2): 39-47.
72. Liao MC, Kim JH, Fruehauf JP. Arachidonic acid and its metabolites as the surveillance differentiation inducers to protect healthy people from becoming cancer patients. *Clin Pharmacol Toxicol Res.* 2021; 4(1): 7-10.
73. Huang M, Ye Y, Chen S, Chai JR, Wang ZY. Use of all trans-retinoic acid in the treatment promyelocytic leukemia. *Blood.* 1988; 72: 567-572.
74. Bourgead MF, Beasoncon F. Induction of 2', 5'-oligoadenylate synthetase by retinoic acid in two transformed human cell lines. *Cancer Res.* 1984; 44: 5355-5360.
75. Mordey JE. Hormone, aging and endocrines in the elderly. In: Felig P, Frohman LA: *Endocrinology and Metabolism.* 4<sup>th</sup> ed. McGraw Hill Inc, Medical Publishing Division. New York. 2001; 1455-1482.
76. Le Coutre P, Mologni L, Cleris L, Marchesi E, Buchdunger E, Giardini R, et al. In vivo eradication of human BCR/ABL-positive leukemia cells with an ABL kinase inhibitor. *J Natl Cancer Inst.* 1999; 91: 163-168.