

Original Research Article

Bio-efficacy of newer insecticides on major insect pests of rice (*Oryza sativa* L.)

ABSTRACT-The present study was carried out at the Farm of Institute of Agricultural Sciences, Banaras Hindu University, Varanasi during 2022-23. During the study on pests of found that stem borer and leaf folder were the main pests infesting paddy. In order to study the bio-efficacy of insecticides; field experiment was laid with six treatments viz., chlorantraniliprole 0.53%G @ 30 g a.i/ha, chlorantraniliprole 0.53%G@39.75g a.i/ha, chlorantraniliprole 0.53%G@49.69 g a.i/ha, chlorantraniliprole 0.4G@ 40 g a.i/ha and fipronil 0.3G @75 g a.i/ha including the untreated control. In case of stem borer and leaf folder chlorantraniliprole 0.53%GR @39.75g a.i/haprovided significantly better control followed by chlorantraniliprole 0.53% GR @ 49.69g a.i/ha, Fipronil 0.30% GR @ 75g a.i/hachlorantraniliprole 0.53%G @ 29.81 g a.i/ha and Chlorantraniliprole 0.40% GR@ 49.69g a.i/ha. It was concluded that the Bio-efficacy of insecticidal treatments against major pests of paddy showed that chlorantraniliprole 0.53%G 39.5g a.i/ha was first best insecticidal treatment against both Lepidopteran insect pests of paddy.

KEYWORD- Chlorantraniliprole, sucking pests, Fipronil, etc

INTRODUCTION

Rice (*Oryza sativa* L.) is a monocotyledonous crop, which belongs to the family Poaceae and genus *Oryza*. Among different insect pests, yellow stem borer (YSB), *Scirpophaga incertulas* (Walker) is the most destructive and widely distributed monophagous pest in the Indian subcontinent. It has assumed the number one pest status. These insect pests infest all plant parts and in all growth of paddy and few also transmit viral diseases (Pathak and KHAN 1994). The rice leaf folder *Cnaphalocrocis medinalis* (Guenee), was previously considered a minor pest but in the last two decades, it has gained major pest status. The use of high quantities of nitrogenous fertilizers and the use of insecticides without proper dosage lead some minor pests to develop major pest status. The larvae fold the leaves and scrape the green tissues of the leaves and cause scorching and leaf drying. The yield loss caused by leaf folder was reported to the extent of 5 to 25% (Kulgagod *et al.*, 2011). These two pests are major pests among lepidopterans infesting rice. The yellow stem borer *S. incertulas* (Walker) has assumed the number one pest status and attacks the crop at all growth stages. Yellow stem borer is distributed all over India and is regarded as the most dominating and destructive pest causing about 25-70% reduction in the yield and it causes 27.34% losses annually (Pasulu *et al.* 2002, Catling *et al.* 1987 and Prasad *et al.* 1988). Despite various chemical control measures, varietal resistance in rice has provided a highly practical approach to controlling insect problems. Attempts to control these pests with indiscriminate use of chemical methods have given rise to many problems viz. resurgence, insecticidal resistance and destruction of natural enemies. The use of chemicals to control this pest is beyond the capacity of marginal farmers. In turn, many insecticidal combinations were used to manage pest problems. Judicious use of insecticides and alternation of chemicals with different modes of action is suggested to reduce insecticide resistance (Mehrotra, 1992). So, the newer insecticides with diversified modes of action against these pests will significantly play a vital role in insect resistance management.

MATERIALS AND METHODS

A field trial was laid out in Randomised Block Design (RBD) with 3 replications and 9 treatments including untreated control during Kharif 2022 to evaluate the bio-efficacy of certain insecticides against insect pests in rice crops and following crop production practices were followed to raise and maintain the crop as per package of practices. The soil in the research plot is sandy clay loam with a neutral pH. The experimental plot is a medium fertile land with assured irrigation and drainage

Comment [T1]: try adapting it to the journal template

Comment [T2]: Enter the driving factors for conducting research

Comment [T3]: enter research objectives

Comment [T4]: enter research objectives

Comment [T5]: enter theoretical sources

Comment [T6]: enter the name of the place and map

Comment [T7]: it is best to complete climate data

facilities. In the present experiment, pests were monitored at regular intervals and when the pest population reached ETL (Economic Threshold Level), insecticides were applied as per the scheduled dates

All the insecticides were in granular form and were mixed with sand and applied by using hand gloves. To avoid intermixing of the treatment, about 20 to 30cm false bund boundaries were prepared all around the plots. All insecticides were directed towards the bottom of the plant. To study the seasonal incidence and efficacy of insecticide, pest population was recorded in different phases i.e. before and after spray. In the case of yellow stem borer, at the vegetative stage number of dead hearts were counted whereas at the reproductive stage number of white ears were counted on 5 randomly selected hill in each treatment at 5th, 10th, 15th day after spray. For counting the population of the rice leaf folder total number of affected leaves to that of the total number of leaves in each hill of individual treatment was recorded. The data for the leaf folder was collected on the 5th, 10th, 15th day after spray. Counts were taken on the number of dead hearts/white ears and the total number of tillers/panicles from 5 randomly selected hills. The damaged leaves and total leaves from 5 randomly selected hills were observed in each plot.

$$\text{Percent Incidence of stem borer} = \frac{\text{Number of dead hearts or white ears}}{\text{total number of tiller or panicle}} \times 100$$

$$\text{Per cent Incidence of leaf folder} = \frac{\text{Number of damaged leaves}}{\text{total number of leaves}} \times 100$$

RESULTS AND DISCUSSION

The field bio-efficacy of certain insecticides against major insect pests of rice during kharif 2022 was assessed. The crop received two sprays during the crop period according to ETL level of pest population. The pretreatment data was recorded one day prior to spraying and the post treatment data at 5TH, 10th, 15th day after each spray. The mean reduction in the population was calculated for analysis.

After the first spray, among all insecticidal treatments chlorantraniliprole 0.53%G @40g a.i/ha considerably reduced (4.56 dead hearts/hill) the yellow stem borer population compared to the control (6.4 dead hearts/hill). The most effective treatment was chlorantraniliprole 0.53%G @40g, which reported the lowest yellow stem borer population, i.e., 5.1, 4.6, 4 dead hearts/hill at 5, 10, 15 DAS respectively which was much better than all other treatments followed by chlorantraniliprole 0.53%G @ 50g a.i/ha i.e 5.3, 5.01, 4.3 dead hearts/hill at 5, 10, 15 DAS respectively was found to be significantly superior to all other treatments followed by fipronil 0.30%G 5.2, 4.7, 4.5 dead hearts/hill at 5, 10, 15 DAS respectively. The least effective pesticide was chlorantraniliprole 0.40%GR which showed mean 5.73 dead heart /hill as compared to 6.53 dead heart /hill in control.

Comment [T8]: It is best to complete the theory that supports the research results

Similar observations were also found in leaf folder affected paddy. Results of impact of insecticidal treatments against *C. medinalis* after first insecticidal application were, least population was observed in chlorantraniliprole 0.53%G @40g a.i/ha which showed 5.77% leaf damage as compared to control 8.10% leaf damage. It was found to be most effective treatment because it showed 5.8, 5.56, 5.97% damage at 5, 10, 15 DAS respectively, followed by chlorantraniliprole 0.53%G @50g a.i/ha which showed 6.84, 6.58, 6.19% leaf damage and fipronil 0.30%GR showed 7.15, 6.94, 6.33% leaf damage at 5, 10, 15 DAS respectively. Least effective among all was chlorantraniliprole 0.53%G @30g a.i/ha which showed mean leaf damage of 7.74% damage as compared to control 8.87% leaf damage.

Comment [T9]: It is best to complete the theory that supports the research results

In the second spray, among all insecticidal treatments chlorantraniliprole 0.53%G @40g a.i/ha considerably reduced (4.93 dead hearts/hill) the yellow stem borer population compared to the control (6.67 dead hearts/hill). Effective treatment was chlorantraniliprole 0.53%G @40g, which reported the lowest yellow stem borer population, i.e., 4.8, 4.3 dead hearts/hill at 5, 10, 15 DAS

respectively which was found the best among all other treatments followed by chlorantraniliprole 0.53%G @ 50g a.i/ha i.e, 5,4.5,4 dead hearts/hill at 5, 10, 15 DAS respectively was found to be significantly superior to all other treatments. followed by fipronil 0.30%GR 5.2,4.7,4.2 dead hearts/hill at 5, 10, 15 DAS respectively. Chlorantraniliprole 0.40%GR was found least effective among all which showed mean 5.2 dead hearts /hill as compared to 6.6 dead hearts /hill in control.

Comment [T10]: It is best to complete the theory that supports the research results

Impact of insecticidal treatments against *C. medinalis* after second insecticidal application were, least population was observed in chlorantraniliprole 0.53%G @40g a.i/ha treated field which showed 3.70% mean leaf damage as compared to control 8.10%(mean) leaf damage. It was found to be most effective treatment because it showed 3.82,3.74,3.56% leaf damage at 5,10,15 DAS respectively, followed by chlorantraniliprole 0.53%G @50g a.i/ha which showed 5.96,5.73,5.39% leaf damage and fipronil 0.30%GR showed 6.2,6.02,5.49% leaf damage at 5, 10, 15 DAS respectively. Least effective among all was chlorantraniliprole 0.53%G @30g a.i/ha which showed mean leaf damage of 7.5% damage as compared to control 8.1% leaf damage.

Comment [T11]: It is best to complete the theory that supports the research results

The efficacy of two sprays was compared using pooled data. The incidence of paddy yellow stem borer was reduced dramatically by all of the insecticides that were tested. It indicated that the average dead hearts before spray were in range of 5.3 to 6.5 dead hearts/plant. chlorantraniliprole 0.53%G @40g a.i/ha was found most significant treatment throughout the experimentation period i.e., 5.1,4.6,4 dead hearts/hill at 5, 10, 15 DAS respectively. The next better treatments were chlorantraniliprole 0.53%G @50g a.i/ha and fipronil 0.30%GR which were equally effective. Followed by chlorantraniliprole 0.53%G @30g a.i/ha Whereas the insecticide chlorantraniliprole 0.40%G @40g a.i/ha was comparatively least effective against paddy yellow stem borer.

Comment [T12]: It is best to complete the theory that supports the research results

Similar pattern of results were also observed in case of paddy leaf folder incidence and percentage of damage significantly reduced in chlorantraniliprole 0.53%G @40g a.i/ha treated plots which showed 5.77% mean leaf damage. Chlorantraniliprole 0.53%G @50g a.i/ha and fipronil 0.30%GR which were equally effective. But chlorantraniliprole 0.40%G @40g a.i/ha was least effective (7.5% leaf damage) in controlling leaf folder population among all insecticides.

Comment [T13]: It is best to complete the theory that supports the research results

Table 1: Effect of insecticidal treatments against *S. incertulus* after 1st spray

Comment [T14]: Do not cut the table

Treatments	Dose (g a.i./ha)	Mean %DH/5hills one day before spray	Mean percent DH per 5 hills at different days after 1 st insecticidal spray			
			5 DAS	10 DAS	15 DAS	Overall Mean
Chlorantraniliprole 0.53% GR	29.81	5.96* (14.13)**	5.4 (13.34)	4.9 (12.79)	4.5 (12.25)	4.933 (12.83)
Chlorantraniliprole 0.53% GR	39.75	5.23 (13.22)	4.8 (12.66)	4 (11.54)	3 (9.97)	3.933 (11.44)
Chlorantraniliprole 0.53% GR	49.69	5.42 (13.46)	5 (12.92)	4.5 (12.25)	4 (11.54)	4.500 (12.25)
Fipronil 0.30% GR	75	5.87 (14.02)	5.2 (13.18)	4.7 (12.52)	4.2 (11.83)	4.700 (12.52)
Chlorantraniliprole 0.40% GR	40	6.23 (14.45)	5.8 (13.94)	5 (12.92)	4.7 (12.52)	5.167 (13.14)
Control	-	5.9 (14.06)	6.5 (14.77)	6.8 (15.12)	6.7 (15.00)	6.667 (14.96)
SE (m) +/-		-	0.148	0.156	0.159	-
C.D. at 5%		-	0.431	0.499	0.507	-

*Mean of 4 replications, ** Figures in paranthesis are angular transformed values, DH = Dead heart

Treatments	Dose (g a.i./ha)	Mean %WE/5hills one day before spray	Mean percent DH per 5 hills at different days after 1 st insecticidal spray			
			5 DAS	10 DAS	15 DAS	Overall Mean
Chlorantraniliprole 0.53% GR	29.81	6.3* (14.54)**	5.8 (13.94)	5.6 (13.69)	5.1 (13.05)	5.5 (13.56)
Chlorantraniliprole 0.53% GR	39.75	5.3 (13.31)	5.1 (13.05)	4.6 (12.38)	4 (11.54)	4.56 (12.33)
Chlorantraniliprole 0.53% GR	49.69	5.4 (13.44)	5.3 (13.31)	5.01 (12.93)	4.3 (11.97)	4.87 (12.75)
Fipronil 0.30% GR	75	5.5 (13.56)	5.2 (13.18)	4.7 (12.52)	4.5 (12.25)	4.80 (12.66)
Chlorantraniliprole 0.40% GR	40	6.5 (14.77)	6.3 (14.54)	5.7 (13.81)	5.2 (13.18)	5.73 (13.85)
Control	-	6.8 (15.12)	6.4 (14.65)	6.5 (14.77)	6.7 (15)	6.53 (14.81)
SE (m) +/-		-	0.173	0.175	0.157	-

Table 2: Effect of insecticidal treatments against *S. incertulus* after 2st spray

Comment [T15]: title adjust the others

Table 3: Effect of insecticidal treatments against *C. medinalis* after 1st spray

Comment [T16]: Do not cut the table

Treatments	Dose (g a.i./ha)	Mean %DH/5hills	Mean percent DH per 5 hills at different days after 1 st insecticidal spray
------------	------------------	-----------------	--

	a.i/ha)	one day before spray	5 DAS	10 DAS	15 DAS	Overall Mean
Chlorantraniliprole 0.53% GR	29.81	7.8 (16.22)	7.6 (16)	6.9 (15.23)	6.8 (15.12)	7.10 (15.45)
Chlorantraniliprole 0.53% GR	39.75	7.9 (16.32)	5.8 (13.94)	5.56 (13.64)	5.97 (14.14)	5.77 (13.90)
Chlorantraniliprole 0.53% GR	49.69	8.1 (16.54)	6.84 (15.16)	6.58 (14.84)	6.19 (14.41)	6.53 (14.81)
Fipronil 0.30% GR	75	8.6 (17.05)	7.15 (15.51)	6.94 (15.27)	6.33 (14.57)	6.80 (15.12)
Chlorantraniliprole 0.40% GR	40	9.2 (17.66)	8.07 (16.50)	7.8 (16.22)	7.37 (15.75)	7.74 (16.15)
Control	-	9.6 (18.05)	9.35 (17.80)	9.08 (17.54)	8.2 (16.64)	8.87 (17.33)
SE (m) +/-	-	-	0.266	0.273	0.189	-
C.D. at 5%	-	-	0.85	0.871	0.603	-

Table 4: Effect of insecticidal treatments against *C .medinalis* after 2nd spray

Treatments	Dose (g a.i/ha)	Mean %leaf damage/5hills one day before spray	Mean percent DH per 5 hills at different days after 1 st insecticidal spray			
			5 DAS	10 DAS	15 DAS	Overall Mean
Chlorantraniliprole 0.53% GR	29.81	6.9 (15.23)	6.68 (14.98)	6.49 (14.76)	6.10 (14.30)	6.42
Chlorantraniliprole 0.53% GR	39.75	6.1 (14.30)	3.82 (11.27)	3.74 (11.15)	3.56 (10.88)	3.70
Chlorantraniliprole 0.53% GR	49.69	6.5 (14.77)	5.96 (14.13)	5.73 (13.85)	5.39 (13.42)	5.6
Fipronil 0.30% GR	75	6.9 (15.23)	6.20 (14.42)	6.02 (14.20)	5.49 (13.55)	5.9
Chlorantraniliprole 0.40% GR	40	8.1 (16.54)	7.87 (16.29)	7.64 (16.05)	7.19 (15.55)	7.5
Control	-	8.6 (17.05)	8.58 (17.03)	8.34 (16.79)	7.60 (16.00)	8.1
SE (m) +/-	-	-	0.208	0.202	0.187	-
C.D. at 5%	-	-	0.664	0.644	0.597	-




FIG1: Bioefficacy of overall mean field bioefficacy of insecticidal treatments on *S. incertulas*

Comment [T17]: adjust to the journal template




FIG 2: BIOEFFICACY OF DIFFERENT INSECTICIDAL TREATMENTS ON *C. medinalis*

Comment [T18]: adjust to the journal template

CONCLUSION

It was concluded that the Bio-efficacy of insecticidal treatments against major pests of paddy showed that chlorantraniliprole 0.53%G 39.5g a.i/ha was first best insecticidal treatment against both Lepidopteran insect pests of paddy. It also suit well for leaf folder. Fipronil can be incorporated in

Comment [T19]: try checking in the adjusted table

integrated pest management practices as it showed persistent toxic effects and gave an effective control for all the major pests of paddy and also improved the yield. Alternatively, chlorantraniliprole 0.53%G 30g a.i also be used both for effective management of chewing and sucking pests like stem borer and leaf folder.

Comment [T20]: try checking in the adjusted table

REFERENCES

- Aulakh, S. S., Randhawa, H. S., & Singh, M. (2016). Bioefficacy of insecticides for management of stem borer and leaf folder on paddy in Punjab, India. *Agricultural Science Digest-A Research Journal*, 36(3), 224-227.
- Bhardwaj, U., Kumar, R., Sarao, P. S., Mandal, K., & Singh, B. (2019). Bioefficacy of chlorantraniliprole against yellow stem borer and leaf folder and persistence in basmati rice. *Pesticide Research Journal*, 31(2), 233-241.
- Catling, H.D., Z. Islam and R. Patrasudhi.(2003) Assessing yield losses in deep water rice due to yellow stem borer, *S.incertulas* in Bangladesh and Thailand. *Crop Prot. J.*,6, 20-27.
- Chormule, A. J., Kharbade, S. B., Patil, S. C., & Tamboli, N. D. (2014). Bioefficacy of new insecticide molecules against rice yellow stem borer, *Scirpophaga incertulas* (Walker). *The ecoscan*, 6, 63-67.
- Dash, A.N. and Mukherjee, S.K. (2003). Insecticidal control of major insect pest of rice. *Pest Management and Economic Zoology*. 11(2): 147-151.
- Dhaka, S.S., Prajapati, C.R., Singh, D.V. and Singh R. (2011). Field evaluation of insecticides and bio-pesticides against Rice leaf folder, *Cnaphalocrocis medinalis* (Guenee). *Ann. Pl. Protec. Sci.* 19(2): 324-326.
- Dhivahar, P. and Dhandapani, N. (2003). Bio-efficacy of the new molecule thiacloprid (Calypso 240 SC) against rice yellow stem borer, *Scirpophaga incertulas* (Walker). *Indian Journal of Plant Protection*.31(1): 144-145.
- Heinrichs, E.A. (1994). Impact of insecticides on the resistance and resurgence of Rice plant hoppers. *Plant hoppers: Their ecology and management*. p. 571-598
- Hurali, S., Prasad, P. R., Shiralli, H., Gowdar, S. B., Mahantashivayogayya, K., & Masthanareddy, B. G. (2023). Evaluation of efficacy of new combi insecticides against lepidopteran pests of rice and its impact on natural enemies. *Journal of Experimental Zoology India*, 26(1).
- Hussain, G., Sultana, T. and Shah, A. (2008). Efficacy and economics of different insecticides against stem borer *Scirpophaga insertulus* in rice crop. *Pakistan j. zool.* Vol. 45(4): 929-933
- Jaglan, M. S., Chaudhary, O. P., Ahlawat, S., & Yadav, J. (2023). Bio-efficacy of chlorantraniliprole 0.4% GR against *Cnaphalocrocis medinalis* and *Scirpophaga incertulas* in rice. *Indian Journal of Agricultural Sciences*, 93(2), 157-162.

Comment [T21]: The reference does not match the theoretical source. Try checking and looking at the journal template

- Jena, M. (2004). Efficacy of new insecticides as seedling root dip treatment against yellow stem borer in Rabi rice. *Indian Journal of Plant Protection*. 32(2): 37-39
- Karthikeyan, K. and Purushothaman, S.M.(2000). Efficacy of carbosulfan against rice yellow stem borer, *Scirpophaga incertulus* (Walker) in rabbi rice. *Indian Journal of Plant Protection*, 28(2): 212-214.
- Khan, A.A and Misra, D.S. (2003). Abundance of spider in relation to biotic and abiotic factors in upland rice ecosystem of Eastern Uttar Pradesh. *Plant Protection Bulletin*, Faridabad. 55(3/4): 23-29.
- Kooner, R., Suri, K. S., & Makkar, G. S. (2022). Efficacy of Takumi 20% WG against Rice Stem Borers and Leaf Folder. *Indian Journal of Entomology*, 1-4.
- Kulgagod, S.D. Hegade, M., Nayak, G.V. Vastrad, A.S., Hugar, P. S. and Basavanagud, K.(2011). Evaluation of insecticide and biorational against yellow stem borer and leaf folder in rice crop. *Karnataka Journal of Agricultural science*. 24(2):244-246.
- Kumar, R., Singh, B., Mandal, K., Kar, A., & Sarao, P. S. (2013). Bioefficacy and fate of fipronil and its metabolites in basmati rice under sub-tropical climatic conditions. *Crop Protection*, 45(1) :41-48.
- Mehrotra, K.N.(1992). Pesticide resistance in insects; Indian Scenario. *In Pest Management and Pesticides*: 17-27.
- Mishra, D.N, Kumar, K. Singh, L.R. (2007). Comparative field efficacy of certain granular insecticides against leaf roller and stem borer in rice crop under Mid-Western Plain Zone of UP. *Environment and Ecology*. 25(4): 938-940.
- Misra, A.K., Singh, S.P.N., Parwez, A. (2005). Incidence of yellow stem borer (*S. incertulas* Wlk.) in different cultivars of boro rice (*Oryza sativa* L.) at different crop age. *Oryza*. 42(4): 329-332.
- Mondal, S. C. A comparative study on the physicochemical properties of selected varieties of rice grains in Bangladesh.
- Murty, M.M.K., Rao, D.V.S. and Azam, K.M. (1985). Efficacy of certain insecticides as foliar formulations in the control of rice stem borer, *Scirpophaga incertulas* and leaf roller, *Cnaphalocrosis medinalis*. *Indian Journal of Entomology*. 47(4): p. 461.
- Pasulu, I.C., N.V. Krishnaiah, G. Kotti and N.R.G. Varma: *IPM in rice mitr*. pp. 45-55 (2002).
- Pathak, M.D. and Khan, Z.R. (1994). Insect pests of rice, Manila, Phillipines. *International Rice Research Institute*.
- Paul, B., Samanta, S., Sen, K., Manger, A., & Samanta, A. (2018). Evaluation of bioefficacy and phytotoxicity of fipronil 5% SC and acetamiprid 20% SP on rice insect pest complex.
- PENROSE, L. Fingerprint Science. *Nature* 210, 1089–1090.
- Prasad, S.S., P.K. Gupta and R.B. Singh (1988) Yield losses in floating rice caused by stem borers. *Int. Rice Res. News*, 13, 38.

- Purohit, M.S., Shah, A.H. and Raman, S. (1988). Chemical control of rice stem borer *Scirpophaga incertulas* Wlk. in Gujarat. Gujarat Agricultural University Research Journal. 13(2): 23-27.
- Rajinder, K., Balwinder, S., Kousik, M., Abhijit, K., & Sarao, P. S. (2013). Bioefficacy and fate of fipronil and its metabolites in basmati rice under sub-tropical climatic conditions. *Crop Protection*, 45, 41-48.
- Samanta, S., Barman, M., & Samanta, N. R. A. (2020). Bio-efficacy trials of carbofuran 3% CG against insect pests of rice. *J. Entomol. Zool. Stud*, 8: 1529-1533.
- . S., & Kaur, H. (2014). Efficacy of Ferra 0.4% GR (chlorantraniliprole) against stem borers and leaf folder insect-pests of basmati rice. *Journal of Environmental Biology*, 35(5), 815.
- Sharma, P. K. (2008). Bio-efficacy of insecticides against paddy insect pests under mid-hill conditions of Himachal Pradesh. *Agricultural Science Digest*, 28(1), 61-62.
- Suri, K. S. (2011). Bioefficacy of some insecticides against rice stem borers and leaf folder infesting rice in Punjab. *Pesticide Research Journal*, 23(2), 190-193.
- Yeligar, R., Kalmath, B. S., Radha, J., Madli, M., & Jyothi, R. (2020). Bio-efficacy of Spinetoram 6% w/v (5.66% w/w)+ Methoxyfenozide 30% w/v (28.3% w/w) SC against lepidopteran insect pests in Rice ecosystem.