

INVENTORY OF SEaweEDS IN SELECTED MUNICIPALITIES IN THE BALICUATRO AREA OF THE PROVINCE OF NORTHERN SAMAR, PHILIPPINES

ABSTRACT

Seaweeds or marine macroalgae is one of the marine resources that are understudied in the province of Northern Samar, the province have twenty-four (24) municipalities and twenty (20) of them have coastal zone which is ideal for marine researches as well as for aquaculture. This study is focused on the present seaweeds in the three (3) selected municipalities in Balicuatro area in the province of Northern Samar. A total of ninety-four (94) taxa were collected/ found in the three (3) sampling sites. Lavezares got the greatest number of seaweeds with sixty (60) taxa, followed by Biri with forty-four (44), and the least was in Allen with thirty-four (34) species. Chlorophyta have thirty-nine (39) species which constitute for 39.36% of the total identified seaweeds, followed by thirty-four (34) red seaweeds (36.17%), and, lastly, Brown macroalgae with only twenty-three (23) species which accounts for 24.47%. The most common marine macroalgae in all sampling sites were: *Hormophysa cuneiformis* which was found in ten (10) sampling sites, followed by *Halimeda macroloba*, *Ulva reticulata*, and *Sargassum oligocystum* which were found in ten (10) sampling sites. In terms of environmental parameters, temperature and salinity of all sampling sites were tested to be within the normal range during the collection period based on established threshold limit. However, in terms of pH, it was detected that two (2) barangays/ sampling sites in Lavezares, namely: Balicuatro and San Juan which only had 7.2 and 7.4 which is below the threshold limit range of seawater pH which is 7.5-8.5. In terms of substrate, rocky substrate is common to all sampling sites which indicates that hard substratum is preferable for the growth, adaptation, and survival of seaweeds.

Keyword: Chlorophyta, Phaeophyta, Rhodophyta, taxa, environmental parameters.

INTRODUCTION

The province of Northern Samar is divided into three areas, namely: Balicuatro, Central, and Pacific area (Galenzoga, 2016). The coastal marine water of the province is enriched by the nutrients coming from the land and it is also characterized by just the right amount of rain and a warm tropical climate. The intertidal/ shallow water are well **were** crafted with developed reef that supports the diverse marine flora and fauna, among these intricate floras in the coastal environment are the mangal community, seagrasses, and seaweeds (Trono, 1997).

In general, the Philippines have 1,065 species of seaweeds (Lastimoso and Santianez, 2020). Those numbers are broadly grouped based on their nutrient and chemical composition into **Chlorophyta (Green algae), Phaeophyta (Brown algae), and Rhodophyta (Red algae)**. The study of seaweeds in the country began in the 1960's when Filipino phycologists participated in the world's discovery of algae and algal researches, specifically in the field of seaweed taxonomy (Baldia *et al.*, 2017). It is during 1960's when Menez studied Philippine marine macroalgae in Hundred Islands (Menez, 1961; Baldia *et al.*, 2017), it was followed by Cordero which brought significant contributions in red algae (Cordero, 1977; Baldia *et al.*, 2017), and Trono on the marine benthic macroalgae (Trono, 1972; Baldia *et al.*, 2017).

At present, there were few published papers and information on the available marine macroalgae in Northern Samar. Out of 24 municipalities in the province, twenty (20) municipalities have shorelines and coastal zones making the province a good place to study seaweeds and since two (2) of the island municipalities are still unexplored when it comes to its seaweed resource. Most recent studies on seaweeds conducted in the province were carried by Baldia *et al* (2017) by which they focused on the diversity assessment of marine macroalgae in Biri and Dalupirit Island and Galenzoga (2019) on the present seaweed resource in Biri Island.

Even though seaweeds are primary plants that do not bear flowers, roots, stems, and leaves they are considered as an important food source commodity and income in the Philippines (Dumilag, 2019). In another study, the carrageenan extract from seaweeds is also used a precursor in the development of antibacterial biofilms that could be used as an alternative to commercially available biofilms (Dagalea, Lim, 2018; Dagalea, Lim, 2021). In 2014, Northern Samar ranked 3rd when it comes to seaweed production in Region VIII which accounts for about 353.86 metric tons from 20.3 hectares of seaweed area/ farm in the province (Pacturan, 2016). Recently, there are current trends on seaweed as a valuable source of bioactive compounds, polysaccharides, antioxidants, minerals, and essential nutrients such as fatty acids, amino acids, and vitamins that could be used as a functional ingredient (Kumar *et al.*, 2021). These compounds gave seaweeds a higher value in the individual diet as a food component and as pharmaceutical supplements (Namvar *et al.*, 2013; Choudhary *et al.*, 2021).

Therefore, it is the purpose of this research to contribute to the initial information on the present species composition of seaweeds in some areas in Northern Samar. Furthermore, the result of this study will add to the existing information on seaweeds in the Philippines.

MATERIALS AND METHODS

Study Site

This study was conducted in three (3) municipalities in the Balicuatroarea of Northern Samar, namely: Allen, Lavezares, and Biri. There were a total of eleven (11) coastal barangays that served as sites for the collection of seaweeds.




Figure 1. The map of the sampling sites. A). Map of Northern Samar highlighting the three (3) municipalities in Balicuatro District; B). Map of Lavezares pinpointing the collection sites/ barangays, namely: Balicuatro, Barobaybay, Bani, San Agustin, and San Juan; C). Map of Biri pinpointing the collection sites/ barangays, namely: Pio del Pilar, Progress, Mc Arthur, San Pedro, and San Antonio; and D). Map of Allen pinpointing the collection site/ barangay of Guinarawayan. Maps were taken from the municipal offices of Allen, Biri, and Lavezares. Northern Samar

Northern Samar is situated in the Eastern Visayas region occupying the eastern section of the Visayas. The province is bordered, clockwise from the North, by the Philippine Sea, Eastern Samar, Samar Sea, and San Bernardino Strait.

Data Gathering Procedure

In every sampling site, 50-100 meter transect line were laid down perpendicular to the rock shore. Random sampling technique were employed during the sampling wherein all of the seaweeds found along the 50-100 meter intertidal zone were collected and identified. All of the seaweeds collected were placed in a ziplock and brought to the laboratory for proper identification of samples.

Identification and Authentication of Samples

Some of the seaweeds were identified *in situ* based on morphological criteria. Phenotypic characteristics of the samples were determined and examined using available literature (Dumilag, 2019, Dumilag and Javier, 2022, Sherwood and Guiry, 2023) to identify samples down to species level. The books of Calumpong (1997) and Trono (1997) were also used for identification and authentication.

Preservation of samples

A portion or whole part of the seaweed samples were placed in a plant press which served as voucher specimens. All samples prior to pressing were washed in a running water to rinsed-off the epiphytes, dirt, and other organisms attached to the seaweed sample. Samples with bulky thalli were soaked in rock salt for a few days to dehydrate and facilitate proper drying. After washing, the seaweed samples were then placed in a neat newspaper and a piece of fabric/ cheesecloth was then placed above the samples to prevent it from sticking to the newspaper.

Determination of Environmental Parameters

Prevailing environmental parameters that characterize the study area were determined during the sampling to the sites, these includes pH, temperature, salinity, and substrate. A water sample was collected at every site and placed in an ice box with ice for its transportation to the College of Science. Bante 900-UK Multiparameter Water Quality Meter was used to determine the pH, temperature, and salinity, while ocular inspection to the sites were conducted regarding the substrate type of the intertidal zone.

RESULTS

Species Composition of Seaweeds

There were a total of ninety-four (94) species of seaweeds identified in three (3) municipalities in the Balicuatro area of Northern Samar. As shown in Table 1, all of the collected samples were represented by the three (3) classes of seaweeds/ macroalgae, namely: Chlorophyta (green algae/ seaweeds), Phaeophyta (brown algae/ seaweeds), and Rhodophyta (Red algae/ seaweeds). Green algae constitute for about thirty-nine (39) (39.36%) out of 94 collected seaweed species, followed by Red algae with thirty-four (34) (36.17%) species, and Brown algae with twenty-three (23) species which accounts for 24.47%.

Table 1. List of Seaweeds in the three municipalities in Balicuatro Area, Northern Samar.

Class	Species	
<p>Chlorophyta (37) 39.36%</p>	<p>Family Anadyomenaceae <i>Anadyomene plicata</i> C. Agardh</p>	
	<p>Family Caulerpaceae <i>Caulerpa brachypus</i> Harvey <i>Caulerpa cupressoides</i> (Vahl) C. Agardh <i>Caulerpa lentillifera</i> J. Agardh <i>Caulerpa racemosa</i> var. <i>macrophysa</i> (Sonder ex Kützing) W.R. Taylor <i>Caulerpa racemosa</i> (Forsskål) <i>Caulerpa serrulata</i> (Forsskål) J. Agardh <i>Caulerpa taxifolia</i> (Vahl) C. Agardh</p>	
	<p>Family Cladophoraceae <i>Chaetomorpha crassa</i> (C. Agardh) Kützing</p>	
	<p>Family Codiaceae <i>Codium geppii</i> O.C. Schmidt <i>Codium intricatum</i> Okamura <i>Codium vermilara</i> (Olivieri) Delle Chiaje</p>	
	<p>Family Dasycladaceae <i>Bornetella oligospora</i> Solms-Laubach <i>Bornetella sphaerica</i> (Zanardini) Solms-Laubach <i>Cymopolia vanbosseae</i> Solms- Laubach <i>Halicoryne wrightii</i> Harvey <i>Neomeris annulata</i> Dickie</p>	
	<p>Family Dichotomosiphonaceae <i>Avrainvillea erecta</i> (Berkeley) A. Gepp & E. Gepp</p>	
	<p>Family Halimedaceae <i>Halimeda macroloba</i> Decaisne <i>Halimeda macrophysa</i> Askenasy <i>Halimeda opuntia</i> (Linnaeus) Lamouroux <i>Halimeda tuna</i> (Ellis and Solander) Lamouroux <i>Halimeda velasquezii</i> W.R. Taylor</p>	
	<p>Family Polyphysaceae <i>Acetabularia dentata</i> Solms-Laubach <i>Acetabularia major</i> Martens</p>	
	<p>Family Siphonocladaceae <i>Boergesenia forbesii</i> (Harvey) J. Feldmann <i>Boodlea composita</i> (Harvey) Brand</p>	
	<p>Family Udoteaceae <i>Chlorodesmis fastigiata</i> (C. Agardh) Ducker</p>	
	<p>Family Ulvaceae <i>Ulva australis</i> Areschoug <i>Ulva clathrata</i> (Roth) <i>Ulva flexuosa</i> Wulfen <i>Ulva intestinalis</i> Linnaeus <i>Ulva lactuca</i> Linnaeus <i>Ulva reticulata</i> Forsskål</p>	
	<p>Family Valoniaceae <i>Dictyosphaeria cavernosa</i> (Forsskål) Borgesen <i>Dictyosphaeria versluysii</i> Weber-van Bosse <i>Valonia fastigata</i> Harvey ex J. Agardh</p>	
		<p>Family Dictyotaceae <i>Canistrocarpus cervicornis</i> (Kützing) De Paula and De Clerk <i>Dictyota dichotoma</i> (Hudson) Lamouroux <i>Dictyota mertensii</i> (Martius) Kützing <i>Padina australis</i> <i>Padina gymnospora</i> (Kützing) Sonder <i>Padina minor</i> Yamada <i>Padina tetrastromatica</i> Hauck</p>
		<p>Family Sargassaceae <i>Hormophysa cuneiformis</i> (J.F. Gmelin) P.C. Sliva</p>

Phaeophyta (23) 24.47%	<i>Sargassum aquifolium</i> (Turner) C. Agardh <i>Sargassum cinctum</i> (J. Agardh) <i>Sargassum cristaefolium</i> C.A. Agardh <i>Sargassum feldmanii</i> Pham-Hoang <i>Sargassum gracillimum</i> Reinbold <i>Sargassum hemiphyllum</i> (Turner) C. Agardh <i>Sargassum oligocystum</i> Montagne <i>Sargassum polyscystum</i> C.A. Agardh <i>Sargassum siliquosum</i> J. Agardh <i>Turbinaria conoides</i> (J. Agardh) Kützing <i>Turbinaria decurrens</i> Bory de Saint-Vincent <i>Turbinaria luzonensis</i> Taylor <i>Turbinaria ornata</i> (Turner) J. Agardh
	Family Scytosiphonaceae <i>Colpomenia sinoua</i> (Mertens ex Roth) Derbs and Solier <i>Hydroclathrus clathratus</i> (C. Agardh) Howe
Rhodophyta (34) 36.17%	Family Bangiaceae <i>Trichogloea requienii</i> (Montagne) Kützing
	Family Champiaceae <i>Coelothrix irregularis</i> (Harvey) Borgessen
	Family Corallinaceae <i>Amphiroa foliacea</i> <i>Amphiroa fragilissima</i> (Linnaeus) Lamouroux <i>Mastophora rosea</i> (C. Agardh) Setchell <i>Jania adhaerens</i> J.V. Lamouroux
	Family Cryptonemiaceae <i>Halymenia durvillei</i> Bory de Saint-Vincent
	Family Cystocloniaceae <i>Hypnea cervicornis</i> J. Agardh
	Family Delesseriaceae <i>Caloglossa cf. beccarii</i> (Zanardini) De Toni
	Family Galaxauraceae <i>Actinotrichia fragilis</i> (Forsskål) Borgesen <i>Dichotomaria 6piculate</i> Kjellman <i>Scinaia hormoides</i> Setchell <i>Tricleocarpa fragilis</i> (Linnaeus) Huisman & R.A.
	Family Gelidiaceae <i>Gelidiella acerosa</i> (Forsskål) Feldmann and Hamel
	Family Gracilariaceae <i>Gelidiopsis intricata</i> (C. Agardh) <i>Gracilaria arcuata</i> Zanardini <i>Gracilaria salicornia</i> (C. Agardh) Dawson <i>Gracilaria textorii</i> (Suringar) De Toni <i>Hydropuntia edulis</i> (S.G. Gmelin) Gurgel & Fredericq
	Family Liagoraceae <i>Ganonema farinosum</i> (J.V. Lamouroux) K.C. Fan & Y.C. Wang <i>Liagora ceranoides</i> J.V. Lamouroux
	Family Kallymeniaceae <i>Euthora cristata</i> (C. Agardh) J. Agardh
	Family Peyssonneliaceae <i>Grateloupia filicina</i> (Lamouroux) C. Agardh
	Family Rhizophyllidaceae <i>Portieria hornemannii</i> (Lyngbye) P.C. Silva
	Family Rhodomelaceae <i>Acanthopora muscoides</i> (Linnaeus) Bory-de Saint-Vincent <i>Acanthopora spicifera</i> (Vahl) Borgesen <i>Laurencia cartilaginea</i> Yamada <i>Laurencia nidifica</i> J. Agardh <i>Laurencia papillosa</i> (C. Agardh) Greville <i>Laurencia tronoi</i> Ganzon-Fortes <i>Ohelopapa flexilis</i> (Setchell) F. Rousseau, Martin-Lescanne, Payri & Le Gall
	Family Solieraceae <i>Betaphycus philippinensis</i> Doty <i>Eucheuma denticulatum</i> (N.L. Burman) Collins and Hervey <i>Kappaphycus striatus</i> (F. Schmitz) L.M. Liao

<i>Ganonema farinosum</i> (J.V. Lamouroux) K.C. Fan & Y.C. Wang												/
<i>Liagora ceranoides</i> J.V. Lamouroux											/	
Family Kallymeniaceae												
<i>Euthora cristata</i> (C. Agardh) J. Agardh				/								
Family Peyssonneliaceae												
<i>Grateloupia filicina</i> (Lamouroux) C. Agardh											/	
Family Rhizophyllidaceae										/		
<i>Portiera hornemannii</i> (Lyngbye) P.C. Silva	/	/										/
Family Rhodomelaceae												
<i>Acanthopora muscoides</i> (Linnaeus) Bory-de Saint-Vincent		/	/		/							/
<i>Acanthopora spicifera</i> (Vahl) Borgesen										/		
<i>Laurencia cartilaginea</i> Yamada								/				
<i>Laurencia nidifica</i> J. Agardh												/
<i>Laurencia papillosa</i> (C. Agardh) Greville		/		/								/
<i>Laurencia tranoi</i> Ganzon-Fortes	/	/		/								
<i>Ohelopapa flexilis</i> (Setchell) F. Rousseau, Martin-Lescanne, Payri & Le Gall							/	/			/	
Family Solieraceae												
<i>Betaphycus philippinensis</i> Doty								/				
<i>Eucheuma denticulatum</i> (N.L.Burman) Collins and Hervey			/	/								
<i>Kappaphycus striatus</i> (F. Schmitz) L.M. Liao								/		/	/	

Legend:

- | | |
|-------------------------|---------------------------|
| 1- Progress | 7- Barobaybay, Lavezares |
| 2- Pio del Pilar | 8- San Agustin, Lavezares |
| 3- McArthur | 9- Bani, Lavezares |
| 4- San Pedro | 10- San Juan, Lavezares |
| 5- San Antonio | 11- Guinarawayan, Allen |
| 6- Balicutro, Lavezares | |

In Lavezares, the barangay with the most number of seaweeds was in Barangay Barobaybay with thirty-one (31) species, followed by Barangay San Juan with twenty-eight (28) seaweed species, Barangay San Agustin have twenty-four (24) seaweeds and Barangay Bani with twenty-three (23), and as mentioned Barangay Balicutro were the least with only twelve (12) species of seaweeds collected. Biri, as mentioned Barangay Pio del Pilar have the most number of seaweeds with thirty six (36) collected seaweeds, followed by Barangay Progress with thirty (30) number of seaweeds, while the number seaweeds collected in Barangays San Antonio, San Pedro, and McArthur were 26, 22, and 17, respectively. In addition, the only barangay sampled in Allen was Guinarawayan with thirty-four (34) number of seaweeds collected and identified.

Most Abundant Seaweed Species

The most abundant seaweed species that was collected and observed in ten (10) sampling sites was *Hormophysa cuneiformis* which was found abundantly growing in current-exposed area in the lower intertidal, it was followed by *Halimeda macroloba* which was found growing abundantly in almost all the sampling sites, though *Ulva lactuca* was found in nine (9) sampling sites they are found growing only in patch and solitary which is in comparison with *Ulva reticulata* which were only found in eight (8) sampling sites but they grow lavishly in all of the sampling sites to where it was collected. *Padina minor* were also collected and observed in almost all the sampling sites, it is present in nine (9) sampling sites and was found widely distributed with other seaweeds in all sampling sites to where it is observed. *Sargassum oligocystum* and *S.*

polycystum were found in nine (9) and eight (8) sampling sites, respectively, they were found abundantly in lower




Figure 2. Most abundant seaweed species: (A). *Halimeda macroloba*; (B). *H. opuntia*; (C). *Ulva lactuca*; (D). *U. reticulata*; (E). *Bornetella sphaerica*; (F). *Padina minor*; (G). *Sargassum oligocystum*; (H). *S. polycystum*; (I). *Hormophysa cuneiformis*; (J) *Canistrocarpus cervicornis*; (K). *Hydropuntia edulis*; (L). *Tricleocarpa fragilis*.

intertidal zone. The most abundant red macroalgae were *Tricleocarpa fragilis* which was found in seven (7) sampling sites and *Hydropuntia edulis* which was found in six (6) sampling sites, *H.edulis* were found widely distributed entangled with other red macroalgae. *Bornetella sphaerica* and *Canistrocarpus cervicornis* were also found in six (6) sampling sites. *B. sphaerica* were found growing in cluster of dead coral rubles.

Environmental Parameters

Selected environmental parameters were measured (Table 3) using Bante 900-UK Multiparameter Water Quality Meter. These environmental parameters are all critical for the adaptation, growth, and survival of most aquatic plants and animals. In terms of pH, the highest was measured in barangay San Pedro and San Antonio, Biri which had 7.9. which within the normal range of 7.5-8.5 (Water watch, 2022). Meanwhile, two (2) barangays in Lavezares tested to be below the threshold limit for pH, it was measured in barangay Balicuatro and San Juan which had 7.2 and 7.4, respectively. In terms of salinity, all of the salinity levels of the sampling sites conform to the threshold range for the seawater which is 30-35 ppt (Water watch, 2022).

Table 3. Environmental Parameters in different sampling sites.

Environmental Parameters	Sampling Sites											Permissible Limit
	Biri					Lavezares					Allen	
	1	2	3	4	5	6	7	8	9	10	11	
pH	7.8	7.8	7.6	7.9	7.9	7.2	7.6	7.6	7.8	7.4	7.7	7.5- 8.5 (Water watch, 2022)
Salinity	34 ppt	35 ppt	35 ppt	33 ppt	34 ppt	35 ppt	35 ppt	35 ppt	34 ppt	35 ppt	34 ppt	30- 35 ppt (Water watch, 2022)
Temperature	26°C	27°C	27.5°C	27.5°C	28°C	28.0°C	26.5°C	26.5°C	26.5°C	26.5°C	26.5°C	26-28 °C (Johnson & Xie, 2010)
Substrate	Rocky	Rocky	Rocky-corally	Rocky	Rocky-corally	Rocky	Rocky-corally	Muddy & Sandy	Sandy	Rocky & Sandy	Rocky-corally	

Legend:

- | | |
|--------------------------|---------------------------|
| 1- Progress | 7- Barobaybay, Lavezares |
| 2- Pio del Pilar | 8- San Agustin, Lavezares |
| 3- McArthur | 9- Bani, Lavezares |
| 4- San Pedro | 10- San Juan, Lavezares |
| 5- San Antonio | 11- Guinarawayan, Allen |
| 6- Balicuatro, Lavezares | |

Furthermore, in terms of water temperature all of the sampling sites also conform to the threshold limit of 26-28°C for seawater. On the other hand, the most common substrate among all the sampling sites was rocky-corally in which it was observed that some macroalgae preferred to have a hard substratum and rocks can provided hard attachment of seaweeds.

DISCUSSION

Recently, Lastimoso and Santianez (2020) reported that there were one-thousand sixty five (1,065) taxa of seaweeds in the Philippines, majority of which is red seaweed with six hundred (600) taxa, followed by green seaweed with two-hundred seventy two (272) taxa, and brown seaweed with one hundred ninety-three (193) taxa. In the present study, ninety-four (94) taxa or species were identified and collected which will be added as new record or information on the available literature of marine flora and presence of some macroalgae in Northern Samar.

There were thirty-seven (37) green seaweeds that were collected in all three (3) sampling sites. Green algae/ seaweeds have a variety of living areas including coral debris, muddy corals, dead

corals, muddy sand, mangroves (Harborne *et al*, 2006; Ghazali *et al*, 2018; Cahyanto *et al*, 2019) one very best example of this, is the genus *Ulva* that have higher abundance than other genus in the intertidal zone. The genus *Ulva* had filamentous thallus that can withstand environmental stress. This genus is also known to be able to live on saltwater or freshwater and has a high salinity tolerance and high reproductive capability (Cahyanto *et al.*, 2019). The abundance of *Ulva* are also dictated by other two factors like nuisance due to anthropogenic activities (Littler and Littler, 1984; Choiet *al.*, 2008; Kim *et al.*, 2016; Baldia *et al.*, 2017) and the effectinfluence of lunar cycle (Lee, 2008). It was observed that all sampling sites have residential areas near the beaches, effluent from houses which contains organic waste might contribute to the abundance of members of family Ulvaceae (White, 1987), since *Cladophora* and *Ulva* are dominant and can thrive in an unstable or polluted environment (Littler and Littler, 1984; Choiet *al.*, 2008; Kim *et al.*, 2016; Baldia *et al.*, 2017). Also, as shown in Table 1 and 2, the genus *Caulerpa* was represented by seven (7) species, in Brgy. Guinarawayan, Allen, *Caulerpa racemosa* was abundantly growing near the reef-edge and gleaners were also observed harvesting *C. racemosa*, it was also observed that the *C. racemos* present in the area have strong short erect branches with crowded ramuli with spherical tips, which is a clear indication that the area is exposed to strong wave actions and current (Trono, 1997). However they are only distributed in the lower intertidal zone due to environmental stress (e.g. desiccation, temperature extremes, and inundation by rain), thus, desiccation sets the upper limit to the species distribution of the genus *Caulerpa* (Cecil *et al.*, 2004).

In addition, the genus *Halimeda* was also composed of five (5) species with *H. macroloba* and *H. opuntia* being two of the most abundant (Fig. 2) and distributed seaweed in almost eight (8) or nine (9) sampling sites, since as a matter of fact *Halimeda* is one of the most abundant algal taxa in the tropical seas (Price *et al.*, 2014). The presence of aragonite which is a natural form of Calcium carbonate on its skeletal structure which is a form plasticity that enables certain species of *Halimeda* to adapt to fluctuating environmental conditions (Vroom *et al.*, 2003; Price *et al.*, 2014) that might contribute to their great abundance in the sampling sites.

On the other hand, red macroalgae were also present in all sampling sites, as a matter of fact they are the more diversified when it comes to number of genera which accounts for about twenty-seven (27), but when it comes to number of species, there were thirty-four (34) in all sampling sites. According to Lee (1999), there are more red seaweeds as compared to any major seaweed groups in the world, this was supported by Lastimoso and Santianez (2020) that 56 % of seaweeds in the country is red seaweeds. Although they can be seen in all latitudes, but red macroalgae are limited to polar and subpolar regions of the world (Baldia *et al.*, 2017). As observed, this group were present in current exposed area due to their multicellular rhizoidal holdfast that serves as a strong attachment of thalli to substrates. In addition, Rhodophyceae have multiple life stages that help in their adaptation for survival. Their triphasic life cycle also explains why compared to brown and green algae, red algae are the most adaptable and morphologically diverse (Lee, 2008; Baldia *et al.*, 2017). As shown in Table 2, *Tricleocarpa fragilis* were present in seven (7) sampling sites by which they are observed growing abundantly, Islam *et al* (2019) has recounted that *T. fragilis* is commonly found attached to rocks, dead corals and shells in shallow areas, moderately exposed to wave action, where it forms large solitary clumps.

In addition, the least group of macroalgae was brown seaweeds which accounts for twenty-three (23) species. As mentioned by Lee (2008) that majority of the brown algae are found in

temperate region rather than in tropical region, it is expected that Philippines has low species diversity of brown algae. Even though there is a low species diversity of brown seaweeds in the province, they can be seen to have the greatest cover in all sampling sites. Most brown algae grows on rocks/ hard substratum which is commonly observed in all sampling sites. Brown macroalgae exhibits larger biomasses (especially *Sargassum*) as compared to other group of marine macroalgae (green and red) which is evident due to their larger size (Johnston, 1969; Baldia *et al.*, 2017). *Sargassum* is the most represented brown macroalgae which is composed of nine (9) species, identification of *Sargassum in situ* is difficult due to its phenotypic plasticity (Gouvea *et al.*, 2022), aside from its variability *Sargassum* have tougher thallus, strong holdfast, and air bladder as compared to other seaweeds that makes them adapt to the harsh wave actions and currents (Hurtado, 1999).

The presence of genus *Padinain* almost all of the sampling sites as well as its variability in cover, might be indicative of anthropogenic nutrient loading from domestic and agricultural runoff. All sites were near populated areas with poor sewage system, and a few backyard pig pens were observed in some of the barangays that served as sampling sites. Fresh and dried biomass of *Padina* have high adsorbing potential for various pollutants (Ansari *et al.*, 2019). Aside from being indicator the wide distribution of the *Padina* can be suspected to its lifecycle and reproduction which is through sporophyte generation. The sporophyte dominance may be due to its greater resistance (longevity of individuals) to water movement that helps them establish their population (Allender, 1977: Baldia *et al.*, 2017), this might be the reason of its great abundance and distribution in almost all sampling sites.

As shown in Table 3, the sampling site which had the lowest pH was detected in barangay Balicutro in Lavezares which only had 7.2 which is lower against the threshold range for seawater pH of 7.5-8.5 (Water watch, 2022). This lower pH in this site could be attributed to lesser number of seaweeds in the area by which there were only twelve (12) identified seaweeds. According to Water watch (2022), human activities such as sewage overflows or runoff, can cause significant short-term fluctuations in pH and long-term impacts can be extremely harmful to plants and animals. Extreme changes in pH, can stress local organisms and may ultimately lead many species to leave the area or die. Aside from pH, salinity and seawater temperature can also be affected by the human activities. The average salinity of ocean water is 35 ppt, but the range from 30-35 ppt can still considered as accepted, as much as possible monitoring of salinity is very important since salinity levels control local species composition. Changes in salinity in a specific area/ region can occur as a result of weather patterns, such as droughts or storms, or they can alert us to events such as increased urban runoff and sewer discharge (Water watch, 2022). Also, as shown in Table 3, the temperature value in all sampling sites were within the normal range of 26-28°C, however, there are some variations of seawater temperature depending on the sites, since as mentioned by Johnson & Xie (2010) that the troposphere may have become less stable and casting doubts on the possibility that the sea surface temperature threshold increases substantially with global warming, by which it can be correlated to human activities.

Rocky substrate was observed to be common in all sampling sites, it is much preferable for the seaweeds to grow (Tsuda, 1972; Wreede, 1976; Baldia *et al.*, 2017) since it gives them a hard attachment. The most common seaweed genera collected near the rocky substrate was *Sargassum* by which even at the reef edge they grow abundantly since according to Hurtado (1999) that macroalgae found in exposed intertidal zone are tougher and shorter with

narrower thalli, therefore they are more prone to wave motion compared to those growing in calmer areas.

CONCLUSION AND RECOMMENDATIONS

Based on the result of the study there were more seaweeds that belongs to Chlorophyta, followed by Rhodophyta, and the least was Phaeophyta. In terms of the municipality with the most number of seaweeds, Lavezares had the most with sixty (60) taxa identified, followed by Biri with forty-four (44), and Allen with thirty-four (34) species. With the result of the measurement of the environmental parameters, pH of seawater really determines the availability seaweeds since some of them are sensitive to variation of pH. As a matter fact it is evident in barangay Balicutro, Lavezares wherein the pH value fell below the threshold limit and only few seaweeds were collected during the sampling. Thus, based on the above mentioned result, the availability of seaweeds in an area can be dictated if the environmental parameters favor for the growth and survival of some seaweeds.

It is recommended to conduct similar studies in other island municipalities of Northern Samar, to determine the uniqueness of the taxa in the province.

COMPETING INTERESTS

The authors declare no competing interests.

ACKNOWLEDGMENT

The authors would like acknowledge the participation of LGU especially the Municipal Agricultural Office (MAO) and the Office of the Mayor in all three (3) municipalities for the release and approval of prior informed consent (PIC) prior to sampling. Also, the participation of some individuals during the collection of seaweeds in the sampling sites. Lastly, great appreciation to the CHED-LAKAS for funding.

REFERENCES

- Allender B.M.(1977). Effects of emersion and temperature upon growth of the tropical brown alga *Padina japonica*, using a tide-simulation apparatus. *Marine Biology*, 40:95–100.
- Ansari, A.A., Ghanem, S.M., Naeem, M. (2019). Brown alga *Padina*: A review. *International Journal of Botany Studies*, 4(1); 01-03.
- Baldia, S.F., Kabling, M.T., Gabriel, C.A., Abeleda, A.N., Aguinaldo, R.C., Clemente, K.J., Cordero Jr., P.A. (2017). Marine macroalgal diversity assessment of Biri Island and Dalupirit Island, Northern Samar. *Acta Manilana*, 65: 29-38.

Baweja, P., Kumar, S., Sahoo, D., and Levine, I. (2016). Biology of seaweeds in Seaweed in Health and Disease Prevention. Academic Press, Cambridge, MA, USA, 41-106

Calumpong, H.P. and Menez, E.G. (1997). Field guide to the common mangroves, seagrasses, and algae of the Philippines. Bookmark, Inc.

Cahyanto, T., Paujiah, E., Rinasti, I., Suryani, Y., Adawiyah, A., & Ulfa, U.A. (2019). Abundance of green algae in the intertidal zone of Karang Papak Coastal, West Java, Indonesia. *Journal of Physics: Conference Series*. <https://doi.org/10.1088/1742-6596/1402/3/033027>.

Cecil, W., Olsen, K., Shrimpton, S., Wimpee, L., Lesichner, J., Osborne-Koch, M., Yamada S., & Helms, A. (2004). Intertidal zonation: does species diversity decrease with tidal height?. <https://scholarsbank.uoregon.edu/xmlui/bitstream/handle/1794/7974/Yamada.pdf?sequence=1&isAllowed=y>.

Choi, H., Lee, K., Wan, X.Q., Yoo, H., Park, H., Kim, J., Chung, I. (2008). Temporal variations in seaweed biomass in Korean Coasts: Woejodo and Jusamdo, Jeonbuk. *Algae*, 23(4): 335-342. <https://doi.org/10.4490/algae.2008.23.4.335>.

Choudhary, B., Chauhan, O.P., Mishra, A. (2021). Edible seaweeds: A potential novel source of bioactive metabolites and nutraceuticals with human health benefits. *Frontiers in Marine Science*. <https://doi.org/10.3389/fmars.2021.740054>.

Cordero Jr., P.A. (1977). Studies on Philippine marine red algae. Special Publications from the s

Dagalea FMS, Cui-Lim KMR. 2018. Antimicrobial properties of biocomposite films from kappa-Carrageenan (kC) filled with Nanorod-rich Zinc Oxide (ZnO-N). *Advance Pharmaceutical Journal*, 3(2). <https://doi.org/10.31024/apj.2018.3.2.5>

Dagalea, F. M. S., & Cui – Lim, K. M. R. (2021). Analysis of Biofilms from Biosynthesized Zinc Oxide Nanoparticles. *International Research Journal of Pure and Applied Chemistry*, 22(1), 92–97. <https://doi.org/10.9734/irjpac/2021/v22i130381>

Dumilag, R.V. (2019). Edible seaweeds sold in the local public markets in Tawi-Tawi, Philippines. *Philippine Journal of Science*, 148(4): 803-811.

Dumilag, R.V., and Javier, R.F. (2022). Ethnobotany of medicinal seaweeds of Ilocos Norte, Philippines. *Philippine Journal of Science*, 151(3): 1135-1156.

Galenzoga, D. (2016). Community Structure of Molluscs in Northern Samar, Philippines. *International Journal of Life Sciences Research*, 4(1): 100-104.

Galenzoga, D. (2019). Seaweed resources in the island town of Biri, Northern Samar. *J. Bio. Innov.*, 8(3):360-363.

Ghazali, M., Kurnianingsih, R., Astuti, S., Sukiman, P. (2018). Jenis Alga Hijau (Chlorophyta) pada ekosistem hutan mangrove di dusun ekas, kabupaten Lombok Timur. *Fish Scientiae*, 8(1) 11-23.

Gouvea, L.P., Horta, P.A., Fragkopoulou, E., Gurgel, C.F., Peres, L.M., Bastos, E., Ramlov, F., Burle, G., Koerich, G., Martin, C.D., Serrao, E.A., Assis, J. (2022). Phenotypic plasticity in *Sargassum* forests may not counteract projected biomass losses along a broad latitudinal gradient. *Ecosystem*, 29: 29-41.

Harborne A.R., Mumby P.J., ZŻychaluk, K., Hedley, J.D., and Blackwell, P.G.(2006). Modeling the beta diversity of coral reefs. *Ecology*, 87(11): 2871-2881.

Hurtado A.Q. (1999). Sargassum studies in Currimao, Ilocos Norte, Northern Philippines I. Seasonal variations in the biomass of *Sargassum carpophyllum* J. Agardh, *Sargassum ilicifolium* (Turner) C. Agardh and *Sargassum siliquosum* J. Agardh (Phaeophyta, Sargassaceae). *Southeast Asian Fisheries Development Center*. 42:321–25

Islam, M., Hoq, E., Haque, M.A., Khan, MS., Hazan, Z., Ali, M.Z., & Mahmud, Y. (2019). Seaweeds of Bangladesh coast: abundance, distribution, and taxonomic list. *Bangladesh Fisheries Research Institute, Ministry of Fisheries and Livestock*.

Johnson, N.C, & Xie, S.P. (2010). Changes in the sea surface temperature threshold for tropical convection. *Nature Geoscience*, 3:842-845. <https://www.nature.com/articles/ngeo1008/>

Kumar, Y., Tarafdar, A., Badgujar, P.C. (2021). Seaweed as a source of natural antioxidants : therapeutic activity and food applications. *Journal of Food Quality*. <https://doi.org/10.1155/2021/5753391>.

Lastimoso, J.M., and Santianez, WJ. (2020). Updated checklist of the benthic marine macroalgae in the Philippines. *Philippine Journal of Science: Special Issue on Biodiversity*, 150(1): 29-92.

Lee R.E. (1999). *Phycology*. UK: University Press, Cambridge.

Lee, R.E. (2008). *Phycology 3rd Edition*. University Press, Cambridge.

Litler, M.M., Litler, D.S. (1984). Relationships between macroalgal functional form groups and substrata stability in a sub-tropical rocky-intertidal system. *Journal of Experimental Marine Biology and Ecology*, 74(1):13-34. [https://doi.10.1016/0022-0981\(84\)90035-2](https://doi.10.1016/0022-0981(84)90035-2).

Menez, E.G. (1961). The marine algae of the Hundred Islands, Philippines. *Philippine Journal of Science*, 90:37-86.

Pacturan, J.E. (2016). Eastern Samar seaweed industry assessment and proposed development interventions for smallholder growers. Report for the AsiaDHRRA Seaweed Study for Eastern Samar.

PhilAtlas. (2023). Northern Samar. <https://www.philatlas.com/visayas/r08/northern-samar.html#sources>.

Price, N.N., Hamilton, S.L., Tootell, S.T., Smith, J.E. (2014). Species-specific consequences of ocean acidification for the calcareous tropical green algae *Halimeda*. *Marine Ecology Progress Series*, 440: 67-78. <https://doi:10.3354/meps09309>

Sherwood, A.R., and Guiry, M.D. (2023). Inventory of seaweeds and seagrasses of the Hawaiian Islands. *Polynesian Seaweeds Biodiversity, Biogeography, and Evolution*, 12(2). <https://doi.org/10.3390/biology12020215>

Trono, G.C. (1972). Notes on some marine benthic algae in the Philippines. *Kalikasan, Philippine Journal of Biology*, 1: 126-147.

Trono, G.C. (1997). Field guide and atlas of the seaweed resources of the Philippines. Bookmark, Inc., Makati City, Philippines, 97.

Tsuda R. T. (1972). Morphological, zonal and seasonal studies of two species of *Sargassum* on the reefs of Guam. In: Proceeding of the 7th International Seaweed Symposium, 40–44. (Sapporo, Japan: University of Tokyo Press).

Vroom P.S., Smith, C.M., Coyer, J.A., Walters, L.J., Hunter, C.L., Beach, K.S., Smith, J.E. (2003) Field biology of *Halimeda tuna* (Bryopsidales, Chlorophyta) across a depth gradient: Comparative growth, survivorship, recruitment, and reproduction. *Hydrobiologia*, 501: 149–166.

Waterwatch. (2022). pH, salinity, and temperature. <https://sfyl.ifas.ufl.edu/media/sfylifasufledu/miami-dade/documents/sea-grant/Temperature,-Salinity-and-pH.pdf>

White, A. (1987). Philippine Coral Reefs A Natural Guide. New Day Publishers, Quezon City 3008, 91 .

Wreede R.E.(1976). The phenology of three species of *Sargassum* (Sargassaceae, Phaeophyta) in Hawaii. *Phycologia* 1976; 15(2):175–83. DOI:10.2216/i0031-8884-15-2-175.1