

Phenolic compound content and antioxidant activity of extracts from the leaves of *Anogeissus leiocarpus* (Combretaceae), a plant used in the north of Côte d'Ivoire for the traditional treatment of gastrointestinal disorders in broiler chickens.

Abstract :

Anogeissus leiocarpus (DC.) Guill. & Perr. (Combretaceae) is a plant found in West Africa, from Senegal to Cameroon and extending to Ethiopia and East Africa. In the north of the Ivory Coast, this plant is commonly used by livestock farmers to treat various pathologies, including digestive and gastrointestinal disorders.

The aim of this study was to determine the phenolic compound content and antioxidant properties of 70% hydroethanol and aqueous extracts of *A. leiocarpus* leaves. Phenolic compound contents were determined by spectrophotometric methods. (Marinova and al and WOOD and al). Antioxidant activities were assessed using ABTS, DPPH and FRAP tests. (Choong and al ; Parejo and al ; Le andal).

The 70% hydroethanolic and aqueous extracts of *A. leiocarpus* presented roughly equivalent concentrations of total polyphenols (608.41 and 558.25 mg EAG/g respectively). However, the hydroethanolic extract was richer in flavonoids (539.60 mg EQ/g) than the aqueous extract (388.95 mg EQ/g). In addition, the average inhibition rate of the ABTS radical in the presence of the 70% hydroethanol extract was 11.56 μ M Trolox equivalent/g, compared with 8.32 μ M Trolox equivalent/g with the aqueous extract. The average chelation percentages were 31.93% (70% hydroethanol extract) and 20.92% (aqueous extract). As for the reduction of the DPPH radical, the IC₅₀ for the 70% hydroethanol and aqueous extracts were 4.4 mg/mL and 6.0 mg/mL respectively.

This study confirms that *Anogeissus leiocarpus* is a plant containing phenolic compounds. The presence of these phenolic compounds in this plant, together with its antioxidant activities with low values (IC₅₀), could justify its widespread use by livestock farmers in northern Côte d'Ivoire, particularly those rearing broiler chickens.

INTRODUCTION

The traditional African pharmacopoeia constitutes a veritable phytopharmacy that can be used in several areas of health care, for both humans and animals and even for plant (Agaie and al, Belemnaba and al, 2007, Souad andal 2020, Mario and al 2014). In the specific case of animals, various studies have shown the use of plant-based remedies for their therapeutic management.

Formatted: English (United States), Highlight

Comment [N1]: *Anogeissus leiocarpa* (DC.) Guill. & Perr.

Formatted: Highlight

Formatted: English (United States), Highlight

Formatted: English (United States)

Comment [N2]: *Anogeissus leiocarpa* (DC.) Guill. & Perr.

review and update the name *Anogeissus leiocarpa* (DC.) Guill. & Perr.

also, make a brief presentation of the study's problem that justifies the basis of this study

Formatted: English (United States)

Formatted: English (United States)

Formatted: Left, Space After: 10 pt

Comment [N3]: Incongruous citations and references in the summary

Formatted: English (United States)

Formatted: Space After: 10 pt

Formatted: Font: 11 pt, English (United States)

Formatted: English (United States)

Comment [N4]: please formulate a transition between the two parties that justifies the problem

Formatted: English (United States)

Anogeissus leiocarpus (DC.) GUILL. & PERR. (Combretaceae) is a plant found in West Africa, from Senegal to Cameroon and extending to Ethiopia and East Africa (Arbab, 2014). Growing in dry forests and gallery forests (Arbab, 2014; Ouédraogo and al., 2013), its distribution extends from the Sahara to the outer layer of tropical rainforests. The decoction of *Anogeissus leiocarpus* leaves is used to treat small ruminants suffering from gastrointestinal parasitism, according to an ethnoveterinary study conducted in Burkina Faso (Githiori and al., 2005). In the north of Côte d'Ivoire, this plant is commonly used by livestock farmers to treat various pathologies including digestive and gastrointestinal disorders (Kone and al., 2019). Julienne and al (2021) confirmed in their studies that *A. leiocarpus* is used in veterinary medicine, particularly in the treatment of parasitic diseases caused by *Haemonchus contortus*. The methanolic extract of *A. leiocarpus* stem bark has also demonstrated anti-trypanosomiasis activity against four strains of Trypanosoma (Shuaibu and al., 2008) as well as leishmanicidal activity (Shuaibu et al., 2008b). *A. leiocarpus* also has strong antihelminthic activity (Juliennea and al., 2021).

In our previous work, we (Ouattara and al., 2023) evaluated the *in vitro* efficacy of *A. leiocarpus* on multi-resistant strains of *Salmonella typhimurium* isolated from the droppings of farmed broilers (Ouattara and al., 2023). Extracts from this plant showed good antimicrobial activity against these multi-resistant *Salmonella typhimurium* germs. Our study also showed the presence of several groups of secondary metabolites, differently distributed, with high levels of phenolic compounds in extracts from the leaves of this plant (Ouattara and al., 2023). Consequently, if farmers are to make better use of this plant in broiler rearing, there is need to design an improved phytomedicine to help combat microbial resistance and oxidative stress effectively. In fact, oxidative stress is unavoidable in poultry production and affects the physiological, behavioural and biochemical state of growing chickens, which can cause their death or deteriorate meat quality (appearance, texture, juiciness, tenderness and odour) (Ali and Li, 2021). Furthermore, curing an animal treated in a farm involves not only eliminating the germ, but also reducing the signs of morbidity generally caused by the massive production of free radicals in the affected organism (Djeridane and al, 2006). However, several synthetic antioxidants are used by broiler breeders as feed additives to combat oxidative stress. Although these enriched food supplements are highly effective against free radicals, they are unfortunately likely to have side-effects and may even be toxic (Maman and al, 2008, Bramen 1975 ; Bougandoura 2012). Plants constitute a natural reservoir of secondary metabolites (Lee and al, 2000 ; Cakir and al 2003 ; Kumaran and al 2007), and have always been heavily involved in the search for new natural antioxidants that

Formatted: English (United States)

Comment [N5]:
please cite the work reference

Formatted: English (United States)

Formatted: English (United States)

are effective and have very few side effects. With this in mind, it would be a great advantage if a plant-based formulation with antioxidant properties could be made available to livestock farmers in the long term. This is the reason why, we conducted the present study to assess the phenolic compound content and antioxidant properties of two extracts of *A. leiocarpus*, a plant from the Ivorian pharmacopoeia that is commonly used by farmers in northern Côte d'Ivoire to treat gastrointestinal disorders in broiler chickens.

MATERIALS AND METHODS

Plant material

It consists of the leaves of *Anogeissus leiocarpus*. They were harvested in the village of Lataha in the Korhogo region (northern Côte d'Ivoire) in March 2022 and authenticated by the Centre National Floristique of the Felix HOUPHOUËT-BOIGNY University in Cocody-Abidjan.

Preparation of aqueous and 70% hydroethanolic extracts of *Anogeissus leiocarpus*

The leaves of *Anogeissus leiocarpus* were washed, cut and dried in the shade for a fortnight. Once dried, the plant material was ground. 100g of this powder was mixed with 1 litre of distilled water. The mixture was homogenised at room temperature in the laboratory using a magnetic stirrer for 24 hours. The homogenate obtained was filtered twice on cotton wool and once on Whatman paper (3 mm). The volume of the filtrate obtained was reduced using a Med Center Venticell oven at 50°C to give a powder that constitutes the total aqueous extract (E.H₂O) (Ouattara and al., 2013).

The same operation was carried out using 70% ethanol instead of distilled water, to obtain the 70% hydroethanol extract (E.HOH) (Zirihi and al., 2003). The two plant extracts obtained were stored in the refrigerator for further testing.

Determination of total polyphenol content

The method of Wood and al (2002) was used for the determination of total polyphenols content. A volume of 2.5 ml of diluted (1/10) Folin-Ciocalteu reagent was added to 30 µL of ethanolic extract. The mixture was kept for 2 min in the dark at room temperature, then 2 mL of calcium carbonate solution (75 g.L⁻¹) was added. The mixture was then placed in

Comment [N6]: We expected to see the prevalence of avian diseases and their consequences on production and the national economy.

Because for us, the study would like to either improve the quality of production or implement an effective protocol of traditional veterinary care, and this is what constitutes the central question

Formatted: English (United States)

Formatted: English (United States)

Comment [N7]: Some ecological characteristics of the sample collection environment; the types or parts of organs removed.

Formatted: English (United States)

Comment [N8]: Why only on dry leaves, without arrow leaves or even stem bark which are very often used in traditional veterinary medicine?

Formatted: English (United States)

Formatted: English (United States)

a water bath at 50°C for 15 min and rapidly cooled. Absorbance was measured at 760 nm, using distilled water as the blank. A calibration line was made with gallic acid at different concentrations. Analyses were carried out in triplicate and the concentration of polyphenols was expressed in milligrams of gallic acid equivalent per gram of extract (mg GAE/g extract).

Formatted: English (United States)

Determination of total flavonoid content

The method of **Marinova and al (2005)** was used to determine total flavonoids. In a 25-mL flask, 0.75 mL of 5% (w/v) sodium nitrite (NaNO₂) was added to 2.5 mL of extract. 0.75 mL of 10% (w/v) aluminium chloride (AlCl₃) was added to the mixture and incubated for 6 min in the dark. After incubation, 5 mL of sodium hydroxide (1N NaOH) was added and the volume was made up to 25 mL. The mixture was shaken vigorously before being assayed using a UV-visible spectrophotometer. The reading was taken at 510 nm. A calibration line was constructed with quercetin at different concentrations. Trials were carried out in triplicate. Flavonoid content was expressed as milligram quercetin equivalent per g extract (mg EQ /g extract).

Determination of the *in vitro* antioxidant activities of aqueous and 70% hydroethanol extracts.

The 2,2'-azinobis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) assay

Formatted: English (United States)

The ABTS radical method is one of the most widely used tests for determining the concentration of free radicals. It is based on the neutralization of a radical - cation resulting from the mono electronic oxidation of the synthetic chromophore 2,2'- azino-bis (3 - ethylbenzothiazoline -6- sulfonic acid)This method is based on the ability of the compounds to reduce the ABTS^{•+} (2,2' azinobis 3 ethylbenzothiazoline 6 sulphonic acid) radical cation. The test was carried out using the method described by **Choong and al (2007)**. The ABTS^{•+} radical-cation was produced by reacting 8 mM ABTS (87.7 mg in 20 mL distilled water) and 3 mM potassium persulphate (0.0162 g in 20 mL distilled water) in a 1 :1 (v/v) ratio. The mixture was then incubated in the dark at room temperature for 12-16 hours. This ABTS^{•+} solution was diluted with methanol to obtain a solution with an absorbance of 0.7 ± 0.02 at 734 nm. A test portion of 3.9 mL of this diluted ABTS^{•+} solution was added to 100 µL of the test compound. After shaking, the mixture was incubated for 6 min in the dark (T=30±2°C). The residual absorbance of the ABTS^{•+} radical was then measured at 734 nm using a UV-visible spectrophotometer and was expected to be between 20% and 80% of the absorbance of the blank. The tests were carried out in triplicate and the results were expressed in µmol

Formatted: English (United States)

Trolox equivalent per litre of extract ($\mu\text{mol TE/L}$). A calibration line was performed with the following concentrations of Trolox : $0.375\mu\text{M}$; $0.5\mu\text{M}$; $0.625\mu\text{M}$; $1\mu\text{M}$; $1.125\mu\text{M}$, $1.375\mu\text{M}$ and $1.5\mu\text{M}$ and the inhibition rate (% I) of ABTS^o+ was expressed as follows (1) :

$$\% I = [(Abs_control - Abs_extract) / Abs_control] \times 100 \quad (1)$$

Abs_control = diluted ABTS absorbance,

Abs_extract = diluted ABTS absorbance + sample,

Antioxidant concentration or activity(2) :

$$(\mu\text{M}\acute{\text{e}}\text{q Trolox}) = (\% I \times d) / (4,99 \times 10) \quad (2)$$

Iron chelation (FRAP)

The chelating capacity of the extracts was measured using the method of **Le and al (2007)**. Solutions of extracts and the reference antioxidant (EDTA 500 μl) were initially mixed with 100 μl FeCl_2 (0.6 mM in distilled water) and 900 μl methanol. After 5 min, 100 μl Ferrozine (5 mM in methanol) was added to the reaction medium. The mixture was shaken well and then left to react for 10 min at room temperature, allowing complexation of the residual iron and formation of a chromophore with an absorption maximum at 562 nm. The negative control contains all the reagents except the test sample, which is replaced by an equal volume of methanol. Readings are taken at 562 nm against a methanol blank. The sequestering effect of the samples on iron is expressed as a percentage of chelation according to the following equation (3) :

$$\% \text{ Chelation} = [(Abs \text{ control} - Abs \text{ test}) / Abs \text{ control}] \times 100 \quad (3)$$

2, 2'-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay

The anti-free radical activity of the two plant extracts was measured using the 2, 2'-diphenyl-1-picrylhydrazyl (DPPH) test according to the method of **Parejo and al. (2000)**. A range of concentrations (0-200 $\mu\text{g/mL}$) of plant extract or gallic acid (reference antioxidant) was prepared. A volume of 2.5 mL of this solution was mixed with 2.5 mL of DPPH (100 μM) prepared in methanol. After homogenisation, the mixture was incubated at room temperature (25°C) in the dark. After 15 min incubation, the absorbance was read at 517 nm against a

Formatted: English (United States)

Formatted: Font: 11 pt, English (United States)

Formatted: English (United States)

Formatted: Font: 11 pt, English (United States)

Formatted: English (United States)

Formatted: English (United States)

"blank" containing only methanol. The percentage inhibition of the DPPH radical was calculated using the following equation :

$$\text{DPPH inhibition (\%)} = (1 - (\text{test OD} / \text{blank OD})) \times 100 \quad (4)$$

The IC₅₀ is the concentration of plant extract or quercetin responsible for 50% inhibition of DPPH radicals. It is determined from the graph showing the percentage of DPPH inhibition as a function of extract and Gallic Acid concentrations.

Statistical analysis

Data analysis and graphical representation were carried out using Graph Pad Prism 8.0.1 (Microsoft, USA). The mean value is accompanied by the standard error of the mean (Mean ± SEM). The difference between two values is considered significant when $P < 0.001$. The results were statistically analysed using a two-way analysis of variance (ANOVA 2)

RESULTS

Phenolic compound content

Determination of phenolic compounds in the two extracts from *A. leiocarpus* leaves showed that both the hydroethanol extract (E.HOH) and the aqueous extract (E.H₂O) had roughly equivalent concentrations of total polyphenols (608.41 and 558.25 mg EAG/g respectively) (Figure 1). In terms of total flavonoids, the hydroethanol extract of *A. leiocarpus* leaves was richer (539.60 mg EQ/g) than the aqueous extract (388.95 mg. EQ/g) (Figure 2).




Figure 1 : Total polyphenol content of extracts from *A. leiocarpus*

E.HOH = hydroethanolic extract 70% ; E.H₂O = Aqueous extract ; ** p (= 0,9404) > 0,005 : différence not significant

Formatted: English (United States)




Figure 2 : Total flavonoid content of extracts from de *A. leiocarpus*

E.HOH = Hydro éthanolic extract 70% ; E.H₂O = Aqueous extract ; ** p (= 0,0027) < 0,005 : significant difference

Formatted: English (United States)

UNDER PEE

ABTS test: antioxidant capacity

The ABTS test, measures the antioxidant capacity in solutions of pure substances or aqueous mixtures. Comparative analysis of these two substances reveals significant differences. For a concentration of 100 mg/mL, the average inhibition rate of the ABTS radical in the presence of the 70% hydroethanol extract was 11.56 μ M Trolox eq/g, whereas it was 8.32 μ M Trolox eq/g with the aqueous extract. For these two extracts, the inhibition rates obtained were much lower compared to that of gallic acid, which recorded a value of 64.71 μ M Trolox eq/g (Table 1).

Chelating power (FRAP)

The average chelation percentages were 31.93% (70% hydroethanol extract) and 20.92% (aqueous extract). However, these values are significantly lower than those obtained with the reference antioxidant (EDTA), whose average chelation percentage was estimated at 96.74%. These results are shown in Table 1.

Table 1 : Anti oxidant activities of *A. leiocarpus* extracts by ABTS and FRAP tests

Extract or Reference molecule	Assays	
	ABTS (mM Eq Trolox/g)	FRAP (% of chelation)
E.H2O	8,32 \pm 0,199	20,92 \pm 0,195
E.HOH	11,56 \pm 0,535	31,93 \pm 0,078
Gallic Acid	64,71 \pm 0,519	
EDTA		96,74 \pm 0,275

E.HOH = 70% hydro ethanolic extract; E.H2O = Aqueous extract

DPPH test

The results of the DPPH radical reduction assay are shown in Figures 3 and 4. The concentrations corresponding to 50% inhibition (CI₅₀) for the hydroethanolic (4.4 mg/mL) and aqueous (6.0 mg/mL) extracts of *A. leiocarpus* are higher than that of the reference antioxidant (Gallic acid), which was 3.6 mg/mL. The IC value₅₀ is inversely proportional to the percentage of DPPH inhibition, so a lower IC₅₀ corresponds to higher antioxidant activity. The two extracts therefore showed significant antioxidant powers, but relatively lower than that of gallic acid (the reference antioxidant).

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font: 11 pt, English (United States)

Formatted: English (United States)

Formatted: English (United States)

Comment [N9]: the results are quite clear but we can please improve the quality of analysis

Formatted: English (United States)

Formatted: English (United States)




Figure 3: Scavenging ability of extracts by DPPH test

E.HOH : Hydroethanolic extract 70% ; E.H2O : Aqueous extract

Formatted: English (United States)




Figure 4: IC₅₀ values of *A. leiocarpus*' extracts by DPPH test

E.HOH : hydro ethanolic extract 70% ; E.H2O : Aqueous extract

Formatted: English (United States)

DISCUSSION

Determination of the phenolic compounds in the two extracts showed that the hydro-ethanolic and aqueous extracts of *A. leiocarpus* leaves had roughly equivalent concentrations of total polyphenols (608.41 and 558.25 mg EAG/g respectively). In terms of total flavonoid content, the hydro-ethanolic extract of *A. leiocarpus* leaves contained a higher level (539.60 mg EQ/g) than the aqueous extract (388.95 mg EQ/g).

Our results are in line with those of **Koné andal in 2006**, who showed that *A. leiocarpus* leaves are rich in total polyphenols and flavonoids. The work by these authors, which looked at a total of six plants, showed that *A. leiocarpus* contained the highest level of total polyphenols (223.13 mg EAG/g, i.e. 4 times more than the leaves of *Vepris heterophylla*, which contained the lowest level of these compounds. This total polyphenol content is still much lower than that which we obtained, which varied from 558.25 mg GAE/g (aqueous extract) to 608.41 mg GAE/g (hydroethanol extract).

In the study by the same authors, the trunk bark of *A. leiocarpus* contained only 26.53 mg EAG/g of polyphenols and 10.30 mg EQ/g of flavonoids. These levels of total polyphenols and flavonoids are also widely lower than those obtained in the present study. *Terminalia macroptera* root bark showed the highest antioxidant activity, followed by *A. leiocarpus* leaf bark in tests using ABTS and DPPH, with respective values of 468.8 μM Eq Trolox/g and 350.30 μM Eq Trolox/g (ABTS) and 361.60 mg/mL and 271.87 mg/mL (DPPH) (**Koné andal, 2009**). **Gheldolf andal, 2002 ; Holasova andal, 2002 and Kumaran andal, 2005** have shown that there is a linear correlation between total polyphenol content and antioxidant activity in these plants.

In another study by **Barku andal (20013)**, the antioxidant activity of methanolic extracts of 5 medicinal plants was assessed using the DPPH test and the ferric ion reducing power (FRAP) test. These were *Amaranthus Spinosus*, *Anogeissus leiocarpus*, *Spondia monbin*, *Corchorus olitorius* and *Mallotus oppositifolia*. The results showed that all these plants have antioxidant activity. However, *Anogeissus leiocarpus* showed the highest antiradical activity ($95.86 \pm 0.1\%$) followed by *Cochorus olitorius* ($94.19 \pm 0.06\%$), as percentages of DPPH radical inhibition ; while *Amaranthus spinosus* recorded the lowest activity ($40.87 \pm 2.5\%$) on the one hand and had the highest reducing power followed by *Spondia monbin* ; *Amaranthus spinosus* having the lowest reducing power, on the other hand. The leaves of *A. leiocarpus* also had the highest levels of total polyphenols (1294.81 ± 30 mg GEA) and flavonoids (540.23 ± 24.5 mg/g), while *Amaranthus spinosus* recorded the lowest levels of these same compounds with

Comment [N10]: in reality these results are not at all comparable. review the terminology used because 223.13 is two to three times lower than the values obtained in this study.

the comparison would be justified if within the framework of this study we compared values on several plant specie

Formatted: English (United States)

Comment [N11]: How can you justify these very significant differences on the same types of organs of the same species. Did you use the same extraction and dosage methods?

Formatted: English (United States)

Formatted: English (United States)

Comment [N12]: Unclear discussion, to be fundamentally reviewed,

Formatted: English (United States)

Formatted: English (United States), Highlight

Formatted: English (United States)

the following respective values: 48.01 ± 2.0 mg GEA and 63.16 ± 107 mg/g. Through this work, these authors have shown that the leaves of *Anogeissus leiocarpus* are rich in total polyphenols and flavonoids on the one hand and constitute an important source of antioxidants on the other.

This study confirms that *Anogeissus leiocarpus* is a plant that contains phenolic compounds. It also justifies our previous work, which consisted of searching for secondary metabolites in hydroethanol and aqueous extracts of *Anogeissus leiocarpus*. Using the triphytochemical method, this study revealed the presence of phenolic compounds, cardiac glycosides, saponins, sterols, terpenes and alkaloids (Ouattara and al., 2023). However, this study also shows that the content of phenolic compounds, flavonoids and antioxidant activities in *Anogeissus leiocarpus* varies from one extract to another, from one organ to another, from one region to another and from one methodology to another. These remarks corroborate those of other authors who stipulate that the levels of phenolic compounds and their activities may depend on several intrinsic and extrinsic factors. These factors generally include the nature of the organ, the harvesting period, the extraction technique used, the solvent used, the geographical location and the method used to assess antioxidant activity (Hanan and al 2022, and Souad and al).

However, the antioxidant activity of extracts from this plant could be attributed to total phenolic compounds and in particular total flavonoids (Djeridane and al, 2006). The presence of phenolic compounds in the leaves of this plant and its interesting antioxidant activity (low IC_{50}) could justify its widespread use by livestock farmers in northern Côte d'Ivoire, particularly those rearing broiler chickens.

Indeed, according to Gerasopoulos and al. (2015), dietary supplementation with phenolic compounds exhibits effective antioxidant activity in broilers. Moreover, it has been reported that vitamin E and C supplementation helps chickens combat oxidative stress and boost their immunity (Minand and al., 2018). For example, *Equol*, which is also an antioxidant compound obtained from the isoflavonoid *daidzein*, a soy isoflavone, can reduce the oxidative load induced by ROS (Liu and al., 2006). *Equol* also protects the intestinal epithelium against oxidative stress by enhancing the expression of antioxidant genes, stimulating the function of antioxidant enzymes and improving antioxidant capacity (Lin and al., 2016). Also, the inclusion of phenolic compounds in poultry farming has been shown to be effective in combating oxidative stress from rearing, processing to consumption of poultry meat. Plants contain powerful antioxidants, (Bidie and al. 2011, Ercetin et al. 2012) so their use in diets as additives could improve digestion in broilers. They are natural sources of antioxidants, and

Comment [N13]: Can you explain to us where and how these results came out?

Formatted: English (United States)

Comment [N14]: it seems to me that only the leaves of the plant are used and that the sampling was done in a specific area

Formatted: English (United States)

Formatted: English (United States), Highlight

Formatted: English (United States)

studies have shown that these plants are much more potent than synthetic antioxidants (Ali and Li, 2021, Bougandoura and al 2012, Bramen andal 1975). In addition, the antimicrobial activities of phenolic compounds have already been demonstrated by several authors. Indeed, the mechanisms of action of certain phenolic compounds involve inhibiting cell wall synthesis, cell membrane function and protein synthesis (Bhuiyan andal.2009 ; Pasril and Yuliasanti 2014). Recently, one of our studies demonstrated the antisalmonial activity of this plant on multi-resistant strains of *Salmonella typhimurium* isolated from the droppings of farmed broilers (Ouattara and al., 2023). The massive use of *Anogeissus leiocarpus* by farmers in the north of Côte d'Ivoire, particularly those rearing broilers, could be justified by its phenolic compound content and antioxidant activities.

CONCLUSION

This study confirms that *Anogeissus leiocarpus* is a plant whose leaves contain variable levels of phenolic compounds. The presence of phenolic compounds in the leaves of this plant, together with its interesting antioxidant activity (IC₅₀ between 4.4 mg/mL and 6.0 mg/mL), could justify its widespread use by livestock farmers in northern Côte d'Ivoire, particularly those rearing broiler chickens. In view of the results obtained, *Anogeissus leiocarpus* could be a potential candidate for the development of improved traditional medicines for the treatment of infectious diseases and oxidative stress in poultry. It would therefore be useful to continue research on this plant with a view to gaining a better understanding of its anti-infectious and antioxidant properties in poultry farming. In the future, it would be interesting for us to study the *in vivo* antioxidant activities of this plant in chicks in order to measure the biomarkers of oxidative stress (catalase, peroxidase, malondialdehyde and nitric oxide).

Comment [N15]: the discussion should highlight the importance of phenolic compounds, and the antioxidant tested and their effects, on the pathogens responsible for digestive diseases. In addition it would be interesting to show if there is a conventional or regulatory dosage.

Formatted: English (United States)

Formatted: English (United States), Highlight

Comment [N16]: the conclusion must highlight the new knowledge and useful information about the content of phenolic compounds in some selected traditional plant leaves and the antioxidant activity of their extracts

Formatted: English (United States)

Formatted: Font: 11 pt, English (United States)

Formatted: English (United States)

REFERENCES

Adama K. (2009). Anthelmintic activity of two tropical plants tested in vitro and in vivo on gastrointestinal strongyles of Mossi sheep from Burkina Fasso. Thesis.

Ali Hassan Nawaz and Li Zhang. (2021). Oxidative stress in broiler chicken and its consequences on meat quality. *International Journal of Life Science Research Archive*, 01(01), 045–054.

Arbab AH (2014). Review on Anogeissus leiocarpus a potent traditional drug. *International Journal of Research in Pharmacy and Chemistry* 4 (3) 465-500.

Barku V Y A, Yaw OB, Ansah EO, Dayie NTKD. (2013) In vitro assessment of antioxidant and antimicrobial activities of methanol extracts of six wound healing medicinal plants. *Journal of Natural Sciences Research*. Vol.3

Belemnaba L (2007). Antihypertensive properties of medicinal plants from Burkina Faso. Comparative study of three traditional medicine plants. DEA dissertation in Pharmacopoeia University of Ouagadougou 156p.

Bhuiyan M.N.I., Chowdhury J.U., Begum J. (2009). Chemical investigation of the leaf and rhizome essential oils of *Zingiber zerumbet* (L.) Smith from Bangladesh. *Bangladesh J Pharmacol* 4:9-12.

Bidie, A. P., N'guessan, B. B., Yapo, A. F., N'Guessan, J. D., & Djaman, A. J. (2011). Antioxidant activities of ten medicinal plants from the Ivorian pharmacopoeia. *Science & Nature*, 8(1-2), 1-12.

Bougandoura N., Bendimerad N. (2012). Evaluation of the antioxidant activity of aqueous and methanolic extracts of *Satureja calamintha* ssp. *Nepeta* (L) Briq Rev "Nature & Technology" B-Agronomic and Biological Sciences, 150p2012.

Bramen, A L, (1975). Toxicology and biochemistry of butylated hydroxyanisole and butylated hydroxytoluene. *J. Am Oil Chem. Soc.* 52, 59-63 PMID: 805808.

Choong T.S.Y., Chuah T.G., Robiah Y., Koay F.L.G. & Azni I. (2007). Arsenic toxicity, health hazards and removal techniques from water: an overview. *Desalination*, 217(1-3): 139-166.

Datta S, Sinha BK, Bhattacharjee S, Seal T. (2019). Nutritional composition, mineral content, antioxidant activity and quantitative estimation of water soluble vitamins and phenolics by RP-HPLC in some less used wild edible plants. *Heliyon*. 28;5(3): e01431. doi: 10.1016/j.heliyon.2019.e01431. PMID: 30976701; PMCID: PMC6441826.

Djeridane A, Yousfi M, Nadjemi B, Boutassouna D, Stocker P, Vidal N. (2006). Antioxidant activity of some Algerian medicinal plant extracts containing phenolic compounds. *Food Chemistry*. 97, 454-460.

Ercetin T, Sezer Senol F, Orhan IE, Toker G. (2012). Comparative assessment of antioxidant and cholinesterase inhibitory properties of the marigold extracts from *Calendula arvensis* L. and *Calendula officinalis* L. *Ind Crops Prod* ; 36:203-8.

Formatted: English (United States)

Formatted: English (United States)

Geldof N and Engeseth NJ (2002) Antioxidant capacity of honeys from various floral sources based on the determination of oxygen radical absorbance capacity and inhibition of in vitro lipoprotein oxidation in human serum samples. *Journal of Agricultural and Food Chemistry*, 50, 3050-3055.

Formatted: English (United States)

Gerasopoulos K, Stagos D, Kokkas S, Petrotos K, Kantas D, Goulas P, Kouretas D. (2015) Feed supplemented with byproducts from olive oil mill wastewater processing increases antioxidant capacity in broiler chickens. *Food and Chemical Toxicology*.

Githiori JB, Athanasiadou S, Thamborg SM. (2006) Use of plants in novel approaches for control of gastrointestinal helminths in livestock with emphasis on small ruminants. *Veterinary Parasitology* 139 308-320.

Hanen F., Chokri H., Ichrak M.; Riadh K. (2022): Evaluation of different procedures for extraction of phenolic compounds from medicinal plant: *Verbena officina*. *Biology Today* 225(3-4) 133-142.

Formatted: English (United States)

Holasova M, Fiedlerova V, Smrcinova H, Orsak, Lachman J, Vavrienova S, Buckweat. (2002). The source of antioxidant activity in functional foods. *Food Research International*. 35, 207-211

Julienne K., Tony T. B. A. S., Pascal A. O., Claude G. H., Basile S. B. K., Fréjus T. A. Z., Ibikounle M., Severin B., Sylvie M. H-A., Patrick A. E. (2021). *Anogeissus leiocarpus* (DC.) Guill. & Perr. (Combretaceae), a medicinal plant traditionally used in small ruminant breeding in West and Central Africa: zootechnical performances, pharmacological activities and chemical compositions (bibliography synthesis), *International Journal of Biosciences*, 19(5): 10-26.

Koné D. (2009) Ethnobotanical survey of six Malian medicinal plants Extraction, identification, quantification of polyphenols: Study of their antioxidant activity. Thesis.

Koné K. H. C., Coulibaly K, Konan K. S. (2019). Plants for medicinal use in sheep farming in Sinématiali (Northern Ivory Coast), *Journal of Animal & Plant Sciences*, 41 (1): 6828-6839.

Kumaran AJ, Karunakaran R. (2005). In vitro antioxidant activities of methanol extracts of five *Phyllanthus* species from India. *LWT* 40, 344-352.

Formatted: English (United States)

Le K, Chiu F. & Ng K (2007) Identification and quantification of antioxidants in *Fructus lycii*. *Food vvv Chemistry*, 105, 353-363.

Liu H, Zhang C, Zeng W. (2006) Estrogenic and antioxidant effects of a phytoestrogen daidzein on ovarian germ cells in embryonic chickens. *Domestic Animal Endocrinology*. 31(3) : 258–268.

Lin X, Jiang S, Jiang Z, Zheng C, Gou Z. (2016) Effects of equol on H₂O₂-induced oxidative stress in primary chicken intestinal epithelial cells. *Poultry Science*.

Marc F., Davin A., Deglène-Benbrahim L., & Ferrand C., (2004). Méthodes d'évaluation du potentiel antioxidant dans les aliments. *Erudit, M/S : médecine sciences*. 20(4) (2004) 458-463.

Formatted: English (United States)

Marinova D., Ribavora F. & Antanassova M. (2005). Total phenolics in bulgarian fruits and vegetables. *Journal of the University of Chemical Technology and Metallurgy*, 40(3) : 255-260.

Mario AG., Marisabel V. Raoul D. Silvia F. Alicia C. (2014) : Effects of Aloe vera coating on postharvest quality of tomato. *Fruits* (69) 117-126.

Min YN, Niu Z, Sun TT, Wang ZP, Jiao PX, Zi BB, Chen PP, Tian DL, Liu FZ. 2018. Vitamin E and Vitamin C supplementation improves antioxidant status and immune function in oxidative-stressed breeder roosters by up-regulating expression of GSH-Px gene. *Poultry Science*.

Okpékon T, Yolou S (2004). Antiparasitic activities of medicinal plants used in Ivory Coast. *Journal of Ethnopharmacology* : 90 : 91-97.

OMS. (2002). Médecine Traditionnelle : Rapport du Secrétariat. Conseil Exécutif. Cent onzième session. Point 5.7 de l'ordre du jour provisoire. EB 111/9 du 12 Décembre

Ouattara, A.L., Abou, O., Karamoko, O., Julien, G.K., and Adama, C. (2023). Antibacterial activities of aqueous and hydroethanolic extracts of *Anogeissus leiocarpus* on the in vitro growth of two multiresistant strains of *Salmonella typhimurium* isolated from broiler chickens. *Mycrobiology Research Journal International*. 33 (7) 38-46.

Formatted: English (United States)

Ouédraogo S, Belemnaba L, Traoré A, Bucher B, Guissou IP. (2011). Etude de la toxicité et des propriétés pharmacologiques de l'extrait aqueux de *Anogeissus leiocarpus* (DC) GUIL.L et PERR (Combretaceae). *Pharmacopée et médecine traditionnelle*. 15.

Parejo I., Codina C., Petrakis C. & Kefalas P., (2000). Evaluation of scavenging activity assessed by Co(II)/EDTA-induced luminal chemiluminescence and DPPH (2,2-diphényl-1-picryl-hydrazyl) free radical assay. *J Pharmacol Toxicol Method*, 44 : 507-512.

Formatted: English (United States)

Pasril Y. and Yuliasanti A. (2014). Daya antibakteri ekstrak daun sirih merah (*Piper crocatum*) terhadap bakteri *Enterococcus faecalis* sebagai bahan medikamen saluran akar dengan metode dilusi. *Insisiva Dental J*. 3 (1) : 88-95.

Sanchez-Moreno C., (2002). Methods used to evaluate the free scavenging activity in foods and biological systems, *Food Sci. And Technol. Inter*. 8(3) (2002) 121-137

Souad A. Moad R. (2020) : Protection by some plant methanol extracts of cherry tomatoes (*Solanum lycopersicum* var. *cerasiforme*) from fongic infection by *Alternaria alternata*. *Biologie Aujourd'hui* 214 (1-2) 55-61.

Stefano Dall' Aqua, Rinaldo Cervellati, Maria Cecilia Loi, Gabriella Innocenti (2008) : Evaluation of in vitro antioxidant properties of some traditional Sardinian medicinal plants : Investigation of the high antioxidant capacity of *Rubus ulmifolius*. 106. 2.

Wood JE, Senthilmohana ST, Peskin AV. (2002). Antioxidant activity of procyanidin-containing plant extracts at different pHs. *Food Chemistry*, 77(2): 155-161.

Comment [N17]: Harmonize and review all references then use the editor's reference style