

# Outline of the pharmacological and phytochemical profile of the jungle jalebi (*Pithecellobium dulce*)

## ABSTRACT

*Pithecellobium dulce* has numerous utilizations which has flexible jobs in the conventional arrangement of medication. A few investigations are being directed concerning the viability of the entire plant or its parts for the treatment of various sicknesses and ailments. Triterpenoids, sterols, flavonoids, tannins, and phytochemicals like quercetin, kaempferol, and dulcitol were recognized from the different pieces of the plant. In this article, the different remedial properties of *P. dulce* have been shown. *P. dulce* is an evergreen sharp tree; each piece of the plant has huge healthful values. The different parts of plant have been shown to display different pharmacological activities like anti-diabetic, locomotor, immunizing agent venom, free radical searching, protease inhibitor, mitigating, anti-bacterial, anti-mycobacterial, abortifacient, spermicidal, anti-convulsant, anti-ulcer, anti-diarrheal, anti-contagious, anti-tubercular, anti-tumor and as well as anti-oxidative properties and safety study like acute and sub-acute toxicity study. Therefore, the current review is cognizant on behalf of pharmacological studies, as well as phytochemicals.

**KEYWORDS:** *Pithecellobium dulce*, phytochemistry, traditional uses, antivenom, abortion, spermicidal safety study.

## INTRODUCTION

The Hindi name of Jungle jalebi is *Pithecellobium dulce* Benth, a woody legume native (northwestern region) of Mexico but now widely distributed all over India, Southeast Asia

and can also be found in South Africa and Australia; small to medium-sized. Height is about 18m and is cultivated throughout India. Seeds of *P. dulce* species are traditionally used worldwide for food because they are highly protein in nature and also contain dietary fibres and unsaturated fatty acid contents(1).

**Synonyms:** *Pithecellobium littorale*, *Pithecellobium dulce* (Roxb.) Benth.

- ✓ **Bark (contain 37 %tannin)** is used for the treatment of astringent in dysentery, dermatitis and inflammation in the eye.
- ✓ **Leaves (contain quercetin, kaempferol, dulcitol and afezilin)** are reported for to have astringent, emollient, abortifacient and antidiabetic properties.
- ✓ **Seeds** are used as glycosides, glycolipids, polysaccharides, steroids, saponins, lipids and phospholipids(2–4).
- ✓ **Roots** take over estrogenic activity.

Recently the studies of plant on alkylated resins from seed oil have been reported. Traditionally fruits have been consumed, well known for their edible fruits; linear, and curved; with lengths from 10 to 13 cm. Normally, the single pod contains ten seeds (irregular in shape).

The chemical compounds present in plants are used as antidiabetics, anticancer, antiinflammation etc. There are several allopathic medicines used in the treatment of diseases but they also have several side effects and drug interactions, so there is a need for herbal treatments that have lesser or no side effect. Here, the review discusses the protection against various diseases and their pharmacological properties of the *P. dulce*(5,6).



**Figure 1. *Pithecellobium dulce* (Jungle jalebi) Plant.**

**FAMILY**

The plant belongs to family Fabaceae. Most woody species are tropical; herbaceous (i.e., nonwoody) species happen principally in mild districts, belongs to family Orchidaceae and assteraceae as well. It contains about 751 genus and 19000 species also(7). Crotalaria, indigofera, astragalus, acacia, and mimosa are the widespread genus. About 19,000 known vegetable species add up to about 7% of blooming plant species. In tropical rainforests and in dry backwoods in the

Americas and Africa most widely recognized family is Fabaceae. Fabaceae is a single monophyletic family, Recent molecular and morphological evidence supports the fact. Here the conclusion has been given not just by the level of interrelation appeared by changed gatherings inside the family contrasted and that found among the Leguminosae. In addition by all the ongoing phylogenetic examinations dependent on DNA groupings(8,9). These investigations confirm that the family Fabaceae is a monophyletic bunch that is firmly identified with the families' polygalaceae, surianaceae and quillajaceae. A few products of the soil roots, various Leguminosae have human nourishment for centuries and their utilization is firmly identified with human advancement. Generic name is derived from the Greek words *pithekos* which means "ape" or "monkey," and *ellobion* stands for "earring," which means coiled shape. Genus is commonly known as blackbeads(10).

## **VERNACULAR NAME**

<b>Language</b>	<b>Name</b>
<b>Hindi:</b>	Jangle jalebi, Vilayati babul, Vilayati imli
<b>English:</b>	Quamachil, Madras thorn, manila tamarind
<b>Tamil:</b>	Kodukkaapuli
<b>Sanskrit:</b>	Kodukkaapuli
<b>German:</b>	Camambilarinde

## **Taxonomical Classification**

<b>Domain:</b>	<i>Eukaryote</i>
<b>Kingdom:</b>	<i>Plantae</i>
<b>Subkingdom:</b>	<i>Tracheobionta</i>

<b>Species:</b>	<i>Pithecellobium dulce</i> (Roxb.) Benth.
<b>Class:</b>	<i>Magnoliopsida</i>
<b>Subclass:</b>	<i>Rosidae</i>
<b>Order:</b>	<i>Fabales</i>
<b>Family:</b>	<i>Fabaceae</i>
<b>Genus:</b>	<i>Pithecellobium</i> Mart.
<b>Botanical name:</b>	<i>Pithecellobium dulce</i> (11).

### **BOTANICAL DESCRIPTION**

In tropical area the tree grows well at low and medium altitudes. The normal rainfall (low as 140mm and as high as 2,200mm) is tolerated. The plant growths need the average temperature (18 - 27.9°C) and also needs a well-drained soil. A pair of straight stout short thorns at the base of the leaf Pinnate leaves with a single pair pinnae, each with a single pair of leaflets often variegated White flowers in globular heads c. 1 cm wide Coiled pinkish pods Shiny black seeds c. 1 cm long, embedded in white pulp. But in favourable soil condition height reach of 10 metres (5 - 6 years). After planting in field other than occasional pruning the tree doesn't need any treatment.

*Pithecellobium dulce* pod's exterior resembles that of tamarind, but it easily peels away like the thin skin of a green bean. The edible flesh may then be separated from the seeds and eaten out-of-hand or prepared in sweet and savoury dishes. The Madras Thorn is a spiny tree that grows up to 20 meters tall. It has an irregular spreading nature making it ideal for a perimeter hedge or living fence. The curvy and spiralled pods are greenish-brown to red or pinkish colour. It is 10-15 cm long x 1-2 cm wide. They are thin and flattened with approximately 10 seeds per pod.<sup>9</sup>the pink, edible pulp of Madras Thorn is both sweet and savoury, with flavours of chestnut and honey. It is the consistency of sticky popcorn, and mildly sour with a cleansing astringent quality on the palate. Fresh Madras Thorn is highly perishable, and the pinkish-white pulp will quickly oxidize once peeled. At room temperature, the fruits keep for three to four days.

### **GEOGRAPHICAL DISTRIBUTION**

This plant originated from different countries is mentioned below:

- ✓ Brazil
- ✓ Argentina
- ✓ Bolivia


✓ Colombia

Jungle jalebi is the only specie that is widely spread outside from its origin. Eighteen species belongs to this genus; geographically distributed in many countries:

✓ India

✓ Huawei

✓ Tropical Africa(12).



**Figure 2.** Worldwide Distribution of *Pithecellobium dulce*. (Jungle jalebi).

### TRADITIONAL USES

- Astringent, homeostasis (bark, pulp of fruit)
- Antipyretic
- Eye inflammation (eye infections and swelling of the eyelids)
- To treat gum ailments,
- Toothache
- Anticancer
- Bile (leaves, also prevent abortions)
- Antiulcer(Seed)

- Haemorrhages
- Chronic diarrhoea
- Tuberculosis.

## PHYTOCHEMISTRY

**Table 1. Plant parts used for the study and their chemical name**

S. No.	Part used	Chemical Name
1.	Bark	Catechol
2.	Wood	Campesterol
3.	Wood	Melacacidin
4.	Seeds	Hederagenin
5.	Seeds	Pitheduloside A
6.	Seeds	Pitheduloside B
7.	Seeds	Pitheduloside D
8.	Seeds	Pitheduloside F
9.	Seeds	Pitheduloside H
10.	Seeds	Pitheduloside I
11.	Seeds	Pitheduloside K
12.	Leaves	Octacosanol
13.	Leaves	Kaempferol
14.	Seeds	Pithogenin
15.	Fruits	Ellagic acid
16.	Fruits	Gallic acid
17.	Fruits	Mandelic acid
18.	Fruits	Ferulic acid
19.	Fruits	Vanillic acid
20.	Fruits	Coumaric acid
21.	Fruits	Rutin
22.	Fruits	Naringin
23.	Peel, leaves	Quercetin
24.	Peel, seeds	Stigmasterol
25.	Fruits, peel	Pinitol
26.	Seeds	Oleanolic acid
27.	Seeds	Echinocystic acid
28.	Leaves	Dulcitol
29.	Stem	Prenylapigenine
30.	Fruits	Daidzein

## PHARMACOLOGY

#### ◆ **Abortion Activity**

The isoflavonoid isolated from root remove, took a stab in female rodents, demonstrated segment subordinate estrogenic development and extending dode weight  $15.5 \pm 0.25$  mg in control group to  $34.2 \pm 0.68$  mg in orally treated rodents dosing (1.25 mg/kg/day for 4 days).

#### ◆ **Antivenom Activity**

Polyphenols from water concentrate of *P. dulce* was tried for their inhibitory exercises against *Naja kaouthia* (NK) venom by In-vitro balance technique.

#### ◆ **Spermicidal Activity**

The saponins of *P. dulce* were besides introduced to tests for the spermicidal property by exhibited to activity in the debilitating against human semen.

#### ◆ **Antiinflammatory Activity**

The saponin (containing two geninacids, echinocystic corrosive and oleanolic destructive with xylose, arabinose and glucose as sugar moieties) procured from results of *P. dulce* has been concentrated against the exudative and proliferative time of provocative reaction in pale cleaned individual rodents by using carrageenan induced oedema and formaldehyde started joint pain models.

#### ◆ **Antitubercular Activity**

leaves extraction (by hexane, chloroform and alcoholic) was read for their anti-mycobacterial action by BACTEC460TB-Radiospirometric framework. The concentrates of the leaves were perused for their anti-mycobacterial activity by BACTEC460TB-Radiospirometric structure. Alcoholic aggregate 20mg/ml the conc. exhibited most raised activity when essentially indistinguishable with standard meds for ex. rifampicin, streptomycin, ethambutol, pyrazinamide andisoniazid(13,14).

#### ◆ **Activity of Protease Inhibitor**

A seed of *P. dulce* is isolation and characterizations of a protease inhibitor are seemed.

#### ◆ **Antimicrobial Activity**

The methanolic and aq. concentrate of *P. dulce* seeds have indicated fungistatic and have fungicidal impacts against plant pathogens like *Fusarium oxysporum*, *Botrytis cinerea*, *Penicillium digitatum*, and *rhizopusstolonifer*. A couple triterpene saponins, pitheduloside A, B, E, F and I frustrated in vitro mycelial progression of *Rhizopus stolonifer* and *Colletotrichum gloeosporioides*. The plant concentrate of the less-polar hexane and polar methanol attempted against different microorganisms and parasites were in the like way observed as incredible(15,16).

#### ◆ **Anti-Ulcer genic Activity**

Anti-ulcer activity is the ulcer control rodents. The activities of H (+), K (+) - ATPase and myeloperoxidase, seen as essentially raised up and saw as lessened in cure pre-treated rodents. Acute gastric ulcer controlling liquor, acetylsalicylic destructive, or hypothermic impediment stress to rodents pre-treated with HAEPD (200 mg/kg body weight for multi day). The effects of *P. dulce* difference standard prescription with omeprazole.

#### ◆ **Activity of Locomotor**

Locomotor activity is called CNS depressant activity. It is watery and alcoholic of leaf of *P. dulce* was evaluated using actophotometer in pale skinned person rodents. In light of addition in the centralization of GABA as a primary concern it shows a CNS depressant movement. The power of alcoholic and watery concentrates of leaf contrasted and chlorpromazine(17,18).

### **SAFETY STUDY**

#### ☐ **Acute toxicity study**

The rodents were fasted for whole with water at free access. For the intense danger, they were randomly partitioned in two gatherings of males, each successively dosed at interims of 48 hour. The controlled group got orally refined water; experimental group was orally treated with stem bark hydroethanolic concentrate of *P. dulce* with extraordinary dose of 5 g/kg. Animals were

watched for general behavioral and body weight changes, hazardous manifestations and mortality inside a time of four hr. for guaranteed indications of harmfulness and in any event once every day for 14 days for postponed indications of poisonous quality. After sedation by inward breath with ether, the spinal-rope of rodents was dis-joined by extending(19).

#### ❑ **Sub-acute toxicity study**

Repeated dose of orally toxicity study was done by OECD rule 407 (OECD, 2008) with slight adjustment without EDTA were centrifuged. The animals were partitioned into three groups of 6 animals each (3 male and 3 female). First group got 10 mL/kg body weight of distilled water and kept as control. Second and third group got extracted portions of 0.5 and 1 g/kg body then weight individually. For twenty eight days the extract were given and at a similar time and saw in any event twice day by day for bleakness and mortality, the animals(B.W.)were evaluated daily. On day 29, after an overnight fast the rodents were anesthetized with ether and blood tests were acquired by retro-orbital cut, utilizing fine cylinders for hematological and biochemical investigations with and without EDTA(20).

## **DISCUSSION AND CONCLUSION**

In the present review we have made an attempt to congregate the taxonomical, botanical and Pharmacological information on *Pithecellobium dulce*, a medicinal plant used in the Ayurveda. *Pithecellobium dulce* has a solid intense in wellbeing advancing, illness forestalling and life delaying properties which has been depicted, explored and checked by current specialists. In any case, the inherently dynamic mixes and the substance dependable have been resolved at this point, and a few components of the activity of *Pithecellobium dulce* are as yet obscure. Along these lines, bioassay-guided separation and distinguishing proof of the bioactive parts must be created to uncover the structure-movement relationship of these dynamic segments. Anyway more examinations are required to investigate the utilization of the plant for commercialization of dynamic elements of organic and home grown medication applications.

## **ETHICAL STATEMENT**

A pharmacist works to advance each patient's well-being in a discreet, kind, and caring way.

## **INFORMED CONSENT**

Using websites, review articles, and other sources to produce research content.

## REFERENCES

1. Narayan JP. Exploratory Studies on Occurrence and Potential Benefits of High Fruit Yielding Reproductive Phenophasic Variants of *Pithecellobium dulce* in Augmenting Fruit Based Semi-arid Agro Forestry Systems. *European J Med Plants*. 2019;28(2):1–17.
2. Jimenez-chino AL, Bugarin A, Zamilpa A, Gives PM de, Villa-mancera A, Olivares-p J, et al. Phenolic Acids and Flavonoids from *Pithecellobium dulce*. *Plants*. 2022;1–13.
3. Aldarhami A, Bazaid AS, Alhamed AS, Alghaith AF, Ahamad SR, Alassmrry YA, et al. Antimicrobial Potential of *Pithecellobium dulce* Seed Extract against Pathogenic Bacteria: In Silico and In Vitro Evaluation. *Biomed Res Int*. 2023;2023.
4. Ta S Ranganamy SK, Shubashini S, Sripathi SK, Ravindran NT, Balashanmugam P. Evaluation of In Vitro Antioxidant and Antibacterial Activity of *Pithecellobium dulce* Benth Fruit Peel. *Artic Int J Curr Res [Internet]*. 2011;(May 2014):378–82. Available from: <https://www.researchgate.net/publication/230688534>
5. Kamat SK, Barde PJ, Raut SB. Evaluation of the estrogenic activity of Indian medicinal plants in immature rats. *Anc Sci Life*. 2015;35(2):90–5.
6. Janeczko A. Estrogens and androgens in plants: The last 20 years of studies. *Plants*. 2021;10(12).
7. Ketcha Wanda GJM, Zemo Gamo F, Njamen D. Medicinal plants of the family of fabaceae used to treat various ailments. *Fabaceae Classif Nutr Compos Heal Benefits*. 2015;(May 2022):1–19.
8. Arigela RK, Ch SR. New synonymy in the genus *Crotalaria* L. (Fabaceae). 2022;(January).
9. Badry MO, Al-Hawshabi OSS, Osman AK. Flora and phytochorology of lahiq governorate of yemen: 1-systematic revision of wild legumes of the family fabaceae. *Egypt J Bot*. 2021;61(2):591–610.
10. Bello MA, Hawkins JA, Rudall PJ. Floral morphology and development in quillajaceae and surianaceae (Fabales), the species-poor relatives of leguminosae and polygalaceae (*Annals of Botany* (2007) 100, (1491-1505)). *Ann Bot*. 2008;101(3):483.
11. Datiles M, Acevedo-rodriguez P. *Pithecellobium dulce* (Manila tamarind). *PlantwisePlus Knowl Bank*. 2022;Species Pages(November).
12. Rajan N, Shukla P. Traditional Utilization and Pharmacological Properties of Medicinal Plants. 2023.
13. Murakami G. Distinct effects of Estrogen on mouse maternal behavior: The contribution of Estrogen synthesis in the brain. *PLoS One*. 2016;11(3):1–16.
14. Velle W. [Induction of abortion by oestrogens in animals. A review (author's transl)]. *Nord Vet Med*. 1974 Oct;26(10):563–71.
15. Kim JY, Park SC, Hwang I, Cheong H, Nah JW, Hahm KS, et al. Protease inhibitors from plants with antimicrobial activity. *Int J Mol Sci*. 2009;10(6):2860–72.
16. Sawasdipuksa N, Lei Z, Sumner LW, Niyomploy P, Sangvanich P. A lysozyme with antifungal activity from *Pithecellobium dulce* seeds. *Food Technol Biotechnol*. 2011;49(4):489–94.
17. Sharifi-Rad M, Fokou PVT, Sharopov F, Martorell M, Ademiluyi AO, Rajkovic J, et al. Antiulcer agents: From plant extracts to phytochemicals in healing promotion. Vol. 23, *Molecules*. 2018. 1–37 p.

18. Brito SA, de Almeida CLF, de Santana TI, da Silva Oliveira AR, do Nascimento Figueiredo JCB, Souza IT, et al. Antiulcer activity and potential mechanism of action of the leaves of *Spondias mombin* L. *Oxid Med Cell Longev*. 2018;2018.
19. Nowland MH, Hugunin KMS, Rogers KL. Effects of short-term fasting in male Sprague-Dawley rats. *Comp Med*. 2011;61(2):138–44.
20. Toyoda K, Shibutani M, Tamura T, Koujitani T, Uneyama C, Hirose M. Repeated dose (28 days) oral toxicity study of flutamide in rats, based on the draft protocol for the “Enhanced OECD Test Guideline 407” for screening for endocrine-disrupting chemicals. *Arch Toxicol*. 2000 May;74(3):127–32.

UNDER PEER REVIEW