

Management of Root Knot nematode *Meloidogyne incognita* on carrot

ABSTRACT

Experiments were conducted to find out the efficacy of certain botanicals like *Azadirachta indica*, *Azadirachta indica*, *Tagetes erecta*, *Ricinus Communis*, *Lantana camara* and *Parthenium hysterophorus* and different biocontrol agents like *Glomus fasciculatum*, consortia (*T. harzianum*, *T. viride* and *Pseudomonas fluorescens*) and vermicompost for the management of *M. incognita* on carrot. The treatment with *A. indica* @ 1.5% (w/w) among the botanicals and the treatment with *G. fasciculatum* @ 300 spores/m² + consortia @ 5ml/l + vermicompost @ 1.25t/ha was found most effective in increasing plant growth parameters and in decreasing the *M. incognita* infection on carrot.

Key words: Botanicals, *Azadirachta indica*, *Azadirachta indica*, *Tagetes erecta*, *Ricinus Communis*, *Lantana camara*, *Parthenium hysterophorus*, biocontrol agents, *Glomus fasciculatum*, *T. harzianum*, *T. viride*, *Pseudomonas fluorescens*, vermicompost, *M. incognita*, carrot.

INTRODUCTION

Carrot (*Daucus carota*) is an important and common vegetable crop grown throughout the world. Out of different plant parasitic nematodes, root knot nematodes (*Meloidogyne* spp) causes considerable damage to the crop with a yield loss of 59.3 percent (Sreenivasan, 2017) due to attack of *M. spp*. Disease management through eco-friendly biological approaches is becoming inevitable which is safe for both user and treated crop, compared to synthetic chemicals.

Comment [Maryam1]: The Abstract is scanty and you should state the most effective, second and least effective treatment.

Comment [Maryam2]: Try to capture the family, origin climatic requirement as well as the importance of carrot before introducing the pest.

Write the statement of the problem and objective,

MATERIALS AND METHODS

Apot experiment was conducted in the net house of Department of Nematology, AAU, Jorhat to study the efficacy of certain botanicals against *M. incognita* on carrot. Earthen pots of 1kg capacity were filled up with sterilized soil and inoculated with J_2 of *M. incognita* @ 1000 J_2 per pot. The soil was incorporated with chopped leaves of *Azadirachta indica*, *Azeratum conizoides*, *Tagetes erecta*, *Ricinus Communis*, *Lantana camara*, *Parthenium hysterophorus* each @ 1.0 and 1.5% (w/w). and the control without application of botanicals. After ten days, three seeds of carrot variety, 'Early nantes' was sown in each pot. Each treatment were replicated five times and arranged in a completely randomized design. The pots were watered regularly for proper decomposition. The treatments taken in the investigations were: T₁: *A. indica* @1.0% (w/w), T₂: *A. indica*@1.5% (w/w), T₃: *A. conizoides* @1.0% (w/w), T₄: *A. conizoides* @1.5% (w/w), T₅: *T. erecta* @1.0% (w/w), T₆: *T. erecta* @1.5% (w/w), T₇: *R. communis* @1.0% (w/w), T₈: *R. communis* @1.5% (w/w), T₉: *L. camara*@1.0% (w/w), T₁₀: *L. camara*@1.5% (w/w), T₁₁: *P. hysterophorus*@1.0% (w/w), T₁₂: *P. hysterophorus* @1.5% (w/w), T₁₃: Control. After ninety days of sowing, plants were uprooted carefully and roots were washed free of soil. Observations on different plant growth parameters viz., length of shoot and tap root, fresh and dry shoot and tap root weight, number of galls and egg masses and final nematode population in soil were recorded.

The microplot experiment was conducted during the *rabi* season of 2015-16 to study the efficacy of *Glomus fasciculatum*, consortia (*T. harzianum*, *T. viride* and *Pseudomonas fluorescences*) and vermicompost for the management of *M. incognita* on carrot. For this microplots infested with *M. incognita* (1-2 J_2 per gram soil) of 1x1 m² size were used and recommended dose of manure and fertilizers were

Comment [Maryam3]: There should be a subsection indicating the coordinate of the experiment

Write all scientific in full for the first time

Comment [Maryam4]: Write in full

Comment [Maryam5]: Write in full for the first time

applied. Initial population of *M. incognita* in different microplots of size 1x1 m² size were recorded. Vermicompost was applied 10 days before sowing. Carrot seeds were sown in rows at the spacing of 30 cm between rows and 10 cm from plant to plant. Consortia and *G. fasciculatum* were applied at the time of sowing. The experiment was laid out in a Randomized block design with following treatments and each treatment was replicated three times. The different treatments were: T₁: Soil application of consortia @ 5ml/ltr/m², T₂: Vermicompost @ 2.5t/ha, T₃: *G.fasiculatum*@ 600 spores/m², T₄: Soil application of consortia @ 5ml/ltr+ vermicompost @ 1.25t/ha, T₅: Soil application of consortia @ 5ml/ltr + *G.fasiculatum*@ 300 spores/m², T₆:Vermicompost @ 1.25t/ha +*G. fasiculatum*@ 300 spores/m², T₇ : Soil application of consortia @ 5ml/ltr+ vermicompost @ 1.25t/ha +*G. fasiculatum*@300spores/m², T₈ : Control. Observations were taken at full maturity. For this 10 plants from each plot were uprooted randomly and roots were washed carefully in tap water to remove adhering soil particles. Records on length of shoot and tap root, fresh and dry weight of shoot and tap root, number of galls per root system, number of healthy and forked tap root per plot, yield per plot (q/ha) and final nematode population in soil were recorded.

Comment [Maryam6]: How many days exactly

Comment [Maryam7]: How many grams of roots were use?

Comment [Maryam8]: What quantity of soil was used?

RESULT AND DISCUSSION

The results of the pot experiment exhibited maximum increase in plant growth in the treatment with *A. indica* @ 1.5% (w/w) (Table 1) . All the treatments were found to be effective in increasing growth parameters of carrot over control. These results are in agreement with John and Hebsy (2000) who reported that bare root dip of brinjal seedlings in neem leaf extract significantly increased the plant growth parameters. This may be due to the presence of azadirachtin, a major nematotoxic compound in neem which

Comment [Maryam9]: There are more current literatures, this is too old

is released through volatilization, exudation, leaching and de-composing of the plant parts that suppress the population of plant parasitic nematode and helps in growth of plant (Akhtar, 2000).

Comment [Maryam10]: Too old

Maximum reduction in number of galls, egg masses and final nematode population in soil were also recorded in the treatment with *A. indica* @ 1.5% (w/w) (Table 2). This may be due to the presence of terthienyl and bithienyl compounds, the toxic principle of marigold, (Uhlenbroek and Bijloo, 1958). Similar findings were also observed by Nelaballe and Mukkara (2013) who reported that leaf extracts of *Daturastramonium*, *Azadiractaindica*, *Calotropisprocera* and *Crotalaria juncea* reduced the infestation of *M. incognita* in mulberry plant among which the highest juvenile mortality (82.8%) was recorded with neem extract. This may be due to the presence of active toxic principles like azadirachtin and nimbin in *A. indica* and thiophene and alpha-terthienyl in *T. erecta*, suppress the nematode population as well as improved plant growth. (Rao and Parmer, 1984).

Comment [Maryam11]: Too old

The result of the microplot experiment revealed that maximum increase in plant growth parameters were recorded in the treatment with *G. fasciculatum* @ 300 spores/m² + consortia @ 5 ml/l + vermicompost @ 1.25 t/ha (Table 3). Nama *et al.* (2015) also reported that talc based formulation of *T. viride*, *T. harzianum*, *P. fluorescens* against *M. incognita*, significantly increased the plant growth characters and reduced nematode reproduction as compared to untreated checks on cowpea. *Trichoderma* sp. is highly rhizosphere competent i.e., able to colonize on roots as it develops, thus promote plant growth. It may also exert several mechanisms such as tolerance to stress through enhanced root and plant development, induced resistance, inactivation of pathogen's enzymes in improving plant growth and suppressing plant pathogens (Weeder *et al.*; 2008).

Maximum decrease in number of galls and final nematode population were also recorded in the treatment with *G. fasciculatum* @ 300 spores/m² + consortia @ 5ml/l + vermicompost @ 1.25t/ha. All the treatments were found to be effective in reducing the number of galls, egg masses and final nematode population over control (Table 4). Similar findings were also recorded by Masadeh *et al.* (2004) who reported that combination of the arbuscular mycorrhizal fungus *G. intraradices* and the fungus *T. virider* retarded the development of the giant cell system, growth of the nematode, consequently reduced production of egg masses and nematode population. The reduction in number of galls, egg masses and also the nematode population in soil in mycorrhizal treatments may be due to increased resistance by mycorrhizal plants to nematodes or due to colonization of the roots by *G. fasciculatum* which mechanically prevents the entry of *M. incognita* into the roots of tomato (Mittal *et al.*, 1991). Reduction of nematode due to *T. harzianum* may be attributed to the direct parasitism of eggs and larvae through the increase in chitinase and protease activity, which could be indicator of eggs and infection capabilities. Sharon *et al.* (2001) reported two mechanisms involved in *Trichoderma* antagonism against root-knot nematode (i) *Trichoderma* produced metabolites with antinematode activity that immobilized J₂ thus reduced root penetration and (ii) direct parasitism by *Trichoderma*.

Comment [Maryam12]: How did you rate the effect on the gall

REFERENCES

- Akhtar, M. (2000) Nematicidal potential of the neem tree *Azadirachta indica* (A. Juss). *Integrated Pest Management Reviews*. **5**: 57-66.
- John, A. and Hebsy, B. (2000). Bare-root dip treatment of brinjal seedlings in phytochemicals for the management of root-knot nematodes (*Meloidogyne incognita*). *J. Trop. Agric.* **38**: 69-72.

Comment [Maryam13]: Write in full. Do so were applicable

- Masadeh, B.; Von Alten, H.; Stoecker, G. G. and Sikora, R. A. (2004). Biocontrol of root-knot nematodes using the Arbuscular mycorrhizal fungus *Glomus intraradices* and the antagonist *Trichoderma viride* in two tomato cultivars differing in their suitability as host for the nematodes. *J. Plant Dis. Protec.* **111**(4): 322-333.
- Mittal, N; Sharma, M.; Saxena, G and Mukherjee, K.G. (1991). Effect of VA mycorrhiza on gall formation in tomato roots. *Pl. Cell Incompatibility News*. **23**:39-43
- Nama, C.P.; Sharma, H. K. and Siddiqui, A. U. (2015). Efficacy of bio-agents against root-knot nematode, *Meloidogyne incognita* infecting cowpea (*Vigna unguiculata* L.) *Journal of biopesticides*, **8**(1): 19-22 .
- Nelaballe, V.K and Mukkara, L.D (2013). Study on the nematicidal Effect Of some local flora on *Meloidogyne Incognita* Chit wood Infesting mulberry. *Intn. J. Chemical, Envnt. and Biol. Sci.* **1**: 2320-4087.
- Rao, K. N. and Parmer, B. S. (1984). A compendium of chemical constituents of neem, neem. *news lett*, **1**: 39-46
- Sharon, E.; Bar-Eyal, M.; Chet, I.; Herrera-Estrella, A.; Kleifeld, D. and Spiegler, Y. (2001). Biological control of the root-knot nematode *Meloidogyne javanica* by *Trichoderma harzianum*. *Phytopathol.* **91**(7): 687-693.
- Sreenivasan, N. (2017). Status of root knot nematode, *Meloidogyne hapla* infection of carrot at Kodaikanal hills of Tamil Nadu, India and its yield loss estimation. *Int. J. Curr. Microbiol. App. Sci.* (2017) **6**(9): 3629-3635.

Uhlenbroek, J.H. and Bijloo, J.D. (1958) Investigation on nematocides. Isolation and structure of a nematocidal principle occurring in Tagetes roots. *Recueil des Travaux Chimiques des Pays-Ba.* 77, 1004-1008

Weeder, C. R.; Shelton, A. M. and Hoffman, M.P. (2008). Biological control. In : *A guide to Natural Enemies in North America*, pp. 1-14.

UNDER PEER REVIEW

Table 1. Effect of botanicals on the plant growth of carrot. (Mean of 5 replications)

Treatments	Shoot length (cm)	Tap root length (cm)	Fresh weight of shoot (g)	Dry weight of shoot (g)	Fresh weight of tap root (g)	Dry weight of tap root (g)
-------------------	--------------------------	-----------------------------	----------------------------------	--------------------------------	-------------------------------------	-----------------------------------

T ₁ : <i>Azadirachta indica</i> @ 1.0% (w/w)	38.76 ^c	18.72 ^c	9.12 ^a	5.54 ^c	28.08 ^b	17.76 ^c
T ₂ : <i>A. indica</i> @ 1.5% (w/w)	43.10 ^a	19.48 ^a	9.22 ^a	6.84 ^a	29.70 ^a	18.94 ^a
T ₃ : <i>Azeratumconizoides</i> @ 1.0% (w/w)	27.00 ⁱ	14.08 ^h	7.16 ^f	3.58 ^g	22.72 ^{fg}	12.90 ⁱ
T ₄ : <i>A. conizoides</i> @ 1.5% (w/w)	28.24 ^h	15.00 ^g	7.48 ^e	4.14 ^f	23.12 ^{ef}	13.00 ⁱ
T ₅ : <i>Tagetes erecta</i> @ 1.0% (w/w)	36.16 ^d	17.58 ^d	8.94 ^a	5.04 ^d	26.12 ^c	17.20 ^d
T ₆ : <i>T. erecta</i> @ 1.5% (w/w)	40.26 ^b	18.00 ^b	9.14 ^a	6.14 ^b	28.08 ^b	18.20 ^b
T ₇ : <i>Ricinus communis</i> @ 1.0% (w/w)	33.70 ^e	16.26 ^f	8.36 ^{bc}	4.04 ^f	24.12 ^d	15.82 ^f
T ₈ : <i>R. communis</i> @1.5% (w/w)	34.08 ^e	17.00 ^e	8.50 ^b	4.46 ^e	26.08 ^c	16.76 ^e
T ₉ : <i>Lantana camara</i> @1.0% (w/w)	29.32 ^g	15.04 ^g	8.00 ^d	4.00 ^f	23.20 ^e	13.92 ^h
T ₁₀ : <i>L. camara</i> @ 1.5% (w/w)	31.62 ^f	16.02 ^f	8.12 ^{cd}	4.06 ^f	24.00 ^d	14.94 ^g
T ₁₁ : <i>Partheniumhysterophorus</i> @1.0% (w/w)	25.00 ^k	13.26 ⁱ	7.00 ^f	3.14 ^h	22.58 ^g	11.98 ^j
T ₁₂ : <i>P. hysterophorus</i> @1.5% (w/w)	26.06 ^j	13.96 ^h	7.10 ^f	3.24 ^h	22.92 ^{efg}	12.08 ^j
T ₁₃ : control	20.84 ^l	10.04 ^j	4.98 ^g	3.12 ^h	18.16 ^h	9.26 ^k
S.Ed (±)	0.29	0.17	0.15	0.11	0.20	0.20
CD _{0.05}	0.58	0.34	0.30	0.23	0.40	0.42

Table2 . Effectof botanicals on *Meloidogyne incognita* on carrot(Mean of 5 replications)

Treatments	No. of galls	No. of egg mass	Final nematode population	% decrease over control
T ₁ : <i>Azadirachta indica</i> @ 1.0% (w/w)	14.80 (3.84) ^k	11.00 (3.08) ^{ij}	121.60 (11.02) ⁱ	61.34
T ₂ : <i>A. indica</i> @ 1.5% (w/w)	14.20 (3.76) ^k	8.60 (3.01) ^j	99.00 (9.94) ^l	68.53
T ₃ : <i>Azeratumconizoides</i> @ 1.0% (w/w)	33.80 (5.81) ^d	25.80 (5.07) ^d	151.60 (12.31) ^d	51.81
T ₄ : <i>A. conizoides</i> @ 1.5% (w/w)	31.80 (5.63) ^e	23.60 (4.85) ^e	144.40 (12.01) ^e	54.10
T ₅ : <i>Tagetes erecta</i> @ 1.0% (w/w)	21.60 (4.64) ^j	9.60 (3.17) ⁱ	101.00 (10.04) ^k	59.94
T ₆ : <i>T. erecta</i> @ 1.5% (w/w)	14.40 (3.79) ^k	9.00 (3.08) ^{ij}	100.80 (10.03) ^k	66.87
T ₇ : <i>Ricinus communis</i> @ 1.0% (w/w)	27.80 (5.27) ^h	17.40 (4.17) ^g	131.20 (11.45) ^h	58.29
T ₈ : <i>R. communis</i> @ 1.5% (w/w)	23.80 (4.87) ⁱ	14.20 (3.76) ^h	110.80 (10.52) ^j	64.78
T ₉ : <i>Lantana camara</i> @ 1.0% (w/w)	30.40 (5.51) ^f	22.80 (4.77) ^e	138.40 (11.76) ^f	56.00
T ₁₀ : <i>L. camara</i> @ 1.5% (w/w)	29.20 (5.40) ^g	19.40 (4.40) ^f	134.20 (11.58) ^g	57.34
T ₁₁ : <i>Partheniumhysterophorus</i> @ 1.0% (w/w)	38.20 (6.18) ^b	30.80 (5.54) ^b	159.00 (12.60) ^b	49.45
T ₁₂ : <i>P. hysterophorus</i> @ 1.5% (w/w)	35.00 (5.91) ^c	27.80 (5.27) ^c	154.20 (12.41) ^c	50.98
T ₁₃ : control	47.40 (6.88) ^a	37.20 (6.09) ^a	314.60 (17.73) ^a	
S.Ed (±)	0.05	0.05	0.03	
CD0.05	0.10	0.11	0.06	


Table3. Effect of *Glomus fasciculatum*, consortia and vermicompost on plant growth and yield of carrot (Mean of 3 replications)

Treatments	Shoot length (cm)	Tap root length (cm)	Number of healthy tap root per plot	Forked tap root per plot (%)	Yield (q/ha)	% increased over control
T ₁ = Soil application of consortia @ 5ml/ltr	41.03 ^e	13.03 ^f	19.00 ^e	32.20 ^b	120.00 ^e	24.16
T ₂ = Vermicompost @ 2.5 ton/ha	41.23 ^e	14.03 ^e	19.66 ^{de}	28.86 ^{bc}	123.33 ^e	27.59
T ₃ = <i>G. fasciculatum</i> @ 300 spores/m ²	39.83 ^f	12.00 ^g	18.00 ^e	31.20 ^b	106.66 ^f	10.34
T ₄ = Soil application of consortia @ 5ml/ltr + vermicompost @ 1.25 t/ha	45.13 ^c	16.23 ^c	22.00 ^c	21.06 ^d	150.00 ^c	55.24
T ₅ = Soil application of consortia @ 5ml/ltr + <i>G. fasciculatum</i> @ 300 spores/m ²	43.03 ^d	15.00 ^d	21.00 ^{cd}	24.40 ^{cd}	136.66 ^d	41.44
T ₆ = <i>G. fasciculatum</i> @ 300 spores/m ² + vermicompost @ 1.25 t/ha	47.93 ^b	18.86 ^b	24.00 ^b	14.40 ^e	166.66 ^b	72.41
T ₇ = Soil application of consortia @ 5ml/ltr + <i>G. fasciculatum</i> @ 300 spores/m ² + vermicompost @ 1.25 t/ha	49.63 ^a	20.56 ^a	26.00 ^a	8.86 ^e	176.66 ^a	82.82
T ₈ = control.	37.83 ^g	10.96 ^h	14.00 ^f	44.43 ^a	96.66 ^g	
S.Ed(±)	0.14	0.21	0.80	2.74	3.72	
CD _{0.05}	0.31	0.47	1.72	5.89	7.99	

Table 4. Effect of *Glomus fasciculatum*, consortia and vermicompost on *Meloidogyne incognita* on carrot (Mean of 3 replications)

Treatments	No. of galls	Final nematode population in soil	% decrease over control	% increase/decreased over INP
T ₁ = Soil application of consortia @ 5ml/ltr	20.83 (4.56)c	183.33 (13.53)c	48.01	-29.91
T ₂ = Vermicompost @ 2.5 t/ha	21.04 (4.58)c	181.00 (13.45)c	48.67	-27.90
T ₃ = <i>G. fasciculatum</i> @ 300 spores/m ²	23.16 (4.81)b	186.66 (13.66)b	47.07	-15.99
T ₄ = Soil application of consortia @ 5ml/ltr + Vermicompost @ 1.25 t/ha	17.43 (4.17)d	171.66 (13.10)e	51.32	-32.26
T ₅ = Soil application of consortia @ 5ml/ltr + <i>G. fasciculatum</i> @ 300 spores/m ²	18.16 (4.26)d	176.66 (13.29)d	49.90	-40.74
T ₆ = <i>G. fasciculatum</i> @ 300 spores/m ² + vermicompost @ 1.25 t/ha	16.90 (4.11)d	169.00 (12.99)f	52.07	-45.67
T ₇ = Soil application of consortia @ 5ml/ltr + <i>G. fasciculatum</i> @ 300 spores/m ² + vermicompost @ 1.25 t/ha	13.38 (3.65)e	163.66 (12.79)g	53.59	-43.21
T ₈ = control.	32.38 (5.69)a	352.66 (18.77)a		+23.45
S.Ed±	0.08	0.04		
CD _{0.05}	0.19	0.09		

UNDER PEER REVIEW



UNDER PEER REVIEW

