

RETINITIS PIGMENTOSA IN AWKA, NIGERIA : A SURVEY STUDY

Abstract

BACKGROUND

Retinitis pigmentosa which causes irreversible blindness and affects all gender is usually inherited.

OBJECTIVES: To describe the incidence and pattern of retinitis pigmentosa at the Eye Unit of Chukwuemeka Odumegwu Ojukwu University Teaching Hospital Awka, Nigeria

MATERIALS AND METHODS: The case files of all the new patients seen at the Eye Unit of the Chukwuemeka Odumegwu Ojukwu University Teaching Hospital Awka between January 2014 to December 2021 were reviewed. Those with clinical diagnosis of Retinitis Pigmentosa were selected and information bordering on age, sex, occupation, disease duration. associated morbidity and fundus findings were extracted and analyzed using descriptive statistics.

RESULTS

Out of the 5461 new patients seen at the Eye Clinic within the study period, 29(0.5%) were diagnosed of Retinitis Pigmentosa. Of the 29 patients 18 (62.1%) were males and 11(37.9%) females, with male to female ratio of 1.6:1. The age range was 7 years to 69 years. Thirteen patients (44.8%) were between 16 years

and 44 years. Median age was 33years while the mean age was 34. 8+15. 2 years. Nine (31.0%) patients were students. The most common complaints of the patients were poor vision, 27(93.1%) and refractive error 8(27.68). Seven patients (24.18) were blind at presentation while retinal pigmentation was the most frequent fundus finding 29 (100%).

CONCLUSION

Retinitis Pigmentosa is a cause of visual impairment and blindness which reduces the quality of life of affected persons. Early diagnosis and visual rehabilitation should be encouraged.

Keywords: Retinitis Pigmentosa, Awka, Nigeria.

Introduction

Retinitis pigmentosa is a group of hereditary disorders characterized by progressive loss of photoreceptor and retinal pigment epithelial (RPE) function^{1,2}. It is a primary pigmentary retinal dystrophy predominantly affecting the rods and cones³. However, the rods are more affected than the cones^{1,2}. The retinal dystrophic lesions of the retinitis pigmentosa is known to cause visual disability in all age groups⁴ and retinitis pigmentosa is a common diagnosis among children in school for the visually impaired in Nigeria^{5,6}. The prevalence varies between 1 in 4000 and 1 in 5000 of the world population.^{1,3} It appears in the childhood and progresses slowly, often leading to blindness in advanced middle age³. More than one million people are affected globally².

Retinitis pigmentosa has been reported to be a common cause of blindness during working life in the industrialized nations of the world.⁷ In Denmark,⁸ retinitis pigmentosa has been noted to be a leading cause of blindness in people aged 20 –

64years accounting for 29.9%. Also in kuwait⁹ retinitis pigmentosa is the leading cause of visual incapacitation in people younger than 60 years of age. In Africa, a study reported retinitis pigmentosa as the most common cause of inherited blindness¹⁰. A Cameroonian study¹¹ reported the frequency of retinitis pigmentosa to be 0.16% with bilateral blindness rate of 30%. In Nigeria, retinitis pigmentosa accounted for 0.69% of all new patients seen at the out patients department in Ibadan¹² and 5.2% of all retinal diseases in Ile-Ife¹³ all in Western Nigeria.

In Onitsha, southeastern Nigeria, Nwosu¹⁴ reported that retinitis pigmentosa constitutes 11.1% of retinal diseases and that it is an important cause of blindness and low vision. Yet in another study, Nwosu et al¹⁵ documented the prevalence of retinitis pigmentosa as 0.6%. Retinitis pigmentosa is therefore of worldwide occurrence^{4-9,12-15} and no race is known to be exempted or at a higher risk³.

Retinitis pigmentosa is a genetic disorder of the eye that causes loss of vision¹⁶. It is one of the most common forms of inherited retinal degeneration². There are multiple genes that when mutated can cause retinitis pigmentosa phenotype¹⁷. Inheritance patterns of retinitis pigmentosa have been identified as autosomal dominant, autosomal recessive, x-linked and maternally (Mitochondrially) acquired and are dependent on specific retinitis pigmentosa gene mutations present in the parental generation¹⁸. Missense mutation of rhodopsin gene mutations present in the parental generation¹⁸. Missense mutation of rhodopsin gene (rhodopsin is a pigment that plays an essential part in the visual transduction cascade, that is vision in low light conditions) most frequently follow autosomal dominant inheritance pattern and accounts for approximately 25% of autosomal dominant form of retinitis pigmentosa^{2,19}. The autosomal dominant variant has the best prognosis, of all the inherited retinitis pigmentosa; it has a later onset and runs a milder cause than autosomal recessive. The x- linked recessive disease is least

common, has an earlier onset and runs the most severe course resulting in total blindness by the 3rd or 4th decade^{20, 21}. In rare occasions, a dominant form of the x-linked mutation will affect both males and females equally²². Several other studies have reported various codon mutations associated with retinitis pigmentosa.^{19,23,24,25,26} Mutation in more than 250 genes has also been linked to retinal dystrophies²⁰. There is also sporadic cases of retinitis pigmentosa. Retinitis pigmentosa can also be classified into typical, atypical and syndromic (those associated with rare systemic disorders) types, but the typical variant is the most common^{12,20,21,25,27,28}. Symptoms of retinitis pigmentosa include trouble with night vision, decreased peripheral vision and as peripheral vision worsens, the affected may experience tunnel or tubular vision¹⁶. The onset of symptoms is generally gradual and often in childhood^{16,29}. However, complete blindness is rare²⁹.

Materials and methods:

This is a retrospective hospital-based study carried out at the Chukwuemeka Odumegwu Ojukwu University Teaching Hospital Awka, Nigeria. The case files of all the new patients seen at the Eye Unit of Chukwuemeka Odumegwu Ojukwu University Teaching Hospital Awka between January 2014 and December 2021 were reviewed. Those with clinical diagnosis of retinitis pigmentosa were selected and information bordering on visual acuity, age, gender, occupation, disease duration, associated morbidity and fundus appearance were extracted. The data were analyzed using descriptive statistics and presented as frequency tables.

Results:

Of the 5461 new patients seen at the Eye clinic of the Chukwuemeka Odumegwu Ojukwu University Teaching Hospital Awka within the study period, 29(0.5%) were diagnosed of retinitis pigmentosa. Eighteen (62.1%) were males and 11(37.9%) females with male to female ratio of 1.6:1. The age range was 7 years to

69 years. Thirteen patients were between the 16-44 years age range. The age range 1 – 15 years and ≥ 65 years had four (13.8%) patients each.

Table 1: Age and Sex distribution of retinitis pigmentosa patients

Age group (years)	Male	Female	Total
1 – 15yrs	3(10.4%)	1(3.4%)	4(13.8%)
16 – 44yrs	9(31.0%)	4(13.8%)	13(44.8%)
45 – 64yrs	5(17.2%)	3(10.4%)	8(27.6%)
≥ 65	1(3.4%)	3(10.4%)	4(13.8%)
Total	18(62%)	11(38%)	29(100%)

Male: female ratio = 1.6:1

mean age = 34.8yrs

The median age was 33 years and the mean age was 34.8 ± 15.2 years. The visual acuity (VA) at the presentation using World Health Organization (WHO) category were as follows: normal vision (27.6%), visual impairment 11(38%) and blindness 10(34.4%). After refraction, some had their visual acuity improved.

Table 2: Visual acuity (VA) at presentation and after refraction

Who, Category	VA at presentation	VA after refraction
Normal vision 6/6 – 6/18	8(27.6%)	13(44.8%)
Visual impairment < 6/18 – 3/60	11(38%)	9 (31.1%)

Blindness, VA < 3/60	10(34.4%)	7 (24.1%)
Total	29(100%)	29(100%)

WHO: World Health Organization

VA: Visual Acuity

Table 3: Occupational and sex distribution of patients with retinitis pigmentosa

Occupation	Frequency	Male	Female
Students	9(31.0%)	6(20.7%)	3(10.3%)
Traders	6(20.7%)	4(13.8%)	2(6.9%)
Artisans	5(17.2%)	3(10.3%)	2(6.9%)
Workers (civil servants/company workers)	7(24.1%)	3(10.3%)	4(13.8%)
Others	2(6.9%)	2(6.9%)	-
Total	29(100%)	18(62%)	11(38.0%)

Table 3 showed occupational distribution of the patients with retinitis pigmentosa. Nine (31.0%) patients were pupils/students, 7(24.1%) were workers (civil servants/company workers), 6(20.7%) traders, while 2(6.9%) were unclassified. 5(17.2%) were artisans.

The complaints of the patients at presentation were many and varied and some had more than one complaint. The chief complaints were poor vision 27(93.1%), poor night vision 23(79.3%) headache 9(31.0%), eye ache 6(20.6%) and diminished vision which had the highest frequency with all the patients complaining of such.

Table 4: Major complaints of retinitis pigmentosa at presentation

	Complaints	Frequency
1	Diminished vision	27(93.1%)
2	Poor vision at night	23(79.3%)
3	Headache	9(31.0%)
4	Eye ache (Eye pain)	8(27.6%)
5	Photophobia	6(20.6%)
	Total	73

Some patients had multiple complaints

The associated co-morbidity found among the retinitis pigmentosa patients were refractive error 8(27.6%) of which 4(13.8%) were myopia, 2(6.9%) hypermetropia while myopic astigmatism and hypermetropic astigmatism were 1(3.5%) each. Three (10.5%) patients had bilateral immature cataract of which 2(7.0%) were male and 1(3.5%) female. One male (3.5%) had unioocular aphakia, which was as a result of couching done by some itinerant doctors who came to their village. Glaucoma recorded 5(17.2%) patients of which 3 (10.38) were males and 2(6.9%) females.a

Table 5: Co-morbidity with retinitis pigmentosa

Co –morbidity	Frequency
Refractive error	8(27.6%)

Myopia	6(20.7%)
Glaucoma	5(17.2%)
Cataract	3(10.5%)
Hypermetropia	2(6.9%)
Unocular Aphakia	1(3.5%)

The fundus findings of the patient with retinitis pigmentosa were perivascular retinal pigmentation 29(100%), retinal vascular attenuation 22(75.9%) and disc pallor 18(62.1%). Glaucomatous optic disc cupping was seen in 5(17.2%) of retinitis pigmentosa patients.

j

Table 6: Fundus finding of the Patients with retinitis pigmentosa

S/N	Fundus Findings	Frequency
1	Bone spicule retinal pigmentation	29 (100%)
2	Retinal vascular attenuation	22 (75.9%)
3	Disc Pallor	18 (62.1%)
4	Disc cupping	5 (17.2%)

Discussion

Retinitis Pigmentosa (RP) comprises a large group of inherited vision disorders that cause progressive degeneration of the retina, the light sensitive membrane that coats the inside of the eyes³⁰. Peripheral or side vision gradually declines and eventually is lost in most cases. However, central vision is usually preserved until late in these conditions³⁰. Retinitis Pigmentosa is generally not a very common disease³¹. The frequency of retinitis pigmentosa in the present study was 0.5% and

this is in consonance with earlier studies by Nwosu¹⁵ (0.6%) and Ashaye¹² (0.69%).

However, other studies^{32,33} carried out on retinitis pigmentosa in same Southern Nigeria as Nwosu¹⁵ and Ashaye¹² did not give report of frequency on their population of studies. Eballe et al¹¹ in Cameroon reported the frequency of 0.16% which is lower than that of the present study and other studies.^{12,15} The higher mean age in the Cameroonian study may have affected the frequency. More males (62.1%) than females (37.9%) were found to have retinitis pigmentosa in the present review and this is in line with other authors^{31,32,33}. The age range mostly affected in this review is 16 – 44years (31.0%) followed by 45 – 64 years (17.2%). Onakpoya et al³³ in a multicentre study in southwestern Nigeria had reported similar trend in age range affected by the retinitis pigmentosa³³. This supports the findings of earlier studies^{7,8,9} that retinitis pigmentosa is a frequent cause of blindness during active life³⁴. This may spell socio-economic doom for the affected, their family and the society in general as the blind years may be prolonged. The occupational distribution of the patients in this study showed that majority of the patients were in their career and economic pursuits and the attendant visual incapacitation arising from retinitis pigmentosa may likely affect their progression in their chosen fields. Parmegigian⁷ and Marmor³⁵ had earlier reported the economic impact of those affected by retinitis pigmentosa on their families and the society at large. The quality of life and livelihood of the affected individuals are also impacted negatively³⁶.

Visual impairment (38%) and blindness (34.4%) were elicited in the present study. The blindness rate in the present study is similar to that of Eballa et al (30%), in Cameroon¹¹. However, this differs from that of Onakpoya et al³³ who reported a

higher rate of 41.7% and the difference could be due to the fact that Onakpoya et al³³ did a multicentre study. Yet while Ukponwan et al³² in Benin Nigeria reported blindness rate of 50%, another author³⁷ in the United States of America (USA) had reported blindness rate of 25%. These differences could generally be attributed to age of presentation of the patients. Onakpoya et al³³ had earlier reported that older patients had higher rate of blindness at presentation. However, the youngest patient, 7 years old, had VA of <3/60 that was not improved by refraction and he dropped out of school. In the same vein, a 30 year old woman in this review also had profound visual loss and was relieved of her duty in her work place. This gives credence to the economic impact of retinitis pigmentosa on the lives of the affected and their families^{7,8,9}.

A mode of inheritance may also have accounted for the severity of the disease in these patients³. However, no positive family history was elicited in this study as opposed to other studies^{14,33}. For the fact that many of the diseased patients in this study fell into the visual acuity of economic and social blindness³⁸, it then becomes necessary to institute the policy of economic and social reorientation and rehabilitation of all the retinitis pigmentosa patients no matter the level of visual acuity at the time of first presentation. Generally, diminished vision (93.1%) was the major symptom that made patients to seek medical help. This finding was similar to the reports of 90% Ukponmwan et al³² and Eballe et al¹¹, 85% of them were working in Cameroon and Benin city respectively^{11,32}. However, the report by Onakpoya et al³³ in a multicentre study in southwestern Nigeria reported 69.8% which was much lower than the 93.1% recorded in the present survey. Edema et al⁶ and Abah et al³⁹ had earlier reported that retinitis Pigmentosa is a common diagnosis among children in schools for the visually impaired in Nigeria. This confirms the fact that retinitis pigmentosa and its co-morbidities are some of the

reasons people attend schools for the physically challenged. Poor night vision (79.3%) was the next disturbing symptom in this review and had literally forced those affected to abandon night or dim light activities. Onakpoya et al³³ and Ukponmwan et al³² had reported night blindness (58.3%) and 56.7% respectively which is lower than that of the present study. The differences could have arisen for the fact that Onakpoya's study was a multicentre study with higher participants. Harton et al² had earlier reported that night blindness is usually the initial symptom followed by loss of visual field and visual acuity as the disease progresses. Other complaints noted in this study include headache (31.0%) eye ache (27.6%) and photophobia (20.6%). Onakpoya³³ reported eye ache (10.4%). These headache, eye ache and photophobia could arise as symptoms of refractive error which is a co-morbidity of retinitis pigmentosa. The prevalence of refractive error (27.6%) myopia (20.7%) and hypermetropia (6.9%) were the findings in this review among the retinitis pigmentosa patients.

Onakpoya³³ had reported prevalence of refractive error (37.5%) which was higher than that of the present survey and both studies (present study and Onakpoya's) found myopia to be the most common refractive error (20.7%) and (71.4%) respectively. Myopia is the most common refractive error seen in retinitis pigmentosa patients. In myopia, pupils are somewhat large and a bit sluggishly reacting and may result in photophobia⁴⁰. It had been documented by LU and SU⁴¹ that patients with retinitis pigmentosa may be misdiagnosed and treated simply as myopic cases. Hypermetropia (6.9%) was noted in this study among the retinitis pigmentosa patients – while Onakpoya et al³³ in their multicentre study reported 11.1%. Other co-morbidities noted in this study include Glaucoma, unocular Aphakia (which was due to coaching). Another author³³ had documented prevalence of glaucoma (11%) which was lower than that of the present study

(17.2%), but reported lens opacity/pseudophakia (10.4%) among retinitis pigmentosa patients which was similar to that of the present study (10.5%). Other authors^{11,32,42,43}, had documented association of these disorders with retinitis pigmentosa patients. Nevertheless, the glaucoma rate in this study is higher than the 7.5% noted in Cameroon¹¹ and 2.3% in people's Republic of China⁴³. The fundus findings of the patients in this study are retinal pigmentation (100%) which is the commonest and this has been collaborated by other authors^{3,33}.

The typical perivascular pigmentation seems to make the diagnosis of retinitis pigmentosa unequivocal. Other fundus findings were retinal vascular attenuation, disc pallor and is similar to that of other authors^{3,33}. Disc cupping, though one of the fundus findings in this study is only secondary to glaucoma which is a co-morbidity of retinitis pigmentosa. No positive family history of retinitis pigmentosa was elicited in this study and this may be attributed to poor family health record among the relatives or the disease has been misdiagnosed in the past⁴¹. It may also mean that the disease has skipped generations. However, retinitis pigmentosa can also cause psychological effects.⁴⁴

Conclusion: Retinitis pigmentosa is a bilateral degenerative, progressive and inheritable disease of the retina that affects people in their active and productive stage of life. It affects the quality of life and livelihood of the affected with severe economic and social denigration. Early diagnosis, occupational, career and economic re-orientation and rehabilitation should be encouraged. Genetic counseling should also be a high point of health education of the affected.

Ethical approval

Ethical approval was sought and granted by the Ethical committee of Chukwuemeka Odumegwu Ojukwu University Teaching Hospital.

REFERENCES

1. Kanski J.J. Degenerative and dystrophies of the fundus. In Kanski J J ed. Clinical ophthalmology, systematic approach 3rd ed. New delhi Butter – worth Heinemann 1997; 381 – 425.
2. Hartong DJ, Berson EL, Dryja TP. Retinitis Pigmentosa. Lancet, 2006; 368:1795-1819. (Pub Med) Google scholar.
3. Khurana AK, Disease of the Retina. In Khurana AK ed. Comprehensive Ophthalmology 5th ed New Delhi, New Age International Ltd, 2012 263 – 306
4. Wan M, Lin H, Baiy et al: Clinical evidence in concurrence of retinitis pigmentosa and glaucoma. Chin Med. J 2011; 124: 1270 – 1274 (Pubmed) Google Scholar
5. Abah ER, Oladigbolu KK, Samalia E, Merali H, Ahmed AO, Abubakar TH. Ophthalmological abnormalities among deaf students in Kaduna. Northern Nigeria. Ann Afri.med 2011; 10:29-3. (pubmed) Google scholar
6. Chude EA, Nwosu SNN, Edema TO, Umuzuruike CN. Causes of visual loss in students attending school for the blind in south-Eastern Nigeria; Niger J Ophthalmol 2018; 28:46-50 (Google scholar).
7. Parmegiani F, Clinics epidemiology and genetics of retinitis pigmentosa. Curr Genomics. 2011; 2:236 – 237 (PMC Free article) (Pubmed) Google scholar
8. Bush H, Vinding T, La Cour M et al. Prevalence and causes of visual impairment and blindness among 9980 Scandinavian adults. The

- Copenhagen city Eye study. *Ophthalmology* 2004; 111; 53 – 61 (Pub med)
Google scholar
9. Al Merjan JI, Pandova MG, Al Ghanim M, et al Registered blindness and low vision in Kuwait *Ophthalmol Epidemiol*, 2005; 12:251 – 257.
 10. Oswald AH Goldblatt J, Sampson G Chokie R. Beighton P. Retinitis Pigmentosa in South Africa. *S Afr Med J* 1985; 68: 863 – 6 (pubmed)
Google scholar
 11. Eballe AO, Koki G. Emche CB Bella LA, Kouam JM, Melong J. blindness and visual impairment in retinitis pigmentosa. A Cameroonian hospital based study. *Clin Ophthalmol* 2010; 4:661-5 (Puc free article) pubmed (Google scholar)
 12. Ashaye A. O presumed hereditary retinal degeneration: Ibadan experience. *West Afri. J. Med* 2005; 24:49-53 (Pub med) (Google Scholar)
 13. Onakpoya OH, Olateju SO, Ajayi JA. Retinal disease in tertiary hospital, the need for establishment of a vitreoretina care unit. *J Natl Med Assoc.* 2008; 100:1286 -1289 (pubmed) Google scholar
 14. Nwosu SNN. Prevalence and pattern of retinal disease at the Guinness Eye Hospital Onitsha Nigeria *Ophthalmol Epidemiol*, 2007; 7:41-8 (Pub Med)
Google scholar
 15. Nwosu SNN, Ndulue CU, Ndulue OI, Uba – Obiano CU, Retinitis Pigmentosa in Onitsha. *Nigeria J West Afr. Coll Surg.* 2020 April – June, 10(2): 30 – 35. Doi:10.4103/Jwas – 65-21 E pub 2022 Mar 26. Pmcid: 35558569; Pmcid Pum 9089809.
 16. Facts about retinitis pigmentosa. National Eye institute. May 2014 retrieved 18 April 2020.
 17. Online Mendelian Inheritance in man (OMIM): Retinitis Pigmentosa RP 268000

18. Rivolta C, Sharon D, Deangelis MM, Dryja TP (2002). Retinitis Pigmentosa and allied disease numerous Diseases, genes and inheritance patterns, Human Molecular Genetics 11(10): 1219 – 27. Doi:10.1093/hmg/11.10.1219.pmid/20/5282
19. Berson, Eliot; Rosner, B; Sandberg, M.A Dryja, JP (1991) Ocular findings in patients with Autosomal Dominant retinitis pigmentosa and a rhodopsin Gene Defect (Pro – 23 - His) Archives of Ophthalmology.109(1):92 - 101.doi:10.100/archiloght.1991.01080094039.pmid.1987956
20. Gregory – Evans K, Pennesi, ME, Weleber RG Retinitis Pigmentosa and allied disorders. In Ryan SJ. editor. Retina 5th ed. Vol.2. New York Elsevier Saunders. 2013 PP 787 – 815 (Google scholar).
21. San Francisco, CA American Academy of Ophthalmology 2011, American Academy of Ophthalmology. Basic and clinical science course section 12 Retina and vitreors 2011 – 2012 pp. 228 – 36 (Google scholar)
22. Prokisch, Holger, Monika, Hellinger, Rosa Meitinger, Thomsa, Rosenberg, Thomas (2007) IVOS – A population – based epidemiological and genetic study of x – linked retinitis pigmentosa” investigative ophthalmology and visual science. 48(9): 4012 – 8. Doi:10.1167/iou.07-0071.pmid17724181.
23. Dryja TP, McGee TL Reichel E, Hahn LB; Cowley, GS; Yandell Dio Saudberg MA, Berson EL (1990). A poink mutation of the rehodopsm gene in one form of retinitis pigmentosa” nature 343 (6256) 364 – 6. Bibcode.1990Batur.43..364D.doi 10.1038/343364aO.pmid 2137202 S2CID4351328.
24. Dryja TP, McGee TL Hahn LB, Cowley GS Olsson JE, Rachiel E, Sandberg MA Bersob EL (1990) “Mutations within the rhodopsin Gene in Patients with Autosomal Dominant retinitis Pigmentosa” New England Journal of

- medicine 323(19): 1302 – 7.doi 10.1056/NEJM 199011083231903.PMID2215617.
25. Berson EL, Rosner B, Sandberg MA, Weigel – Difrancio C, Dryja TP (1991) ocular findings in patients with autosomol dominant retinitis pigmentosa and rhodopsin Proline – 347 – Leucine” American journal of Ophthalmology.111(5):614 -23 doi:10.1016/0002 – 9394(14) 73708 – 0. PMID 2021172.
 26. Inglehearn CF, Bashir R, Lester DH Jay M, Bird AC, Bhattacharya, SS(1991) “A 3-bp deletion in the Rhodopsin gene in Pigmentosa” American Journal of Heiman Genetics. 48(i); 26 – 30 PMC 1682750 PMID 1985460
 27. EzegwuiIR , Umeh RE, Ezepue UF. Causes of childhood blindness. Result form schools for the blind in South, Eastern Nigeria. Br J Ophthamol 2003; 87; 20 -3 PMC Free article) (Pub-med Google scholar)
 28. Haim M. Epidemiology of Retinitis Pigmentosa in Denmark. Acta Ophlthamol Scand.Suppl.2002;233:+1-3
 29. Openshaw A. (Feb. 2008). Understanding Retinitis Pigmentosa (PDF), University of Michigan Kellog Eye Centre. Archived from the original (PDF) on 2017 – 08 – 29. Retrieved 2017 – 12-02
 30. Iannaccone A, Berdia J. Rare Disease Database National Organization for Rare disorders (NORD) 55 Kenosia Ave; (Danbury) CT06810. (203) 744-0100.
 31. Hammel C. Retinitis Pigmentosa. Orphanet. J Rare. Dis. 2006; 1:40.
 32. Ukponmwan CU, Atama A. Retinitis Pigmentosa in Benin – Nigeria. East Afr. Med. J. 2004; 81:254 – 7 (pubmed) Google scholar
 33. Onakpoya OH, Adeoti C O, Oluleye TS Ajayi IA, Majengbasan T, Olarundare OK Clinical presentation and visual status of retinitis

- pigmentosa, A multicentre study in southwestern Nigeria Clin Ophthalmol. 2016;10:1578-83 (PMC free article) Pub med Google scholar.
34. Tsujikawa M, Wada Y, Sukegawa M et al Age at onset curves of retinitis pigmentosa Arch Ophthalmol. 2008; 126:337-340
 35. Marmor MF. Visual loss in retinitis pigmentosa AMJ. Ophthalmol.1990;89:692-698
 36. Briensen S, Roberts H, Finger RP. The impact of visual impairment on health related quality of life in rural Africa. Ophthalmic Epidemiol.2014;21(5):297-306
 37. Grover S. Fishman GA, Alexander KR, Anderson RJ Derlacki DJ visual acuity impairment in patients with retinitis Pigmentosa, Ophthalmology 1996; 103; 1593 – 1600
 38. Khurana AK community Ophthalmology. In Khurana AK ed. Comprehensive Ophthalmology 5th ed. New Delhi; New Age International Ltd. 2012.443 – 458
 39. Abah ER, Oladigbolu K.K, Samaila E. Merali H, Ahmed AO, Abubakar TN, Ophthalmologic abnormalities among deaf students in Kaduna, Northern Nigeria Ann Afr. Med 2011;10:29 -33 (Pubmed) Google scholar.
 40. Khurana AK. Optics and refraction in Khurana AK ed. Comprehensive Ophthalmology 5th ed. New Delhi, New Age International Ltd 2012
 41. LU Y, SUN X, Retinitis Pigmentosa Sine Pigmento masqueraded as myopia. A case report. Medicine 2021;100: e24006 (PMC – free article) (pubmed) Google scholar
 42. Pruett RC. Retinitis Pigmentosa. Clinical observations and correlation. Trans AM Ophthalmol Soc. 1983; 81:693-735
 43. Peng DW Retinitis Pigmentosa associated with glaucoma. Zhonghua Yan Ke za Zhi. 1991; 27:262-264.

44. Michael C Okosa, Richard Uwakwe, Akunne I Apakama, Arinze A Onwuegbuna, Chukwudi C Uzozie, Miriam-Benigna C Amobi, Chinwe A Uzuke. Psychological Effects of Eye Diseases: A Tertiary Eye Center Study. *Journal of Psychiatry and Psychiatric Disorders* 5 (2021): 128-139.