

# Utilization of bumblebee in crop pollination

---

## ABSTRACT

Bumblebees play a vital role in pollination of plants due to their large size and hairiness hence are ideal for picking up and transferring pollen. They can vibrate ('buzz') flowers and have no communication system but can work efficiently in tunnels as they have a better sense of direction. They are 400 times more efficient than honeybees at any pollination chore and are capable of visiting 30 to 50 flowers per minute. They are non-swarming and less aggressive than honeybees. Bumblebees generally forage during the early morning (0530-0800h) and evening (1700-1900h) time. Bumblebees are better pollinators than honeybees in greenhouses and glass houses and they have a great potential to serve as supplemental pollinators in the cultivation of crops such as tomato, eggplant, cucumber, melon, strawberry, pumpkin, cherry, sweet pepper, etc. It improves the yield and quality parameters of the fruits. In India, efforts are being made at Solan to mass rear bumble bees and utilize them in pollination. A few private companies have attempted to introduce newer species of bumblebees in India. In the present era, the global population of bumblebees is declining due to injudicious use of pesticides, loss of natural colony and mechanization in agriculture and climatic effect.

**Keywords:** Bumblebees, pollinator, foraging behavior

## INTRODUCTION

"Bumblebees are social insects, belonging to order Hymenoptera, tribe Bombini and genus *Bombus* in the family Apidae. There are over 250 known species of bumblebees in the world, primarily existing in the Northern Hemisphere although they occur in South America also. They have been introduced to New Zealand and the Australian state of Tasmania". Chauhan et al., (2013)<sup>b</sup> Of the 400 species of bumblebees worldwide (Heinrich, 1979), the six species commonly found are viz., Buff-tailed bumblebee, *Bombus terrestris*; White-tailed Bumblebee, *Bombus lucorum*; Garden Bumblebee, *Bombus hortorum*; Red-tailed Bumblebee, *Bombus lapidaries*; Early Bumblebee, *Bombus pratorum* and Common Carder Bee, *Bombus pascuorum*. In India, *Bombus haemorrhoidalis* is reported by Chauhan et al., (2013)<sup>b</sup> at Solan.

"Bumblebees live in organized colonies in nests typically built in a small hole in the ground or under a clump of grass. The three castes of bumblebees are the queen, female workers and males. Bumblebees have an annual life cycle. Bumblebee colonies infested by pests and diseases like mite, moth, conopid flies, nematode and *Nosema* spp". Chauhan et al., (2013)<sup>b</sup> They are robust, hairy and black with

yellow bands and make a buzzing sound while flying. However, some species have orange or red bands on their bodies or may be entirely black. A rarity, the bumblebee is a warm-blooded insect, by “shivering” her massive flight muscles; she can create heat in her thorax. Then, by contracting her muscles, she circulates the warmth into her abdomen. Bumblebees require pollen as a protein source for feeding their larvae and nectar as a sugar (carbohydrate) source - a fuel for the adults’ flight and to produce wax for nest building.

Bumblebees live in organized colonies in nests typically built in small holes in the ground or under clumps of grass. The size of queen cells is 16.56-20.57 mm in length and 12.55-15.20 mm width, the size of worker cells is 13.16-14.24 mm in length and 10.01-10.95 mm in width. Honey pots are 15.31-20.35 mm in length and 12.91-16.32 mm in width. The three castes of bumblebees are the queen, female workers and males. The body length of the queen is 20-25 mm, the worker is 18-25 mm and that of the male is 14-16 mm. Bumblebees have an annual life cycle and pass through four stages (egg, larva, pupa and adult) during their life span. The queen lays eggs in batches of 5-20 on the ball of pollen in the pots, seals it with wax, and incubates the eggs to keep them warm. Eggs hatched in 2.6 days after laying and the larval period is 17.2 days while the average pupal period was 8.6 days. The average total period from egg to adult is 28.4 days. Bumblebee colonies are infested by pests and diseases like mites, moths, conopid flies, nematodes and *Nosema* spp. Bumblebees forage on different kinds of flowers visiting about 8-12 flowers per minute. To collect pollen and nectar from a single flower it spends 4-8 seconds (Chauhan *et al.*, 2013<sup>a</sup>).

### **Role of Bumblebee as a Pollinator:**

“Bumblebees act as an important pollinator of plants due to being large, very hairy and so are ideal for picking up and transferring pollen. Bumblebees are active at temperatures near 5°C as well as on cloudy, foggy and rainy days. It will fly in winds of up to 40 mph (70 km/hr). The bumblebees can vibrate (‘buzz’) flowers and have no communication system. Bumblebees are non-swarming and less aggressive than honeybees. Bumblebees work better in tunnels as they have a better sense of direction. They are 400 times more efficient than honeybees at any pollination chore, capable of visiting 30 to 50 flowers per minute. Bumblebee generally forages during early morning (0530-0800h) and evening (1700-1900h) time” (Chauhan *et al.*, 2013<sup>b</sup>). Bumblebees are good pollinators in greenhouses and poly house and are of great economic importance in greenhouse cultivation of such as tomato, brinjal, strawberry, cucumber, melon, sweet pepper, cherry and pumpkin than honeybees.

### **Bumblebee in crop pollination:**

Abak *et al.* (1995) observed that the bumblebee pollinated plants of eggplant and tomato recorded significantly higher fruit yield (23% in eggplant and 17% in tomato), number of fruits per m<sup>2</sup> and number of seeds per fruit. However, there was no significant differences observed in fruit quality parameters like total soluble solids content, pH of fruit juice, titratable acidity, fruit diameter and fruit length in both crops as compared to control. In case of pepper, Abak *et al.* (1997) reported that “average yield, weight of fruit, diameter of fruit and the numbers of seeds increased by 4.0%, 10.0%, 6.0% and 12.5%, respectively in the bumblebee pollination as compared to control in greenhouse”.

“Tomato flowers pollinated by *B. vosnesenskii* produced larger fruit than manually pollinated flowers as evidenced by significant increases in fruit weight, height, diameter and seed count. Overall, bumblebee treatments (bumblebees and bumblebees plus manual) resulted in significantly greater fruit weight, seed count, fruit diameter and fruit height compared with non-bee treatments (manual and no-pollination). In contrast, fruit roundness did not differ between treatments” (Dogterom *et al.*, 1998).

In cucumber and watermelon, the flowers visited by bumblebees had significantly lower abortion rates and higher seed sets than those visited by honeybees (Stanghellini *et al.*, 1998).

Eggplants pollinated by bumblebees yielded higher (25%) than those that had vibration treatment with bigger fruit sizes (14% in weight and 7% in length) and four times higher number of seeds per fruit (Abaket *et al.*, 2000). Bumblebee pollinated plants of strawberries produced 30 per cent higher yield than control plants (Paydaset *et al.*, 2000).

Significantly higher tomato yield (4.12 kg m<sup>-2</sup>) in bumblebee pollinated plants with an increased yield of 89.9% and 60.9% over vibration treatment and growth regulator application respectively was observed. The quality of tomato like number of seeds per fruit, volume of fruits and number of carpels were also significantly higher (Dasgan *et al.*, 2004). Similarly, visits by bumblebees produced larger and heavier fruits and higher yields of sweet pepper as compared to self-pollination (Serrano and Jose, 2006).

In strawberries, the pollination done by bumblebees had a significant impact on the number and weight of marketable fruits produced. The fruit yields were higher in comparison to the control group. In control group, 30% of the fruits produced were deformed due to insufficient pollination (Dimouet *et al.*, 2008).

Chauhan, 2011 noted that in cucumbers pollination through bumblebee produced high quality fruits than without pollinators. Bumblebee supplementation led to higher pumpkin yields: 16.5 lbs/plant compared to 15.2 lbs/plant with honey bee supplementation and 13.1 lbs/plant in non-supplemented fields in New York (Nault *et al.*, 2011).

The tomato plants pollinated by bumblebees produced fruits with greater number of seeds compared to those pollinated mechanically or not pollinated at all. “Maturation time was longer and sugar content, fresh weight and seed count were higher in bumblebee pollinated flowers than those pollinated

manually or with no supplemental pollination; but did not differ with flowers pollinated mechanically”(Vergara and Buendia, 2012).

The positive impact of the bumblebees in bell pepper was found through the 3.77% increase in the number of fruits per plant and 24.6% higher fruit weight as well as 13.51% and 21.52% increases in fruit length and fruit breadth respectively. An increase of 23.84% in the number of healthy fruits, 113.64% more seed number and 5.44% increase in 1000 seed weight and 89.42% higher fruit yields per m<sup>2</sup> over control plots (Thakur, 2018). According to Marques *et al.*, (2019) fruit set was significantly higher in *B. terrestris* introduced field of almond as compared to control plots.

### **Impact of pesticides on bumblebee population**

“According to a study colonies treated with neonicotinoid (imidacloprid) suffered a significantly reduced growth rate and an 85% decline in production of new queens as compared to control colonies. It was suggested that the negative impact of these chemicals was considerable on wild bumblebee populations across the developed world”(Whitehorn *et al.*, 2012).

“Bumble bee, *Bombus terrestris* workers survived fewer days when exposed with syrup dosed at 98 µg thiamethoxam kg<sup>-1</sup>, while production of brood (eggs and larvae) and consumption of syrup and pollen in microcolonies were significantly reduced by thiamethoxam only at the two highest concentrations (39, 98 µg kg<sup>-1</sup>). In contrast, no detectable effect of thiamethoxam at levels typically found in the nectars of treated crops (between 1 and 11 µg kg<sup>-1</sup>) and also brood production in worker bumble bees is more sensitive to imidacloprid than thiamethoxam” (Laycock *et al.*, 2014). Baronet *et al.*, 2017 reported that “exposure of thiamethoxam cause 26% reduction in the proportion of queens that laid eggs, and advanced the timing of colony initiation”.

### **Conclusion:**

Pollination is one of the most important ecological processes on the planet. Among pollinators, bumblebees have gained importance as an efficient pollinator due to their large size and very hairy makes them more ideal for picking up and transferring pollen of many commercial crops grown under greenhouse, glasshouse, poly houses and open field. Bumblebees are better pollinators than honeybees in greenhouses and glass house and it have a great potentiality to serve as a supplemental pollinator in cultivation of crops such as tomato, eggplant, cucumber, melon, strawberry, pumpkin, cherry, sweet pepper *etc.* and resulted in to produced higher yield and increase the number of fruit and seed, fruit diameter, fruit length, fruit weight and also improve the quality of the fruit in crops and thus can render farming more profitable. Pesticides including neonicotinoid (imidacloprid) have been proven to the cause of bumblebee population decline and may be avoided in zones with crops that require pollination by bumblebees.

## References:

- Abak, K., Dasgan, H. Y., Ikiz, O., Uygun, N., Sayalan, M., Kaftanoglu, O. and Yeninar, H. (1997). Pollen production and quality of pepper grown in unheated greenhouses during winter and the effects of bumblebees (*Bombus terrestris*) pollination on fruit yield and quality. *Acta Horticulture*. 437: Proc. Int'l symp. on pollination. Pp. 303-308.
- Abak, K., Ozdogan, A. O., Dasgan, H. Y., Derin, K. and Kaftanoglu, O. (2000). Effectiveness of bumble bees as pollinators for eggplants grown in unheated greenhouses. *Acta Horticulture*, 514: Proc. XXV IHC- part 4. Pp. 197-203.
- Abak, K., Sari, N., Paksoy, M., Kaftanoglu, O. and Yeninar, H. (1995). Efficiency of bumble bees on the yield and quality of eggplant and tomato grown in unheated glasshouses. *Acta Horticulture*, 412: I Int'l symp. on Solanacea for Fresh Market. Pp. 268-274.
- Baron, G. L., Jansen, V. A. A., Vincent A. A., Brown, M. J. F. and Raine, N. E. (2017). Pesticide reduces bumblebee colony initiation and increases probability of population extinction. *Nature Ecology & Evolution*, 1 (9): 1308–1316.
- Chauhan, A. (2011). Refinement of bumblebees rearing technology and its use in cucumber pollination. M. Sc. Thesis, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, India, 98.
- Chauhan, A., Katna, S. and Rana, B. S. (2013<sup>a</sup>). Ethological studies on bumble bees under subtropical conditions in India. *Journal of Medical Science And Clinical Research*, 1 (3) : 125-129.
- Chauhan, A., Katna, S., Rana, B. S. and Sharma, H. K. (2013<sup>b</sup>). Studies on pests and diseases of bumble bee (*Bombus haemorrhoidalis* Smith) in India. *Journal of Medical Science And Clinical Research*, 1 (2) : 93-98.
- Dasgan, H. Y., Ozdogan, A. O., Kaftanoglu, O. and Abak, K. (2004). Effectiveness of bumblebee pollination in anti-frost heated tomato greenhouses in the Mediterranean basin. *Turkish Journal of Agriculture and Forestry*, 28 : 73-82.
- Dimou, M., Taraza, S., Thrasyvoulou, A. and Vasilakakis, M. (2008). Effect of bumble bee pollination on greenhouse strawberry production. *Journal of Apicultural Research. Bee World*, 47 (2) : 99-101.
- Dogterom, M. H., Matteoni, J. A. and Plowright, R. C. (1998). Pollination of greenhouse tomatoes by the North American *Bombus vosnesenskii* (Hymenoptera: Apidae), *Journal of Economic Entomology*, 91 (1) : 71-75.
- Heinrich, B. (1979). Majoring and minoring by foraging bumblebees *Bombus Vagans*. *Ecology*, P 245-255.
- Laycock, I., Cotterell, K. C., O'Shea-Wheller, T. A., Cresswell, J. E. (2014). Effects of the neonicotinoid pesticide thiamethoxam at field-realistic levels on microcolonies of *Bombus terrestris* worker bumble bees. *Ecotoxicology and Environmental Safety*, 100 : 153-158.

- Marques A, Juan A, Ruiz M, Traveset A, Leza M. (2019). Improvement of almond production using *Bombus terrestris*(Hymenoptera: Apidae) in Mediterranean conditions. *Journal of Applied Entomology*, **143** (10): 1132-1142.
- Nault, B. A., Artz, D. R. and Petersen, J. D. (2011). Proc. of the New England vegetable and fruit conf., Pp.45-47.
- Paydas, S., Eti, S., Kaftanoglu, O., Yasa, E., Derin, K. (2000). Effects of pollination of strawberries grown in plastic greenhouses by honeybees and bumblebees on the yield and quality of the fruits. *Acta Horticulturae*, 513: 443–451.
- Serrano, A. R. and Jose, G. M. (2006). Quality Fruit Improvement in Sweet Pepper Culture by Bumblebee Pollination. *Scientia Horticulturae*, 110 : 160-166.
- Stanghellini, M. S., Ambrose, J. T. and Schultheis, J. R. (1998). Using commercial bumble bee colonies as backup pollinators for honey bees to produce cucumbers and watermelons. *HortTechnology*, **8** (4) : 590-594.
- Thakur, S. (2018). Studies on bumblebees (*Bombus haemorrhoidalis* Smith) pollination in *Capsicum annuum* L. under protected cultivation. M.Sc. Thesis, Department of Entomology, Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India.
- Vergara, C. H. and Buendia, P. F. (2012). Pollination of greenhouse tomatoes by the Mexican bumblebee *Bombus ephippiatus* (Hymenoptera: Apidae). *Journal of Pollination Ecology*, **7** (4) : 27-30.
- Whitehorn, P. R., Stephanie, C., Wackers, F. L. and Dave, G. (2012). Neonicotinoid pesticide reduces bumble bee colony growth and queen production, *Science express*. 29 March : 351-352.

\*\*\*\*\*