

# Studies on Genetic Divergence ~~analysis~~ Analaysis in Black gram genotypes (*Vigna mungo* L. Hepper)

**Comment [u1]:** The manuscript should revise by a native speaker. English needs to be improved

## Abstract

**Comment [u2]:** In one or two sentences, state the importance of the subject and then state the purpose and results

The present investigation on Genetic Divergence and clustering patterns was carried out with twenty-eight diverse genotypes of Black gram using Mahalanobis's  $D^2$  Statistics. The experiment was conducted in Randomized Block Design with three replications at Plant Breeding Farm, Department of Genetics and Plant Breeding, Faculty of Agriculture, Annamalai University, Chidambaram. Observations were recorded on nine various Biometrical characters viz. Plant height at maturity (PH), length of the pods (LP), number of branches per plant (NBPP), number of clusters per plant (NCP), number of pods per cluster (NPPC), number of pods per plant (NPPP), number of seeds per pod (NSPP), 100 seed weight (SW), and seed yield per plant (SY). Analysis of variance revealed a substantial amount of variability for all eleven characters. Mahalanobis ( $D^2$ ) static revealed considerable genetic diversity among the genotypes. Genotypes were grouped into 6 clusters. Cluster VI was the largest including 11 genotypes. The ~~intra-intra~~ cluster distance for cluster IV is the maximum and the lowest intra cluster distance was reported for cluster II. The highest inter cluster distance was revealed between cluster I and III and then between clusters II and III. ~~The~~ This clearly indicates that the genotypes included in this cluster are having a broad spectrum of genetic diversity and could very well be used in the hybridization programme of black gram for improving yield. The minimum ~~inter-inter~~ cluster distance was revealed between clusters I and II. The percentage contributed to total divergence by various characters are seed yield was higher (50.997) followed by 100 seed weight (29.456) and number of pods per plant (8.121). It is suggested that, crosses among the parents belonging to most divergent clusters would be expected to manifest maximum heterosis and also wide variability of genetic architecture.

**Comment [u3]:** Do you have any recommendations for the beneficiaries and the scientific community?

## Introduction

Black gram (*Vigna mungo* L. Hepper), also called urd bean is a member of the Asian *Vigna* crop species (Delic *et al.*, 2009). Black gram is also known as Mash in India and is a grain

legume domesticated from *V. mungo var. silvestris*. The major portion of black gram is utilized in making dal, curries, soup, sweets and snacks. Its seeds contain approximately protein (24- 26%), carbohydrates (60%), fat (1.5%), fiber (3.5- 4.5%), ash (4.5-5.5%), minerals, amino acids and vitamins. It is the richest sources of phosphoric acid being 5-10 times richer than other crops. It belongs to family fabaceae. Center of genetic diversity for black gram is found in India (Zeven *et al.*, 1982). It is grown in an area of about 5.44 million hectares in India with production of 3.56 million tonnes and productivity of 653 kg/ha (Anonymous, 2017-18). In spite of its importance, the productivity of this crop is relatively low. India is the largest producer and consumer of black gram in the world. Black gram has been distributed mainly in tropical to sub-tropical countries where it is mainly grown in India, Pakistan, Sri-Lanka, Burma, and some countries of South East Asia. The development of new varieties depends largely on the availability of genetic variability in the base material and the extent of variability for the desired character. The development of cultivated species and breeding of new varieties typically relies on the available biological diversity in existing genotypes (Datta and Lal, 2011). Limited variability has been exploited in varietal development programmes in black gram (Jayamani and Sathya, 2013). The breeding progress has been slow and uneven because several desirable traits need to be combined for developing appropriate plant type for a particular growing region and cropping system in black gram. Genetic diversity is a pre-requisite for any crop improvement program as it helps in estimating and establishing genetic relationship in germplasm collection, identifying diverse parental combinations to create segregating progenies with maximum genetic variability and superior recombinations for further selection and introgressing desirable genes from diverse germplasm. The  $D^2$  analysis proposed by Mahalanobis (1936) is an effective tool in quantifying the degree of genetic divergence among the genotypes. Keeping the above in view, the present study was undertaken to identify the best performing germplasm of black gram based on quantitative traits using Mahalanobis  $D^2$  statistics and Tocher's method. In the present study, genetic divergence and clustering pattern of the black gram genotypes for selection of suitable parents so as to utilize them in hybridization programme, extended to study the genetic parameters attributing to yield.

## Materials and Methods

The material under investigation consisted of twenty-six genotypes of black gram (*Vigna mungo* L. Hepper) were grown in kharif 2022 at the Plant Breeding Farm, Department of

**Comment [u4]:** First, state the different methods to achieve the purpose of the study, and then state the advantages of your preferred method. Otherwise, the purpose of the research is incompletely stated.

**Comment [u5]:** June to November

Genetics and Plant Breeding, Faculty of Agriculture, Annamalai University is situated in 11.39° North latitude 79.71° East longitude and 12 km away from Bay of Bengal with an altitude of 6m above sea level. The climate is relatively warm and humid. The genotypes used are listed at table-1. Experiment was done according to randomized block design with three replications, All the recommended package of practices was followed for raising healthy and productive crop with proper agronomic measures. Each plot consisted of a three row of 5m length with a spacing of 25 cm between rows and 15 cm between plants was maintained by proper thinning. Observations were recorded on single plant basis. For recording single plant observations, five competitive plants from each replication were randomly selected. Average of these five plants in respect of plant height at maturity, length of the pods, number of branches per plant, number of clusters per plant, number of pods per cluster, number of pods per plant, number of seeds per pod, 100 seed weight, and seed yield per plant was used for statistical analysis. Mahalanobi's  $D^2$  statistic was employed to assess the genetic diversity. The genotypes were collected from Tamil Nadu Agriculture University (Coimbatore), National Pulses Research Centre (Vamban) and some local regions of Tamilnadu.

**Comment [u6]:** Meteorological statistics should be given in a table or a sentence.

**Table 1. List of Black gram Germplasm used for Genetic diversity analysis**

Accession No.	Genotypes	Accession No.	Genotypes
G1	K 1	G15	Omalur local
G2	Erode local	G16	Mettur local
G3	ADT 3	G17	Madurai local
G4	Shankari local	G18	VBN 5
G5	Chidambaram local	G19	PAIYUR 1
G6	Sirkali local	G20	TMV 1
G7	Salem local	G21	CO 1
G8	ADT 5	G22	CO 2
G9	VBN 3	G23	CO 5
G10	T 9	G24	CO 6
G11	APK 1	G25	VBN 1
G12	VBN 4	G26	VBN 2
G13	VBN 6	G27	Aatur local

G14	VBN 7	G28	Sethiyathoppu local
-----	-------	-----	---------------------

## Results and Discussion

The mean sums of squares of nine different traits are analysed using ANOVA. High significant differences for all characters under study among the twenty-eight black gram genotypes were found in analysis of variance, at 5 % level of significance indicating the presence of sufficient considerable genetic variability among different genotype. These results were in agreement with the findings of Balachandran *et al.* (2010), Kumar *et al.* (2015), Priyanka *et al.* (2016), Rolaniya *et al.* (2017) and Nagmi and Lal (2017). The pattern of clustering proved the existences of significant amount of variability. It is obvious that the genotypes have grouped into different cluster irrespective of their geographical origins. It means that the genetic constitution of the varieties was more important than their origin and distribution (Mehandi *et al.*, 2013). Genetic divergence analysis was widely used to determine the genetic relationship among the genotypes and find out the suitable genotypes for future breeding programme. Genetic diversity analysis also helps in tagging and elimination of the duplicate accessions from genetic stock. On the basis of divergence 28 genotypes under investigation have been grouped into six distinct clusters given at Table 2, indicating a wide range of diversity in the experimental materials.

**Table 2. Composition of D<sup>2</sup> clusters for 28 Black gram Germplasms**

Clusters	List of Genotypes of Black gram under each clusters	Total number of genotypes
I	G1, G2, G4, G10	4
II	G7, G16, G27	3
III	G3, G8, G11	3
IV	G5, G6, G15, G17, G28	5
V	G19, G20	2
VI	G9, G12, G13, G14, G18, G21, G22, G23, G24, G25, G26	11

Among the five clusters, cluster VI was the largest including 11 genotypes followed by cluster IV had 5 genotypes, cluster I had 4 genotypes, cluster II and cluster III had 3 genotypes each and cluster V had 2 genotypes. These findings suggested that genetic divergence may not be strongly influenced by geographic diversification. The genotypes in a cluster came from a variety

of geographic origins, demonstrating that genetic diversity is not necessarily correlated with geographic divergence. The divergence among the genotypes in the same cluster is indicated by the divergence within the cluster. Inter cluster divergence, one on either hand, denotes the divergence between the genotypes of various two groups. Based on main character relations, a critical analysis of clusters revealed that they were heterogeneous within and between one another. The lower  $D^2$  value between their characters indicated a close genetic relationship between these genotypes in one cluster and those in the other cluster. These findings are supported with the results obtained by Manikannan *et al.* (2000), Lad *et al.* (2005), Konda *et al.* (2007), Majumdar *et al.* (2011) and Panigrahi *et al.* (2014) in black gram.

**Table 3. Average inter and intra cluster  $D^2$  values for 28 Black gram genotypes**

Clusters	I	II	III	IV	V	VI
I	8.923	5.865	80.791	22.35	52.134	20.86
II		1.994	80.438	18.315	44.364	16.342
III			4.487	51.231	20.003	50.762
IV				22.85	28.199	20.21
V					6.491	25.223
VI						16.596

It is revealed in table-3, that the intra cluster distances were found Lower than the inter cluster distance revealing a considerable amount of genetic diversity among the genotypes under various clusters. The highest inter cluster distance was revealed between cluster I and III (80.791) followed by cluster II and cluster III (80.438), cluster I and cluster V (52.134), cluster III and cluster IV (51.231), cluster III and cluster VI (50.762), cluster II and cluster V (44.364), cluster IV and cluster V (28.199), cluster V and cluster VI (25.223) and between cluster I and cluster IV (23.383). The genotypes grouped in these clusters can be used in breeding programme in order to get a wide spectrum of variability and transgressive segregates. The minimum distance between clusters I and II (5.865) indicated that they were genetically closure clusters. Selection of parents from such clusters may be avoided because it may result in narrow genetic base. This result is supported by findings of Lad *et al.* (2005), Konda *et al.* (2007), Umadevi and Ganesan (2007) and Katna and Verma (2003) in black gram. The intra cluster distance among

various clusters exhibited maximum intra cluster distance for cluster IV (22.85) and lowest intra cluster distance was reported for cluster II (1.994). The maximum intra cluster distance was because of wide genetic diversity among its genotypes. The chance of developing good segregates by crossing the genotypes of the same cluster showing low value for intra cluster distance are very low. Therefore, it would be logical to attempt crosses between the genotypes of clusters separated by larger inter cluster distance. The little diversity and selection of parents within the clusters having higher mean for a particular character may also be useful for further developing high yielding black gram varieties.

**Table 4. cluster mean of 28 black gram genotypes for observed traits**

Clusters	PH	NBPP	NCPP	NPPC	NPPP	NSPP	LP	SW	SY
I	30.12	2.56	7.90	4.21	32.67	4.56	4.60	3.99	6.86
II	28.52	2.51	8.12	4.43	33.31	4.74	4.64	4.21	6.31
III	31.32	3.01	6.87	5.90	36.21	6.27	4.98	5.28	9.07
IV	28.90	2.53	7.95	4.64	34.61	4.62	4.50	4.52	6.65
V	27.86	2.33	5.91	5.17	29.97	4.83	4.86	5.11	6.71
VI	29.33	2.78	7.21	5.03	33.44	4.73	4.72	4.41	6.70

(Plant height at maturity (PH), length of the pods (LP), number of branches per plant (NBPP), number of clusters per plant (NCPP), number of pods per cluster (NPPC), number of pods per plant (NPPP), number of seeds per pod (NSPP), 100 seed weight (SW), and seed yield per plant (SY))

The cluster mean for each eleven characters are presented in Table 4. The genotypes of cluster III were higher for 100 seed weight (5.28) and pod length (4.98) followed by cluster V (5.11) and (4.86) respectively. The genotypes fall in the cluster III were having the desirable direction of characters *i.e.*, plant height (31.32), number of branches per plant (3.01), number of pods per plant (36.21), number of pods per cluster (5.90), number of seeds per pod (6.27) and seed yield per plant (9.07). followed by cluster I for plant height, number of branches per plant, number of clusters per plant, number of pods per plant are high. It is suggested that, crosses among the parents belonging to most divergent cluster would be expected to noticeable maximum heterosis and also wide variability of genetic architecture. Thus, the crosses between the genetically diverse genotypes of cluster I and III, I and II, I and V, III and IV, III and VI, II

and III, III and V, IV and V, V and VI, I and VI are expected to exhibit high heterosis and are also likely to produced new combinations with desired characters to get desirable segregates with higher yield for developing superior varieties of black gram.

**Table 5. contribution of different characters for genetic diversity of Black gram**

Sl.No	Observed Characters	Percentage of contribution
1	Plant Height at Maturity	2.988
2	Number of Branches	1.005
3	Number of clusters per plant	2.220
4	Number of pods per cluster	0.782
5	Number of pods per plant	8.121
6	Number of seeds per plant	0.754
7	Length of the pod	3.661
8	100 seed weight	29.456
9	Seed yield	50.997

The percentage contributed to total divergence of various characters are presented in Table 5. The results showed that the contribution of seed yield was higher (50.997) followed by 100 seed weight (29.456) and number of pods per plant (8.121) then other characters contribution is pod length was (3.661), plant height (2.988), number of clusters per plant (2.220), number of branches per plant (1.005) and lower contribution was made by number of seeds per pod (0.754) and number of pods per cluster (0.782) towards the genetic divergence.

### **Conclusion**

The 28 black gram germplasms used for the genetic diversity assessment demonstrated a significant range of variance. The 28 black gram germplasms were divided into 6 clusters, which was consistent with the Mahalanobis  $D^2$  clustering pattern. The parents for the hybridization programme should be chosen based on the genetic distance, the contribution of various characters to the total divergence, and the magnitude of the cluster means for various characters performing with the greatest heterosis. Thus the crosses between the genetically diverse genotypes of cluster I and III, I and II, I and V, III and IV, III and VI, II and III, III and V, IV

and V, V and VI, I and VI are expected to exhibit high heterosis and are also likely to produce new combinations with desired characters to get desirable segregates with higher yield for developing superior varieties of black gram.

## Reference

Anonymous. (2017-18). Pulses Revolution from Food to Nutrition, Directorate of Pulse Development, Dept. of Agri., Co-operation and Farmer Welfare. Ministry of Agriculture and Farmer Welfare, Govt. of India.

Balachandran, D., Mullainathan, L. Velu, S. and Thilagavathi, C. (2010). Study of genetic variability, heritability and genetic advance in blackgram. African J. of Biotechnology.9: 19, 2731-2735.

Datta J, Lal N. Genetic differentiation in *Cicer arietinum* L. and *Cajanus cajan* L. Millspaugh using SSR and ISSR marker systems. Advanced Biotech, 2011; 11(5):39-44.

Delic, D, Stajkovic, O, Kuzmanovic, D, Rasulic, N, Knezevic, S, Milicic, V. 2009. The effects of rhizobial inoculation on growth and yield of *Vigna mungo* L. in Serbian soils. Biotechnology in Animal Husbandry, 25(5-6): 1197-1202.

Jayamani, P. and Sathya, M. 2013. Diversity in pod characters in black gram [*Vigna mungo* (L.) Hepper]. Legume Res., 36(3): 220-223.

Kanta G, and Verma, S. (2003). Genetic divergence studies in black gram. National J. of Plant Improvement.5:1, 38-42.

Konda, C.R., Salimath, P.M. and Mishra, M.N. (2007). Genetic diversity estimation in black gram. Legume Res. 30: 3, 212-214.

Kumar, G. V., Vanaja, M., Lakshmi, N. J. and Maheshwari, M. (2015). Studies of variability, heritability and genetic advance for quantitative traits in black gram (*Vigna mungo* (L.) Hepper). Agric. Res. J. 52(4): 28-31.

Lad D.B., Punde, P.B., and Dahake, K.D. (2005). Genetic divergence studies in blackgram. J. of Maharashtra Agri. Universities.30: 2, 183-186.

**Comment [u7]:** Use recent references. 6 years have passed from your newest references.

**Comment [u8]:** Don't use Anonymous references

Nagmi, P. and Lal, G.M. (2017). Estimates of genetic variability and heritability for yield and yield component traits in blackgram [*Vigna mungo* (L.)]. *International Journal of agriculture Sciences*, 9(36): 4550-4552.

Priyanka S., Rangaiah S. and Showkath Babu B. M. (2016). Genetic Variability Estimates of Quantitative and Qualitative Traits in Black Gram. *International Journal of Agriculture Sciences*, Volume 8, Issue 40, pp.-1821-1824.

Umadevi, M. and Ganesan, N. M. (2007). D<sup>2</sup> analysis for yield and quality characters in blackgram (*Vigna mungo* (L.) Hepper). *Legume Res.*, 30 (3):197-200.

Zeven, A.C., and De Wet J.M.J. (1982). *Dictionary of cultivated plants and their regions of diversity*. Centre for Agricultural Publication and Documentation, Wageningen.

UNDER PEER REVIEW