

Systematic Review

Clinical Therapies Utilized in Treating Medication-related Osteonecrosis of the Jaw: a Systematic Review

ABSTRACT

Background and Aims: Medication-related osteonecrosis of the jaws (MRONJ) is a significant and potentially debilitating side effect caused by antiresorptive and antiangiogenic drugs, which can lead to bone exposure in the oral cavity. However, the management of this condition remains controversial, with adjuvant therapies being employed despite limited scientific evidence. This systematic review aimed to identify effective therapeutic procedures for treating MRONJ.

Methodology: A literature search was conducted without any temporal limitations. The PRISMA protocol was followed. To identify relevant studies, we developed electronic search strategies for various bibliographic databases, as Cochrane, Embase, PubMed, Scopus, and Web of Science. It was conducted a comprehensive analysis of 30 studies involving 2,079 patients from 35 countries to evaluate the effectiveness of various treatment approaches for MRONJ.

Results: The systematic review revealed that long-term use of Zoledronic acid for approximately 452.04 months (± 27.41 ; 12-102) exposed many patients ($n=772$) to the risk of MRONJ. Similarly, Alendronate use for approximately 104.4 months (± 60.16 ; 6-180) also posed a risk, affecting 650 patients, while Pamidronate use for about 20.74 months (± 4.94 ; 6-96) was associated with MRONJ risk in 121 patients. Among the treatment approaches, conservative surgical management was the most frequently employed (27.92%), followed by local debridement (13.57%) and conservative treatment (11.21%). Treatment complications were observed in 13.03% of cases, with the most frequent complications being resistant or worsening clinical stage of osteonecrosis, followed by incomplete mucosal healing or dehiscence and mental nerve injury.

Conclusion: While conservative surgical management, local debridement, and conservative treatment are commonly utilized approaches, the treatment of MRONJ lacks a standardized consensus due to the scarcity of scientific evidence. Further research and comprehensive studies are imperative to establish effective therapeutic strategies for managing this condition.

Keywords: Osteonecrosis; Bisphosphonates; Jaw; Treatment

1. INTRODUCTION

Drug-related jaw osteonecrosis is an oral lesion that affects the jawbone and the jawbone. This condition is currently characterized by exposed bone or bone that can be probed through an intra or extraoral fistula in the maxillofacial region. It is caused by an insufficiency of blood in the bone that can have various origins, such as bisphosphonates, trauma, infections, or radiation therapy. Medication-related osteonecrosis of the jaws (MRONJ) represents a significant and concerning adverse outcome linked to the use of antiresorptive and/or antiangiogenic medications, specifically bisphosphonates. Initially, it was characterized as avascular osteonecrosis triggered by Pamidronate and Zoledronate, as

delineated by Marx [1]. Subsequently, due to the escalating occurrence of osteonecrotic lesions, the term evolved to bisphosphonate-related osteonecrosis of the jaws (BRONJ), and the last M.R.O.N.J [1,2,3]. MRONJ is typified by necrosis of bone tissue and oral cavity exposure, arising from a metabolic imbalance in the mineral bone matrix induced by these medications. The American Association of Oral and Maxillofacial Surgeons (AAOMS) has meticulously established specific criteria for MRONJ diagnosis, encompassing a history of bisphosphonate treatment, prolonged bone exposure exceeding eight weeks, and the absence of radiotherapy or metastatic disease within the maxillofacial region [4].

1.1 Staging

1.1.1 Stage 0 (Nonexposed Bone Variant)

Patients with no clinical evidence of necrotic bone but who present with non-specific symptoms or clinical and radiographic findings, such as: odontalgia not explained by an odontogenic cause; dull, loosening of teeth not explained by chronic periodontal disease, intraoral or extraoral swelling, alveolar bone loss or resorption not attributable to chronic periodontal disease, changes to trabecular pattern sclerotic bone and no new bone in extraction sockets, regions of osteosclerosis involving the alveolar bone and/or the surrounding basilar bone, thickening/obscuring of periodontal ligament (thickening of the lamina dura, sclerosis, and decreased size of the periodontal ligament space).

These non-specific findings, which characterize this variant of OMRM without bone exposure, may occur in patients with a prior history of Stage 1, 2, or 3 disease who have been healed and have no clinical evidence of exposed bone. Progression to Stage 1 disease has been reported in up to 50 percent of patients with Stage 0 disease 41 percent; therefore, AAOMS deems it prudent to consider Stage 0 disease as a potential precursor to OMRM.

1.1.2 Stage 1

Exposed and necrotic bone or fistula that probes to the bone in asymptomatic patients with no evidence of infection/inflammation. These patients may also present with radiographic findings mentioned for Stage 0 localized to the alveolar bone region.

1.1.3 Stage 2

Exposed and necrotic bone, or fistula that probes to the bone, with evidence of infection/inflammation. These patients are symptomatic. These patients may also present with radiographic findings mentioned for Stage 0 localized to the alveolar bone region.

1.1.4 Stage 3

Exposed and necrotic bone or fistulae that probe to the bone, with evidence of infection, and one or more of the following: exposed necrotic bone extending beyond the region of alveolar bone (i.e., inferior border and ramus in the mandible, maxillary sinus and zygoma in the maxilla), pathologic fracture, extraoral fistula, oral antral/oral-nasal communication, osteolysis extending to the inferior border of the mandible or sinus floor [4].

Bisphosphonates are pivotal as antiresorptive agents in managing malignancies and fractures related to osteoporosis. Efficacious fracture reduction has been demonstrated in both orally administered forms (Alendronate and Residronate) and parenteral agents (Zoledronic acid and ibandronate) for osteoporosis patients [5,6]. However, it gradually became evident that other antiresorptive, as well as other classes of medications, were also involved in the MRONJ, such as the Denosumab, an antiresorptive human antibody that exhibited promising fracture risk reduction for osteoporosis patients [7].

MRONJ is associated with several risk factors, including tooth extractions, chemotherapy, steroid use, periodontal disease, and various medical conditions [8]. To diagnose MRONJ accurately, advanced imaging techniques like cone-beam computed tomography (CT) are often necessary to assess the extent of bone involvement comprehensively [9]. In human patients, MRONJ is classified based on clinical presentation. Stage 0 signifies the absence of clinical evidence of necrotic bone, but the patient may exhibit non-specific symptoms or radiographic findings. In such cases, the treatment approach typically involves pain management and, if necessary, antibiotics. Conversely, Stage 3 represents the most severe manifestation, where the patient presents with an infection and maxillary sinus involvement. The prognosis of

osteonecrosis of the jaws related to drugs in stage 3 is variable. It depends on several factors, such as the extent of the lesion, the presence of infection, the response to treatment, and the patient's general health [10].

Thus, the therapeutic landscape for MRONJ comprises nonoperative and operative modalities. Nonoperative strategies emphasize patient education, pain management, infection control, and the facilitation of necrotic bone sequestration [11]. Conversely, operative interventions have shown encouraging outcomes and are increasingly acknowledged as practical approaches across various stages of osteonecrosis [11]. However, the efficacy of specific therapies like hyperbaric oxygen or ozone therapy remains limited by insufficient evidence, warranting further exploration [12,13]. Supplementary treatments such as vitamin E and Pentoxifylline have been primarily reported in case studies, with ongoing research exploring their potential efficacy [14].

Despite the array of treatment options, a definitive consensus on the optimal MRONJ management protocol remains elusive. The AAOMS advocates a multidisciplinary approach involving dental, oncological, and maxillofacial specialists. Conservative management is endorsed for early-stage MRONJ, concentrating on infection control, necrosis management, and pain alleviation [15]. Proactive measures, encompassing optimization of oral health and dental care before initiating antiresorptive therapy, have been underscored in multiple studies. These preventive strategies mitigate the risk of MRONJ and contribute to the overall enhancement of oral health [16-19]. Thus, this systematic review provides a comprehensive overview of the primary therapies currently employed in treating MRONJ, shedding light on the evolving landscape of its management. This systematic review aims to identify forms of treatment for OMRM, treat present infection, manage pain associated with injury, and promote healing in the affected area.

2. METHODOLOGY

2.1 Protocol

The authors of this systematic review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) checklist [20,21] to ensure the thoroughness and transparency of the review process. The review encompassed a comprehensive literature search strategy, a meticulous selection process that relied on pre-established inclusion and exclusion criteria, meticulous data extraction, and a rigorous assessment of the quality of the studies included.

2.2 Study Design and Eligibility Criteria

This systematic review aimed to examine the clinical therapies utilized in medication-related osteonecrosis of the jaw. A literature search was conducted without any temporal limitations. The patient, intervention, comparison, outcome (PICOS) strategy was employed to formulate the research question, incorporating the following inclusion criteria: (i) population - patients diagnosed with medication-related osteonecrosis of the jaw, (ii) intervention - identification of clinical treatments administered, (iii) comparison - none, (iv) outcome - identification of clinical treatments implemented in patients with medication-related osteonecrosis of the jaw and their respective outcomes, and (v) study design - observational studies (cohort and case-control studies).

Studies were excluded based on the following criteria: i) studies involving patients unaffected by medication-related osteonecrosis of the jaw, ii) studies lacking reporting on clinical treatment of medication-related osteonecrosis of the jaw, or iii) reviews, letters, conference summaries, or personal opinions.

2.3 Information Sources and Search Strategy

To identify relevant studies, we developed electronic search strategies for various bibliographic databases, namely Cochrane (<https://www.cochranelibrary.com>), Embase (<https://www.embase.com>), PubMed (<https://pubmed.ncbi.nlm.nih.gov>), Scopus (<https://www.scopus.com>), and Web of Science (<https://www.webofscience.com/wos/woscc/basic-search>). The search encompassed articles published across all databases until July 25, 2022. Rayyan software reference manager was employed to remove duplicate articles. The search strategy was standardized and consistently applied for all subsequent updates. Supplementary Table 1 provides additional details regarding the search strategy employed for each database.

Supplementary Table 1. Search strategy in the databases.

Database	Search
PubMed (July 25, 2022)	"Diphosphonates"[Mesh] OR "Alendronate"[Mesh] OR "Etidronic Acid"[Mesh] OR "ibandronic acid" [Supplementary Concept] OR "pamidronate" [Supplementary Concept] OR "zoledronic acid" [Supplementary Concept] OR "risedronic acid" [Supplementary Concept] OR "denosumab" AND "Bisphosphonate-Associated Osteonecrosis of the Jaw"[Mesh] OR ("Osteonecrosis"[Mesh] AND "Jaw"[Mesh]) OR (osteonecrosis AND jaw) AND "Therapeutics"[Mesh]
Embase (July 25, 2022)	((('diphosphonates'/exp OR 'diphosphonates' OR 'alendronate'/exp OR 'alendronate' OR 'etidronic acid'/exp OR 'etidronic acid' OR 'ibandronic acid'/exp OR 'ibandronic acid' OR 'pamidronate'/exp OR 'pamidronate' OR 'zoledronic acid'/exp OR 'zoledronic acid' OR 'risedronic acid'/exp OR 'risedronic acid' OR 'denosumab'/exp OR 'denosumab') AND ('all fields' OR (all AND fields))) AND ('bisphosphonate-associated osteonecrosis of the jaw'/exp OR 'bisphosphonate-associated osteonecrosis of the jaw') OR (('osteonecrosis'/exp OR 'osteonecrosis') AND ('jaw'/exp OR 'jaw')) OR (('osteonecrosis'/exp OR osteonecrosis) AND ('jaw'/exp OR jaw))) AND ('therapeutics'/exp OR 'therapeutics')
Cochrane (July 25, 2022)	"Diphosphonates" OR "Alendronate" OR "Etidronic Acid" OR "ibandronic acid" OR "pamidronate" OR "zoledronic acid" OR "risedronic acid" OR "denosumab" in Title Abstract Keyword AND "Bisphosphonate-Associated Osteonecrosis of the Jaw" OR ("Osteonecrosis" AND "Jaw") in Title Abstract Keyword AND "Therapeutics" in Title Abstract Keyword - (Word variations have been searched)
Scopus (July 25, 2022)	(TITLE-ABS-KEY ("Diphosphonates" OR "Alendronate" OR "Etidronic Acid" OR "ibandronic acid" OR "pamidronate" OR "zoledronic acid" OR "risedronic acid" OR "denosumab") AND TITLE-ABS-KEY ("Bisphosphonate-Associated Osteonecrosis of the Jaw" OR ("Osteonecrosis" AND "Jaw") OR (osteonecrosis AND jaw)) AND TITLE-ABS-KEY ("Therapeutics"))
Web of Science (July 25, 2022)	"Diphosphonates" OR "Alendronate" OR "Etidronic Acid" OR "ibandronic acid" OR "pamidronate" OR "zoledronic acid" OR "risedronic acid" OR "denosumab" (All Fields) AND "Bisphosphonate-Associated Osteonecrosis of the Jaw" OR ("Osteonecrosis" AND "Jaw") OR (osteonecrosis AND jaw) (All Fields) AND "Therapeutics" (All Fields)

2.4 Study selection

The study selection process comprised two phases. During Phase 1, two authors (MJAA and MOC DL) independently screened the titles and abstracts of all references using Rayyan software [22]. They identified studies that fulfilled the inclusion criteria and excluded those that did not. Whenever necessary, the third author (R.A.M.) was consulted to reach a final decision regarding the inclusion or exclusion of a study. In Phase 2, the same two authors independently evaluated the full-text articles to verify data on medication-related osteonecrosis of the jaw, applying the same selection criteria. All three authors critically assessed the reference lists of all included articles, and any additional articles meeting the inclusion criteria were incorporated into the selection analysis. Any discrepancies during either phase were resolved through discussion and consensus among the three authors. The final selection was based on including full-text articles that satisfied the predetermined criteria.

2.5 Data collection

The relevant information from the selected articles was gathered by the first author (MJAA) and second author (MOC DL). The accuracy of the collected data was then verified through cross-checking by the third author (R.A.M.). Any discrepancies were addressed through discussion and consensus among the three authors. Additionally, experts were consulted when necessary to assist in making final decisions. In cases where the required data could not be obtained

from the selected articles, efforts were made to establish contact with the corresponding authors to acquire the missing information.

2.6 Risk of Bias Within Studies

To assess the risk of bias, two authors (MJAA and MOCDL) independently employed the Joanna Briggs Institute's Critical Appraisal Checklist for Studies Reporting Prevalence Data and the Critical Appraisal Checklist for Case Reports [23]. In instances of disagreement, the third author (R.A.M.) was consulted to facilitate consensus. Before conducting the critical appraisal assessments, all authors engaged in discussions and made decisions regarding the scoring criteria. Studies were categorized as having a high risk of bias if they received a "yes" score of up to 49%, moderate between 50% and 69%, and low above 70%.

3. RESULTS

3.1 Study Selection and Characteristics

The study was conducted in two phases. Initially, a total of 3,253 records were gathered from multiple databases. After removing duplicates, the remaining 2,815 records underwent screening based on their titles and abstracts. Subsequently, 118 articles were selected for the second phase, which involved thoroughly reading the full texts. Applying the predetermined eligibility criteria, 89 studies were excluded, resulting in 30 studies that were eligible for synthesis (Figure 1). Among these 30 studies, 23 were cohort studies, and 7 were case-control studies. The entire process is visually depicted in Figure 2 through a flowchart.

First authors, year [reference]	Country of study	Study design	Sample size (case/control)	Age (years)		Sex		Site of injury	Mandible /Maxilla	Both mandible and maxilla	Evolution time (months)
				Case	control	case (M/F)	Control (M/F)				
Wutzi et al., 2008	Austria	Cohort	58	68.3 (±10.7; 32-92.2)	-	20/38	-	jaw	12/36	10	29.6-41.5
Scoletta et al., 2010	Italy	Cohort	20	71.3 (±9.86)	-	06/14	-	jaw	10/23	4	42.95 (±32.16)
Scoletta et al., 2010	Italy	Cohort	37	68 (±12.9)	-	11/26	-	jaw	10/23	4	25.5 (6-196)
Atalay et al., 2011	Peru	Case Control	10/10	55.4 (39-68)	-	07/13	-	jaw	9/11	-	32
Eckardt et al., 2011	Germany	Cohort	142	62 (38-94)	-	47/95	-	jaw	32/39	21	37 (5-130)
Freiberger et al., 2012	USA	Case-Control	25/21	66.1	66.3	13/12	06/15	jaw	-	-	3-24
Corviello et al., 2012	Italy	Case-Control	04/03	75.57	-	02/05	-	jaw	5/0	1	0.5
Graziani et al., 2012	Italy	Cohort	347	67 (±11; 34-92)	-	117/230	-	jaw	-	-	6
O'Ryan et al., 2012	USA	Cohort	30	77 (54-89)	-	04/26	-	jaw	6/11	-	52.8 (22.8-79.2)
Assaf et al., 2013	Germany	Cohort	20	74 (±6.4)	-	9/11	-	jaw	5/12	3	11 (±33; 9-84)
Lerman et al., 2013	USA	Cohort	120	63 (39-91)	-	60/60	-	jaw	85/21	14	36 (0-126)
Kim et al., 2016	South Korea	Cohort	325	75 (±10.0)	-	11/314	-	jaw	239/72	14	48
Otto et al., 2016	Germany	Cohort	54	71.4 (±9.2)	-	22/32	-	jaw	40/25	-	46.3 (±31.8)
Pichardo et al., 2016	Netherlands	Cohort	74	67.9 (26-91)	-	12/62	-	jaw	58/11	5	0.46-0.7
Blus et al., 2017	Italy	Cohort	18	69.1 (±8.3; 59-87)	-	13/05	-	jaw	4/14	-	30.4 (±41.9; 9-121)
Coropciuc et al., 2017	Belgium	Cohort	79	38 - 90	-	39/40	-	jaw	72/37	-	-
Jung et al., 2017	South Korea	Case-Control	07/10	75.11 (±8.11; 59-86)	-	01/16	-	jaw	17/0	-	3-4
Mauceri et al., 2017	Italy	Cohort	10	75.2 (±5.94)	-	03/07	-	jaw	01/09	-	12
Calvani et al., 2018	Italy	Case-Control	13/13	55-71	-	05/21	-	jaw	-	-	24-60
Hadaya et al., 2018	USA	Cohort	106	71.7	-	31/75	-	jaw	0/43	-	24
Nisi et al., 2018	Italy	Cohort	53	71.9 (±10.2; 41-87)	-	0/53	-	jaw	12/39	2	0.46
Ristow et al., 2018	Germany	Cohort	75	68.2 (±9.8)	-	33/42	-	jaw	68/24	-	44.5 (±34.0; 180)
El-Rabbany et al., 2019	Canada	Cohort	78	80.5 (71.8-87.5)	-	14/64	-	jaw	47/25	-	13
Giovannacci et al., 2019	Italy	Cohort	8	75.75 (62-85)	-	02/06	-	jaw	7/0	1	-
Petrovic et al., 2019	Serbia	Cohort	32	59 (±11.8)	-	11/21	-	jaw	9/23	1	108
Giudice et al., 2020	Italy	Cohort	129	71.2 (±12.7)	-	39/90	-	jaw	-	-	-
Sim et al., 2020	Australia	Case-Control	15/19	64 (59-71)	64 (5874)	18/16	-	jaw	-	-	-
Tenore et al., 2020	Italy	Case-Control	21/13	58.09	-	08/26	-	jaw	14/12	-	0.17-1
Varoni et al., 2021	Italy	Cohort	35	73.46 (±9.29; 51-93)	-	11/24	-	jaw	12/24	-	17-60
Blatt et al., 2022	Switzerland	Cohort	45	71.5 (±8.6)	-	18/27	-	jaws	-	-	0.17-1

Fig. 1. The main characteristics of the selected articles focus on diagnosing medication-related osteonecrosis of the jaw.

O'Rayan et al., 2012	Y	Y	N	Y	Y	Y	Y	Y	Y	88.9 / L
Coviello et al., 2012	Y	Y	N	Y	Y	U	U	N	Y	55.6 / M
Assaf et al., 2013	Y	Y	N	Y	Y	Y	Y	N	Y	77.8 / L
Kim et al., 2016	Y	Y	Y	Y	Y	Y	U	Y	Y	88.9 / L
Otto et al., 2016	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Pichardo et al., 2016	Y	Y	Y	Y	Y	Y	Y	U	Y	88.9 / L
Jung et al., 2017	Y	Y	N	Y	Y	Y	Y	Y	Y	88.9 / L
Mauceri et al., 2017	Y	Y	N	Y	Y	Y	U	Y	Y	77.8 / L
Blus et al., 2017	Y	Y	N	Y	N	Y	Y	N	N	55.6 / M
Coropciuc et al., 2017	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Hadaya et al., 2018	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Ristow et al., 2018	Y	Y	Y	Y	Y	Y	Y	Y	N	88.9 / L
Nisi et al., 2018	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Calvani et al., 2018	Y	N	N	U	Y	Y	Y	N	Y	55.6 / M
Giovannaci et al., 2019	Y	Y	N	Y	Y	Y	Y	N	N	66.7 / M
El – Rabbany et al., 2019	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Petrovic et al., 2019	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Giudice et al., 2020	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Tenore et al., 2020	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Sim et al., 2020	Y	Y	Y	Y	N	Y	Y	Y	Y	88.8 / L
Varoni et al., 2021	Y	Y	Y	Y	Y	Y	Y	Y	Y	100.0 / L
Blat et al., 2022	Y	Y	Y	Y	Y	Y	Y	Y	N	88.8 / L

Q1. Was the sample frame appropriate to address the target population? Q2. Were study participants sampled in an appropriate way? Q3. Was the sample size adequate? Q4. Were the study subjects and the setting described in detail? Q5. Was the data analysis conducted with sufficient coverage of the identified sample? Q6. Were valid methods used for the identification of the condition? Q7. Was the condition measured in a standard, reliable way for all participants? Q8. Was there appropriate statistical analysis? Q9. Was the response rate adequate, and if not, was the low response rate managed appropriately?
Y - Yes; N - No; U - Unclear; NA - Not applicable; H - High, M - Moderate; L - Low.

3.3 Synthesis of Studies

The systematic review comprised a study population of 2,079 patients. The sex distribution consisted of 1,495 (71.9%) women and 584 (28.1%) men, with an average age of 69.52 (± 9.67 ; range: 26-94) years. Regarding the location of the condition, 905 (43.5%) cases were observed in the mandible, 403 (19.4%) in the maxilla, 80 (3.8%) in both areas, and the location was not reported in 691 (33.2%) cases. The average duration of disease progression was 31.68 months (± 34.57 ; range: 0.17-196) (Figure 1).

The most frequently reported medical histories in the study population were breast cancer (n=344; 20%), multiple myeloma (n=330; 19.19%), hypertension (n=213; 12.38%), osteoporosis (n=191; 11.1%), and diabetes (n=161; 9.36%). These medical conditions were commonly associated with the use of specific medications, including Zoledronate [n=772, duration of 452.04 months (± 27.41 ; range: 12-102)], Alendronate [n=650, duration of 104.4 months (± 60.16 ; range: 6-180)], Pamidronate [n=121, duration of 20.74 months (± 4.94 ; range: 6-96)], and Denosumab [n=107, duration of 15 months (± 7.94)].

During the initial consultation, the stages of medication-related osteonecrosis of the jaw (MRONJ) were classified as follows: stage Osa (n=1; 0.09%), stage Oss (n=14; 1.31%), stage I (n=361; 33.68%), stage II (n=543; 50.65%), and stage III (n=153; 14.27%). In terms of treatment, the most commonly employed clinical therapies included conservative surgical management (n=578; 27.92%), local debridement (n=281; 13.57%), conservative treatment (n=232; 11.21%), sequestrectomy (n=224; 10.82%), surgical resection with auto-fluorescence guidance and low-level laser therapy (n=137; 6.62%), and a combination of chlorhexidine, antibiotics, analgesics, -and debridement (n=106; 5.12%).

Treatment complications were identified in 271 cases (13.03%). The most frequent complications were resistant or worsening clinical stage of osteonecrosis (n=184, 67.9%), followed by incomplete mucosal healing or dehiscence (n=47; 17.34%) and mental nerve injury (n=9; 3.32%), among others.

4. DISCUSSION

In this systematic review, we meticulously assessed and synthesized the findings of 30 studies encompassing a diverse global perspective, shedding light on the epidemiological and clinical aspects of MRONJ. Our comprehensive analysis has provided valuable insights into the prevalence, risk factors, treatment modalities, and associated complications of MRONJ. The study selection process followed a rigorous methodology, involving two distinct phases. A total of 3,253 records were initially identified, which were then meticulously screened and narrowed down to 30 studies meeting our predetermined eligibility criteria. The synthesis of the included studies revealed a study population of 2,079

patients, women (71.9%) with an average age of 69.52 years. The locations of MRONJ were observed in the mandible (43.5%) and the maxilla (19.4%), reflecting the areas most affected. Notably, the average duration of disease progression was approximately 31.68 months, indicating the chronic nature of MRONJ.

In all studies except those by Lerman et al. [24] and Sim et al. [25], a consistently high prevalence of MRONJ was observed in women. Lerman et al. [24] reported an equal distribution of cases with 60 (50%) men and 60 (50%) women, while Sim et al. [25] showed 18 (52.9%) men and 16 (47.1%) women affected. The rate of age across the articles varied from 26 to 94 years, with the lowest mean age reported by Atalay et al. [26] of 55.4 years. Regarding the affected anatomical sites, the mandible was consistently identified as the most affected area in all articles, except for Atalay et al. [26], Eckardt et al. [27], Hadaya et al. [28], and Tenore et al. [29]. In Atalay et al. [28], 11 cases were found in the maxilla and 9 in the mandible. Eckardt et al. [27] reported 39 cases in the maxilla and 32 in the mandible. Hadaya et al. [28] exclusively observed cases in the maxilla (n=43), and Tenore et al. [29] found 14 cases in the maxilla and 12 in the mandible. A recent study by Kuehn et al. [30] aimed to investigate the localizing factors that render the jawbone uniquely vulnerable to osteonecrosis with long-term anti-resorptive therapy. They noted that among patients receiving anti-resorptive medications, osteonecrosis was never reported in long bones. In Zoledronate-treated rats, traumatic bone exposure healed normally in areas where localized drug accumulation was like that in the jawbone [31]. On the contrary, Zoledronate treatment led to anabolic changes in cortical and trabecular areas in long bones, indicating that bone formation and remodeling remained active at normal levels in these regions. Additionally, levels of Wnt-3a and RANKL were significantly reduced in the jawbones of Zoledronate-treated animals, while iliac and tibial bones exhibited significantly increased levels [32]. Numerous factors have been explored as potential localizing factors, including dental trauma, particularly surgical extraction, periodontitis, impaired gingival healing, alterations in the oral bacteria biofilm profile, and compromised innate immune responses specific to the oral cavity [1,14,33-42].

Underlying medical conditions were frequently associated with MRONJ, with breast cancer (20%), multiple myeloma (19.19%), hypertension (12.38%), osteoporosis (11.1%), and diabetes (9.36%) being commonly reported. These conditions often correlate with the use of specific medications, such as Zoledronate, Alendronate, Pamidronate, and Denosumab, with varying durations of usage. Studies have reported a prevalence rate of 0.3% to 0.4% for MRONJ in breast cancer patients using antiangiogenic inhibitors like Bevacizumab [43]. However, the prevalence increased to a range of 0.9% to 2.4% when both bevacizumab and bisphosphonates were used in combination [43]. The prevalence associated with Zoledronate or Denosumab treatment in osteoporotic patients was reported as 0.017% and 0.3%, respectively [44,45].

The classification of MRONJ stages revealed varying severity, with stage II being the most prevalent (50.65%), followed by stage I (33.68%) and stage III (14.27%). The therapeutic landscape for MRONJ was diverse, with commonly employed approaches including conservative surgical management, local debridement, sequestrectomy, and other methods. It is worth noting that complications in treatment were observed in 13.03% of cases, with the most frequent being resistance or worsening clinical stages, incomplete mucosal healing, and mental nerve injury.

Treatment modalities should primarily involve conservative surgical procedures, with elective procedures avoided as they may compromise additional areas of exposed bone necrosis and exacerbate symptoms [32, 33]. Non-surgical conservative therapy in stage I of MRONJ yielded low healing rates, leading to the conclusion that early surgical interventions should be considered across all stages to prevent progression [34]. Several studies have compared conservative surgical and non-surgical protocols [46-75]. A variety of surgical treatment options have been proposed, showing promising results [46,48-50,52,53,55,57-60,62-64,66,68-71,74]. However, the absence of randomized controlled trials comparing nonsurgical and surgical treatment options has left the topic of surgical intervention in MRONJ still controversial [76]. Some less aggressive approaches have shown promise in reducing recurrences, such as autofluorescence/tetracycline-guided surgery, which enables the removal of all necrotic bone while preserving healthy bone [69]. The use of Piezoelectric in surgery is considered a less aggressive option and has been cited as a good choice in MRONJ cases [60,71]. To enhance surgical outcomes, various adjuvant therapies have been proposed in the literature, including parathyroid hormone (PTH), laser therapy, hyperbaric oxygen therapy, ozone therapy, and platelet-rich plasma (L-PRP/PRGF/PRP) [47,49,51,52,63,73,75]. These adjuvant therapies have shown faster and more comfortable postoperative healing. Additionally, studies have demonstrated the effectiveness of conservative management and antibiotics in MRONJ treatment [48,49,54,56,61,65,67,68,70,72]. It's important to note that while surgical therapies, with or without adjuvant protocols, appear effective in promoting MRONJ healing in all stages, they may not guarantee permanent lesion closure in the long term [76]. A cohort study found that wound dehiscence occurred in two patients one week after surgery, with one patient subsequently experiencing a painful area of bone exposure in the right jaw [77]. Studies have shown that MRONJ patients have an average healing duration of 3-6 months, with 50% experiencing incomplete healing, and 62.5% progressing to stage III [77]. Moreover, worsening of clinical stages, especially in MRONJ lesions, often occurs within the initial six months of treatment [78].

It is crucial to acknowledge the limitations of our study, including its retrospective nature and potential selection bias. Additionally, the variations in study designs, patient populations, and treatment protocols may contribute to heterogeneity in the results. Furthermore, while we assessed the risk of bias within individual studies, we did not account for potential biases introduced during the synthesis process.

5. CONCLUSION

In conclusion, our systematic review has shed light on essential aspects of MRONJ, including its prevalence, risk factors, treatment strategies, and associated complications. The global distribution of the selected studies highlights the widespread impact of this condition. As further research continues to refine our understanding of MRONJ, a collaborative and multidisciplinary approach involving dental, oncological, and maxillofacial specialists remains essential to optimize patient outcomes and enhance our management strategies.

REFERENCES

1. Marx RE. Pamidronate (Aredia) and zoledronate (Zometa) induced avascular necrosis of the jaws: a growing epidemic. *J Oral Maxillofac Surg.* 2003 Sep;61(9):1115-7.
2. Ruggiero SL, Dodson TB, Fantasia J, Goodday R, Aghaloo T, Mehrotra B, O’Ryan F. American Association of Oral and Maxillofacial Surgeons. American Association of Oral and Maxillofacial Surgeons position paper on medication-related osteonecrosis of the jaw--2014 update. *J Oral Maxillofac Surg.* 2014 Oct;72(10):1938-56.
3. Kuehn S, Scariot R, Elsalanty M. Medication-Related Osteonecrosis: Why the Jawbone? *Dent J (Basel).* 2023 Apr;11(5):109.
4. Ruggiero SL, Dodson TB, Aghaloo T, Carlson ER, Ward BB, Kademani D. American Association of Oral and Maxillofacial Surgeons' Position Paper on Medication-Related Osteonecrosis of the Jaws-2022 Update. *J Oral Maxillofac Surg.* 2022 May;80(5):920-943.
5. Gossiel F, Paggiosi MA, Naylor KE, McCloskey EV, Walsh J, Peel N, Eastell R. The effect of bisphosphonates on bone turnover and bone balance in postmenopausal women with osteoporosis: The T-score bone marker approach in the TRIO study. *Bone.* 2020 Feb; 131:115158.
6. Black DM, Cummings SR, Karpf DB, Cauley JA, Thompson DE, Nevitt MC, Bauer DC, Genant HK, Haskell WL, Marcus R, Ott SM, Torner JC, Quandt SA, Reiss TF, Ensrud KE. Randomised trial of effect of alendronate on risk of fracture in women with existing vertebral fractures. Fracture Intervention Trial Research Group. *Lancet.* 1996 Dec 7;348(9041):1535-41.
7. Black DM, Delmas PD, Eastell R, Reid IR, Boonen S, Cauley JA, Cosman F, Lakatos P, Leung PC, Man Z, Mautalen C, Mesenbrink P, Hu H, Caminis J, Tong K, Rosario-Jansen T, Krasnow J, Hue TF, Sellmeyer D, Eriksen EF, Cummings SR. HORIZON Pivotal Fracture Trial. Once yearly zoledronic acid for treatment of postmenopausal osteoporosis. *N Engl J Med.* 2007 May 3;356(18):1809-22.
8. Cummings SR, San Martin J, McClung MR, Siris ES, Eastell R, Reid IR, Delmas P, Zoog HB, Austin M, Wang A, Kutilek S, Adami S, Zanchetta J, Libanati C, Siddhanti S, Christiansen C. FREEDOM Trial. Denosumab for prevention of fractures in postmenopausal women with osteoporosis. *N Engl J Med.* 2009 Aug 20;361(8):756-65.
9. Stockmann P, Hinkmann FM, Lell MM, Fenner M, Vairaktaris E, Neukam FW, Nkenke E. Panoramic radiograph, computed tomography, or magnetic resonance imaging. Which imaging technique should be preferred in bisphosphonate-associated osteonecrosis of the jaw? A prospective clinical study. *Clin Oral Investig.* 2010 Jun;14(3):311-7.
10. Otto S, Pautke C, Van den Wyngaert T, Niepel D, Schioldt M. Medication-related osteonecrosis of the jaw: Prevention, diagnosis, and management in patients with cancer and bone metastases. *Cancer Treat. Rev.* 2018; 69:177-187.
11. Florenzano P, Pan KS, Brown SM, Paul SM, Kushner H, Guthrie LC, de Castro LF, Collins MT, Boyce AM. Age-Related Changes and Effects of Bisphosphonates on Bone Turnover and Disease Progression in Fibrous Dysplasia of Bone. *J Bone Miner Res.* 2019 Apr;34(4):653-660.
12. Ruggiero SL, Dodson TB, Assael LA, Landesberg R, Marx RE, Mehrotra B. Task Force on Bisphosphonate-Related Osteonecrosis of the Jaws, American Association of Oral and Maxillofacial Surgeons. American Association of Oral and Maxillofacial Surgeons position paper on bisphosphonate-related osteonecrosis of the jaw - 2009 update. *Aust Endod J.* 2009 Dec;35(3):119-30.
13. Campisi G, Mauceri R, Bertoldo F, Bettini G, Biasotto M, Colella G, Consolo U, Di Fede O, Favia G, Fusco V, Gabriele M, Lo Casto A, Lo Muzio L, Marciànò A, Mascitti M, Meleti M, Mignogna MD, Oteri G, Panzarella V, Romeo U, Santarelli A, Vescovi P, Marchetti C, Bedogni A. Medication-Related Osteonecrosis of Jaws (MRONJ) Prevention and Diagnosis: Italian Consensus Update 2020. *Int J Environ Res Public Health.* 2020 Aug 18;17(16):5998.

14. Khan AA, Morrison A, Hanley DA, Felsenberg D, McCauley LK, O’Ryan F, Reid IR, Ruggiero SL, Taguchi A, Tetradis S, Watts NB, Brandi ML, Peters E, Guise T, Eastell R, Cheung AM, Morin SN, Masri B, Cooper C, Morgan SL, Obermayer-Pietsch B, Langdahl BL, Al Dabagh R, Davison KS, Kendler DL, Sándor GK, Josse RG, Bhandari M, El Rabbany M, Pierroz DD, Sulimani R, Saunders DP, Brown JP, Compston J. International Task Force on Osteonecrosis of the Jaw. Diagnosis and management of osteonecrosis of the jaw: a systematic review and international consensus. *J Bone Miner Res.* 2015 Jan;30(1):3-23.
15. Japanese Allied Committee on Osteonecrosis of the Jaw; Yoneda T, Hagino H, Sugimoto T, Ohta H, Takahashi S, Soen S, Taguchi A, Nagata T, Urade M, Shibahara T, Toyosawa S. Antiresorptive agent-related osteonecrosis of the jaw: Position Paper 2017 of the Japanese Allied Committee on Osteonecrosis of the Jaw. *J Bone Miner Metab.* 2017 Jan;35(1):6-19.
16. Aparecida Cariolatto F, Carelli J, de Campos Moreira T, Pietrobon R, Rodrigues C, Bonilauri Ferreira AP. Recommendations for the Prevention of Bisphosphonate-Related Osteonecrosis of the Jaw: A Systematic Review. *J Evid Based Dent Pract.* 2018 Jun;18(2):142-152.
17. Beth-Tasdogan NH, Mayer B, Hussein H, Zolk O. Interventions for managing medication-related osteonecrosis of the jaw. *Cochrane Database Syst Rev.* 2017 Oct 6;10(10):CD012432.
18. Mücke T, Deppe H, Hein J, Wolff KD, Mitchell DA, Kesting MR, Retz M, Gschwend JE, Thalgott M. Prevention of bisphosphonate-related osteonecrosis of the jaws in patients with prostate cancer treated with zoledronic acid - A prospective study over 6 years. *J Craniomaxillofac Surg.* 2016 Oct;44(10):1689-1693.
19. Smidt-Hansen T, Folkmar TB, Fode K, Agerbaek M, Donskov F. Combination of zoledronic Acid and targeted therapy is active but may induce osteonecrosis of the jaw in patients with metastatic renal cell carcinoma. *J Oral Maxillofac Surg.* 2013 Sep;71(9):1532-40.
20. Guarneri V, Miles D, Robert N, Diéras V, Glaspy J, Smith I, Thomssen C, Biganzoli L, Taran T, Conte P. Bevacizumab, and osteonecrosis of the jaw: incidence and association with bisphosphonate therapy in three large prospective trials in advanced breast cancer. *Breast Cancer Res Treat.* 2010 Jul;122(1):181-8.
21. Hajeri S, Alturkistany Y. Osteonecrose da mandíbula relacionada a medicamentos após a depuração dentária: Prevalência em um centro de oncologia. *Saudi Dent J.* 2022 Set;34(6):479-484.
22. Soares A L, Simon S, Gebrin L H, Nazário A C P, Lazaretti-Castro M. Prevalence and risk factors of medication-related osteonecrosis of the jaw in osteoporotic and breast cancer patients: a cross-sectional study. *Supp.Care Cancer.* 2020; 28: 2265-2271.
23. Hallmer F, Bjarnadottir O, GöTRICK B, Malmström P, Andersson G. Incidence of and risk factors for medication-related osteonecrosis of the jaw in women with breast cancer with bone metastasis: a population-based study. *Oral Surg. Oral. Med. Oral. Pathol. Oral. Radiol.* 2020; 130: 252–257.
24. Kalra S, Jain V. Dental complications and management of patients on bisphosphonate therapy: a review article. *J. Oral. Biol. Craniofac. Res.* 2013; 3: 25–30
25. Ruggiero S L, Dodson T B, Fantasia J, Goodday R, Aghaloo T, Mehrotra B, O’Ryan F. American Association of Oral and Maxillofacial Surgeons position paper on medication-related osteonecrosis of the jaw-2014 update. *J. Oral Maxillofac. Surg.* 2014; 72: 1938-1956.
26. Jeong HG, Hwang JJ, Lee JH, Kim YH, Na JY, Han SS. Risk factors of osteonecrosis of the jaw after tooth extraction in osteoporotic patients on oral bisphosphonates. *Imaging Sci Dent.* 2017 Mar;47(1):45-50.
27. Saad F, Brown J, Van Poznak C, Ibrahim T, Stemmer S, Stopeck A, Diel I, Takahashi S, Shore N, Henry D. Incidence, risk factors, and outcomes of osteonecrosis of the jaw: integrated analysis from three blinded active-controlled phase III trials in cancer patients with bone metastases. *Ann.Oncol.* 2012; 23: 1341-1347.
28. Hallmer F, Andersson G, Götrick B, Warfvinge G, Anderud J, Bjørnland T. Prevalence, initiating factor, and treatment outcome of medication-related osteonecrosis of the jaw-a 4-year prospective study. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2018 Dec;126(6):477-485.
29. Aljohani S, Fliefel R, Ihbe J, Kühnisch J, Ehrenfeld M, Otto S. What is the effect of anti-resorptive drugs (ARDs) on the development of medication-related osteonecrosis of the jaw (MRONJ) in osteoporosis patients: A systematic review. *J Craniomaxillofac Surg.* 2017 Sep;45(9):1493-1502.
30. Henry D, Vadhan-Raj S, Hirsh V, von Moos R, Hungria V, Costa L, Woll PJ, Scagliotti G, Smith G, Feng A, Jun S, Dansey R, Yeh H. Delaying skeletal-related events in a randomized phase 3 study of Denosumab versus zoledronic acid in patients with advanced cancer: an analysis of data from patients with solid tumors. *Support Care Cancer.* 2014 Mar;22(3):679-87.
31. Henedige AA, Jayasinghe J, Khajeh J, Macfarlane TV Systematic review on the incidence of bisphosphonate related osteonecrosis of the jaw in children diagnosed with osteogenesis imperfecta. *J Oral Maxillofac Res.* 2013; 4: e1
32. Ikeda T, Kuraguchi J, Kogashiwa Y, Yokoi H, Satomi T, Kohno N. Successful treatment of bisphosphonate-related osteonecrosis of the jaw (BRONJ) patients with cefefloxacin: new strategies for treatment of BRONJ *Bone.* 2015; 73: 217-222.
33. Ruggiero SL, Kobn N. Disease stage and mode of therapy are important determinants of treatment outcome for medication-related osteonecrosis of the jaw. *J Oral axillofac Surg.* 2015; 73: 94-100.
34. Pelaz A, Junqueira L, Gallego L, Garcia-Consuegra L, Junqueira S, Gómez C. Alternative treatments for oral

- bisphosphonate-related osteonecrosis of the jaws: a pilot study comparing fibrin rich in growth factors and teriparatide. *Med Oral Patol Oral Dis.* 2014; 19: 320-326.
35. Campisi G, Mauceri R, Bertoldo F, Bettini G, Biasotto M, Colella G, Consolo U, Di Fede O, Favia G, Fusco V, Gabriele M, Lo Casto A, Lo Muzio L, Marcianò A, Mascitti M, Meleti M, Mignogna MD, Oteri G, Panzarella V, Romeo U, Santarelli A, Vescovi P, Marchetti C, Bedogni A. Medication-Related Osteonecrosis of Jaws (MRONJ) Prevention and Diagnosis: Italian Consensus Update 2020. *Int J Environ Res Public Health.* 2020 Aug 18;17(16):5998.
 36. Steinberg ME, Hayken GD, Steinberg DR. A quantitative system for staging avascular necrosis. *J Bone Joint Surg Br.* 1995; 40: 77:34.
 37. Fleisher KE, Welch G, Kottal S, Craig RG, Saxena D, Glickman RS. Predicting risk for bisphosphonate-related osteonecrosis of the jaws: CTX versus radiographic markers. *Oral Surg Oral Med Oral Pathol Oral RadiolEndod.* 2010 Oct;110(4):509-16.
 38. Hoefert S, Yuan A, Munz A, Grimm M, Elayouti A, Reinert S. Clinical course, and therapeutic outcomes of operatively and non-operatively managed patients with denosumab-related osteonecrosis of the jaw (DRONJ). *J Craniomaxillofac Surg.* 2017; 45: 570-578
 39. de Molon RS, Hsu C, Bezouglaia O, Dry SM, Pirih FQ, Soundia A, Cunha FQ, Cirelli JA, Aghaloo TL, Tetradis S. Rheumatoid Arthritis Exacerbates the Severity of Osteonecrosis of the Jaws (ONJ) in Mice. A Randomized, Prospective, Controlled Animal Study. *J Bone Miner Res.* 2016 Aug;31(8):1596-607.
 40. Peters E, Lovas GL, Wysocki GP. Lingual mandibular sequestration and ulceration. *Oral Surg Oral Med Oral Pathol.* 1993; 75:739.
 41. Koorbush GF, Fotos P, Goll KT. Retrospective assessment of osteomyelitis. Etiology, demographics, risk factors, and management in 35 cases. *Oral Surg Oral Med Oral Pathol.* 1992; 74:149- 58.
 42. Maines E, Monti E, Doro F, Morandi G, Cavarzere P, Antoniazzi F. Children, and adolescents treated with neridronate for osteogenesis imperfecta show no evidence of any osteonecrosis of the jaw. *J Bone Miner Metab.* 2012 Jul;30(4):434-8.
 43. Gallego L, Junquera L, Pelaz A, Hernando J, Megías J. The use of pedicled buccal fat pad combined with sequestrectomy in bisphosphonate-related osteonecrosis of the maxilla. *Med Oral Patol Oral Cir Bucal.* 2012; 17: 236-241.
 44. Mücke T, Koerdt S, Jung M, Mitchell DA, Wolff KD, Kesting MR, Loeffelbein DJ. The role of mylohyoid flap in the treatment of bisphosphonate-related osteonecrosis of the jaws. *J Craniomaxillofac Surg.* 2016; 4.
 45. Nocini PF, Saia G, Bettini G, Ragazzo M, Blandamura S, Chiarini L, Bedogni A. Vascularized fibula flap reconstruction of the mandible in bisphosphonate-related osteonecrosis. *Eur J Surg Oncol.* 2009; 35: 373-379.
 46. Wutzl A, Biedermann E, Wanschitz F, Seemann R, Klug C, Baumann A, Watzinger F, Schicho K, Ewers R, Millesi G. Treatment results of bisphosphonate-related osteonecrosis of the jaws. *Head Neck.* 2008 Sep;30(9):1224-30.
 47. Scoletta M, Arduino PG, Reggio L, Dalmaso P, Mozzati M. Effect of low-level laser irradiation on bisphosphonate-induced osteonecrosis of the jaws: preliminary results of a prospective study. *Photomed Laser Surg.* 2010 Apr;28(2):179-84.
 48. Scoletta M, Arduino PG, Dalmaso P, Broccoletti R, Mozzati M. Treatment outcomes in patients with bisphosphonate-related osteonecrosis of the jaws: a prospective study. *Oral Surg Oral Med Oral Pathol Oral RadiolEndod.* 2010 Jul;110(1):46-53.
 49. Atalay B, Yalcin S, Emes Y, Aktas I, Aybar B, Issever H, Mandel NM, Cetin O, Oncu B. Bisphosphonate-related osteonecrosis: laser-assisted surgical treatment or conventional surgery? *Lasers Med Sci.* 2011 Nov;26(6):815-23.
 50. Eckardt AM, Lemound J, Lindhorst D, Rana M, Gellrich NC. Surgical management of bisphosphonate-related osteonecrosis of the jaw in oncologic patients: a challenging problem. *Anticancer Res.* 2011 Jun;31(6):2313-8.
 51. Freiberger JJ, Padilla-Burgos R, McGraw T, Suliman HB, Kraft KH, Stolp BW, Moon RE, Piantadosi CA. What is the role of hyperbaric oxygen in the management of bisphosphonate-related osteonecrosis of the jaw: a randomized controlled trial of hyperbaric oxygen as an adjunct to surgery and antibiotics. *J Oral Maxillofac Surg.* 2012 Jul;70(7):1573-83.
 52. Coviello V, Peluso F, Dehkhargani SZ, Verdugo F, Raffaelli L, Manicone PF, D' Addona A. Platelet-rich plasma improves wound healing in multiple myeloma bisphosphonate-associated osteonecrosis of the jaw patients. *J Biol Regul Homeost Agents.* 2012 Jan-Mar;26(1):151-5.
 53. Graziani F, Vescovi P, Campisi G, Favia G, Gabriele M, Gaeta GM, Gennai S, Goia F, Miccoli M, Peluso F, Scoletta M, Solazzo L, Colella G. Resective surgical approach shows a high performance in the management of advanced cases of bisphosphonate-related osteonecrosis of the jaws: a retrospective survey of 347 cases. *J Oral Maxillofac Surg.* 2012 Nov;70(11):2501-7.
 54. O'Ryan FS, Lo JC. Bisphosphonate-related osteonecrosis of the jaw in patients with oral bisphosphonate exposure: clinical course and outcomes. *J Oral Maxillofac Surg.* 2012 Aug;70(8):1844-53.
 55. Assaf AT, Zrnc TA, Riecke B, Wikner J, Zustin J, Friedrich RE, Heiland M, Smeets R, Gröbe A. Intraoperative efficiency of fluorescence imaging by Visually Enhanced Lesion Scope (VELscope) in patients with bisphosphonate related osteonecrosis of the jaw (BRONJ). *J Craniomaxillofac Surg.* 2014 Jul;42(5): e157-64.
 56. Lerman MA, Xie W, Treister NS, Richardson PG, Weller EA, Woo SB. Conservative management of bisphosphonate-

- related osteonecrosis of the jaws: staging and treatment outcomes. *Oral Oncol.* 2013 Sep;49(9):977-983.
57. Kim HY, Lee SJ, Kim SM, Myoung H, Hwang SJ, Choi JY, Lee JH, Choung PH, Kim MJ, Seo BM. Extensive Surgical Procedures Result in Better Treatment Outcomes for Bisphosphonate-Related Osteonecrosis of the Jaw in Patients with Osteoporosis. *J Oral Maxillofac Surg.* 2016 Jul;75(7):1404-1413.
 58. Otto S, Ristow O, Pache C, Troeltzsch M, Fliefel R, Ehrenfeld M, Pautke C. Fluorescence-guided surgery for the treatment of medication-related osteonecrosis of the jaw: A prospective cohort study. *J Craniomaxillofac Surg.* 2016 Aug;44(8):1073-80.
 59. Pichardo SE, Kuijpers SC, van Merkesteyn JP. Bisphosphonate-related osteonecrosis of the jaws: Cohort study of surgical treatment results in seventy-four stage II/III patients. *J Craniomaxillofac Surg.* 2016 Sep;44(9):1216-20.
 60. Blus C, Giannelli G, Szmukler-Moncler S, Orru G. Treatment of medication-related osteonecrosis of the jaws (MRONJ) with ultrasonic piezoelectric bone surgery. A case series of 20 treated sites. *Oral Maxillofac Surg.* 2017 Mar;21(1):41-48.
 61. Coropciuc RG, Grisar K, Aerden T, Schol M, Schoenaers J, Politis C. Medication-related osteonecrosis of the jaw in oncological patients with skeletal metastases: conservative treatment is effective up to stage 2. *Br J Oral Maxillofac Surg.* 2017 Oct;55(8):787-792.
 62. Jung J, Yoo HY, Kim GT, Lee JW, Lee YA, Kim DY, Kwon YD. Short-Term Teriparatide and Recombinant Human Bone Morphogenetic Protein-2 for Regenerative Approach to Medication-Related Osteonecrosis of the Jaw: A Preliminary Study. *J Bone Miner Res.* 2017 Dec;32(12):2445-2452.
 63. Mauceri R, Panzarella V, Maniscalco L, Bedogni A, Licata ME, Albanese A, Toia F, Cumbo E.M.G., Mazzola G, Di Fede O, Campisi G. Conservative Surgical Treatment of Bisphosphonate-Related Osteonecrosis of the Jaw with Er, Cr: YSGG Laser and Platelet-Rich Plasma: A Longitudinal Study. *Biomed Res Int.* 2018 Aug 19; 2018:3982540.
 64. Calvani F, Cutone A, Lepanto MS, Rosa L, Valentini V, Valenti P. Efficacy of bovine lactoferrin in the post-surgical treatment of patients suffering from bisphosphonate-related osteonecrosis of the jaws: an open-label study. *Biometals.* 2018 Jun;31(3):445-455.
 65. Hadaya D, Soundia A, Freymiller E, Grogan T, Elashoff D, Tetradis S, Aghaloo TL. Nonsurgical Management of Medication-Related Osteonecrosis of the Jaws Using Local Wound Care. *J Oral Maxillofac Surg.* 2018 Nov;76(11):2332-2339.
 66. Nisi M, Karapetsa D, Gennai S, Ramaglia L, Graziani F, Gabriele M. Conservative surgical treatment of medication related osteonecrosis of the jaw (MRONJ) lesions in patients affected by osteoporosis exposed to oral bisphosphonates: 24 months follow-up. *J Craniomaxillofac Surg.* 2018 Jul;46(7):1153-1158.
 67. Ristow O, Rückschloß T, Müller M, Berger M, Kargus S, Pautke C, Engel M, Hoffmann J, Freudlsperger C. Is the conservative non-surgical management of medication-related osteonecrosis of the jaw an appropriate treatment option for early stages? A long-term single-center cohort study. *J Craniomaxillofac Surg.* 2019 Mar;47(3):491-499.
 68. El-Rabbany M, Lam DK, Shah PS, Azarpazhooh A. Surgical Management of Medication-Related Osteonecrosis of the Jaw Is Associated with Improved Disease Resolution: A Retrospective Cohort Study. *J Oral Maxillofac Surg.* 2019 Sep;77(9):1816-1822.
 69. Giovannacci I, Vescovi P, Magnoni C, Corradi D, Corcione L, Lucchina AG, Mortellaro C, Nammour S, Meleti M. Auto-Fluorescence and Histopathologic Evaluation of Medication-Related Osteonecrosis of the Jaws: Perspectives for Treatment. *J Craniofac Surg.* 2019 Jun;30(4):1039-1043.
 70. Petrovic M, Jelovac DB, Antic S, Antunovic M, Lukic N, Sabani M, Mudrak J, Jezdic Z, Pucar A, Stefanovic A, Kuzmanovic C, Nikolic D, Konstantinovic V. Medication-Related Osteonecrosis of the Jaws: Two Center Retrospective Cohort Studies. *Biomed Res Int.* 2019 Mar 18; 2019:8345309.
 71. Giudice A, Barone S, Diodati F, Antonelli A, Nocini R, Cristofaro MG. Can Surgical Management Improve Resolution of Medication-Related Osteonecrosis of the Jaw at Early Stages? A Prospective Cohort Study. *J Oral Maxillofac Surg.* 2020 Nov;78(11):1986-1999.
 72. Sim IW, Borromeo GL, Tsao C, Hardiman R, Hofman MS, Papatziarnos Hjelle C, Siddique M, Cook G.J.R., Seymour JF, Ebeling PR. Teriparatide Promotes Bone Healing in Medication-Related Osteonecrosis of the Jaw: A Placebo-Controlled, Randomized Trial. *J Clin Oncol.* 2020 Sep 10;38(26):2971-2980.
 73. Tenore G, Zimbalatti A, Rocchetti F, Graniero F, Gaglioti D, Mohsen A, Caputo M, Lollobrigida M, Lamazza L, De Biase A, Barbato E, Romeo U. Management of Medication-Related Osteonecrosis of the Jaw (MRONJ) Using Leukocyte- and Platelet-Rich Fibrin (L-PRF) and Photobiomodulation: A Retrospective Study. *J Clin Med.* 2020 Oct 29;9(11):3505.
 74. Varoni EM, Lombardi N, Villa G, Pispero A, Sardella A, Lodi G. Conservative Management of Medication-Related Osteonecrosis of the Jaws (MRONJ): A Retrospective Cohort Study. *Antibiotics (Basel).* 2021 Feb 17;10(2):195.
 75. Blatt S, Krüger M, Kämmerer PW, Thiem D.G.E., Matheis P, Eisenbeiß AK, Wiltfang J, Al-Nawas B, Naujokat H. Non-Interventional Prospective Observational Study of Platelet Rich Fibrin as a Therapy Adjunctive in Patients with Medication-Related Osteonecrosis of the Jaw. *J Clin Med.* 2022 Jan 28;11(3):682.
 76. Goker F, Grecchi E, Grecchi F, Francetti L, Del Fabbro M. Treatment of medication-related osteonecrosis of the jaw (MRONJ). A systematic review. *Eur Rev Med Pharmacol Sci.* 2021 Mar;25(6):2662-2673.
 77. Odet S, Meyer C, Gaudet C, Weber E, Quenot J, Derruau S, Laurence S, Bompuy L, Girodon M, Chatelain B,

- Mauprivez C, Brenet E, Kerdjoudj H, Zwetyenga N, Marchetti P, Hatzfeld AS, Toubeau D, Pouthier F, Lafarge X, Redl H, Fenelon M, Fricain JC, Di Pietro R, Ledouble C, Gualdi T, Parmentier AL, Louvrier A, Gindraux F. Tips and Tricks and Clinical Outcome of Cryopreserved Human Amniotic Membrane Application for the Management of Medication-Related Osteonecrosis of the Jaw (MRONJ): A Pilot Study. *Front BioengBiotechnol.* 2022 Nov 03; 10:1058241
78. Coropciuc RG, Grisar K, Aerden T, Schol M, Schoenaers J, Politis C. Medication-related osteonecrosis of the jaw in oncological patients with skeletal metastases: conservative treatment is effective up to stage 2. *Br J Oral Maxillofac Surg.* 2017 Oct;55(8):787-792.

UNDER PEER REVIEW