

IMPACT OF 1 YEAR OF LARGE-SCALE ANTIRETROVIRAL DRUG TREATMENT WITH LOW THERAPEUTIC COVERAGE ON THE ANTHROPOMETRIC PROFILES IN THE RURAL DISTRICT HOSPITAL OF EASTERN NIGERIA.

Comment [d1]: Modification to title.

Abstract:

The study aimed at assessing the evolution of body mass index of (BMI) and treatment outcome of HIV adult-patients who were on antiretroviral therapy (ART) over a period of one year in a community district hospital with high prevalence of HIV attendance. One year retrospective follow-up study of the recruited patients was carried out using data collected from their treatment cards. One hundred and fifty-one patients were recruited for the study but eleven patients dropped as a result of clinical emergencies and breach of study protocol. Ninety-two (66%) were females and the rest adult males. Study patients were on four different WHO clinical stages at various numerical strengths. There were significant ($p \geq 0.05$) synchronized linear increment of BMI and positive treatment outcomes following treatment with antiretroviral therapy (ART).

Comment [d2]: Delete to space

Comment [d3]: Explain to HIV full form.

Comment [d4]: WHO stand for ?

BACKGROUND.

UNAIDS estimates that by mid 1998, over 30 million people were living with HIV/AIDS¹. More than 90% of these people live in developing countries, where majority of the infected population have no access to any kind of medicaments and discrimination also is very high. While international agencies and federal government of Nigeria supply both antiretroviral drugs and anti-parasitic drugs free of charge to the people living with the virus through health institutions, discrimination and poor coverage to hinterlands where the infection is more prevalent, pose a great challenge. The infection has continued to be a major global public health problem, having killed more than 35 million lives so far in 2017 while 940,000 people died of HIV-related causes globally².

Comment [d5]: First time explain to short form words.

However, with increasingly widespread use of potent combination antiretroviral therapy (ART) since 1996, substantial improved prognosis of people infected with HIV in sub-Saharan Africa has been observed. Administration of ART to people living with HIV and monitoring and evaluation of HIV/AIDS treatment programmes

critically depend on regular and complete patient follow-up. Individual treatment decisions can then be made in the light of clinical and laboratory results, and treatment response, complication and mortality rates can be accurately estimated at the programme level.

Comment [d6]: Mention to citation

Body weight of people living with HIV has been an important diagnostic and evaluation measure. Low body mass index (BMI) is recognized as one of the first criteria for the clinical definition of AIDS. The current WHO clinical staging of the infection also includes moderate unexplained weight loss (<10%) presumed or measured body weight while clinical stage 2 is unexplained severe weight loss (>10%) of body weight or BMI < 18.5kg/m². Clinical stages 3 & 4 are unexplained severe wasting syndrome which is used to define advanced HIV infection³.

Comment [d7]: Delete to extra space

The BMI of people living with HIV infection is an important predictor of ART outcome,⁴⁻⁶ including the prediction of the CD4 cells change⁷ and death⁹⁻¹⁰. A negative change in BMI of patients was also found to be an dependent predictor of drop-out from care or non-response to the first line of treatment regimen and therefore, may demand need to change to second line treatment regimen. However, the BMI of patients with HIV is dependent on gender and also affected by drugs used to treat opportunistic infections such as isoniazid, cotrimoxazole, CD4 cells count and socio-economic profile.

Comment [d8]: Need to space between isoniazid and cotrimaxazole drugs.

Using data from a district health institution designated for diagnosis and management of HIV infection, we examine some anthropometric profiles of patients who have consistently been on ART for 1 year as an important and easily calculable tool to predict positive treatment outcome of HIV patients on ART.

MATERIALS AND METHODS.

The correlation between gain in anthropometric status as a predictor of antiretroviral drugs (ARDs) efficiency was prospectively evaluated in all those who had been on ART over the period of March 13, 2012 to April 15, 2013 at Udi District Hospital in the Eastern geographical zone of Nigeria. This hospital is centrally situated and very proximate to Oji province where high prevalence of HIV infection had been reported. Screening and enrollment for the study began

March 14, 2012 after approval of the protocol by the management of ethics hospital committee.

The study entails participants being willing to present themselves for weight, height and haematocrit evaluations. All the clients underwent anthropometrics measurement carried by a trained medical laboratory assistant. Clients' weights were measured without shoes, using Seca scales, which had intervals of 0.5kg, height was measured to the nearest 0.1cm using a calibrated Scale consisting of a wooden platform with a scale and a sliding head piece. To reduce intra-individual error, weight and height were measured twice and the mean value was used for analysis. Weights for age; z-score was used to denote underweight as an overall indicator for malnutrition. Height for; z-score was used as indicator for stunting (chronic malnutrition). Weight for height; z-score was used as an indicator for wasting (acute malnutrition). The z-scores were calculated based on median values of the National Centre for Health Statistics(NCHS), Ref, population. Blood from finger prick was used to evaluate haemoglobin status spectrophotometrically using Drabkins reagents. As outlined in the study protocol, participants (clients) must be willing and able to give informed consent, to access the drugs regularly without missing any appointments as prescribed by the care givers for the 12 months and without relocating their residence until the participants were followed-up for the 12 months and clients who were highly cachexic or debilitated by the infection were not recruited initially to minimize drop outs.

RESULT.

A total of 140 adult patients aged between 22 and 58 years who were on ART treatment regimen were recruited in the study. Out of the number (140), 92 (66%) were female. At the beginning of the study, we started with 151 adult patients but 11 patients out of this number dropped out, following emergencies and non-adherence to the treatment regimen respectively. Greater number of the study population was self employed 57 (41%) while the least affected was unemployed populace (8%). Body weight and haemoglobin level of all the

Comment [d9]: Typing error

Comment [d10]: Typing error

Comment [d11]: Typing error,

patients who adhered strictly to the study protocol were measured on each visit for the 1 year study period apart from the one taken at the recruitment day.

Most of the patients had formal education of different levels 125 (89%). Seventy (50%) patients had WHO clinical stage II, 38 (27%) had WHO stage I while 32 (23%) had WHO clinical stage III and IV. There were significant synchronized linear increment of BMI and positive treatment outcomes following treatment with antiretroviral therapy (ART) . The baseline socio-demographic characteristic of the study patient is shown in table 1.

|

| _____ **Comment [d12]:** Delete extra space

Table 1: Baseline socio-demographic characteristics of adult patients with HIV on ART in Udi district hospital. Eastern Nigeria, January 2020- December 2020.

Comment [d13]: Modification to table alignment .

Variables	Frequency	Percentage
Age (years)		
22-31	35	25
32-41	62	44

42-51	27	19
>51	16	11
Sex		
Female	92	66
Male	48	34
Educational status		
Non formal education	15	11
Primary education	55	39
Secondary education	52	37
Tertiary education	18	13
Occupation		
Government employed	25	18
Private employed	32	23
Self employed	57	41
Unemployed	11	8
Student	15	11
WHO stages		
Stage I	38	27
Stage II	70	50
Stage III and IV	32	23

Functional status		
Working	125	89
Ambulatory	15	11
CPT given (960mg)		
Yes	95	68
No	45	32
INH		
Yes	10	7
No	130	93

Note: CPT – Cotrimoxazole prophylaxis therapy

ART – Antiretroviral therapy

INH – Isoniazid.

Table 2: Baseline characteristics of the study patients.

Comment [d14]: Modification to table alignment and font size.

Hb (range in g/dl)	Pre- treatment Frequency	Post treatment frequency	P-value
6 – 8	85	32	p> 0.05
9 – 11	50	83	
>11	5	25	”
Weight (kg)			”
30 – 39	39	18	
40 – 49	40	44	”
50 – 59	59	65	”
>59	8	13	”
LFT			
Bilirubin (mmol/l)			
Total < 0.5 – 1.0	139	140	P< 0.05
Conjugated <0.5	139	140	”
Total >1.0	1	0	”
Alanine transaminase – ALT(mmol/l) < 3- 15mol/l	140	140	”
Aspartate transaminase AST <3 -18	140	140	”
Urea (mmol/l)	140	140	”

Creatinine (mmol/l)	140	140	
CD4 cells <100- 198	140		”
>250		140	

Comment [d15]: Delete

DISCUSSION.

Clients presenting with autoimmune deficiency syndrome are at risk of developing severe anaemia, sarcoma, etc and dying. Oral ART treatment is therefore often recommended to ensure rapid antiretroviral bioavailability, and thus expedite reduction of the infecting retroviral biomass and therefore tune-up cluster of differentiation type 1V (CD4) cells as sub population of T-lymphocytes.

Comment [d16]: Mention to citation

This study carried out in district hospital attended by a community earlier reported to have high prevalence of HIV infection, showed direct correlation between BMI of patients on ART over time and positive treatment outcomes. There is a BMI on enrollment of patients studied and their BMI on treatment over time. Absolute compliance to prescribed treatment regimen was found to be a significant prediction of positive BMI changes. Gradual increment in BMI of study patients is attributed to efficiency of ART on HIV virus. However, the BMI of ambulatory and bedridden patients did not improve in synchrony with the duration of treatment received. This is attributed to their inability to move about which prevent them from accessing nutrition and balanced diet of their choice. This class of patient is susceptible to minor infections and osteoporosis because of institutionalization¹². More so, ART related metabolic changes cause hyperlipidanaemia, insulin resistance diabetes which enhances average weight gain¹³.

The BMI of patients at WHO stages 111 and 1V did not improve. This finding corroborates a multicentre study in a resource-poor settings which shows

individuals who had high clinical status (WHO stages 111 and 1V) had poorer weight gain when compared with weight change in patients at lower WHO stage

14.

The duration of ART treatment with addition of isoniazid prophylaxis therapy (IPT) and cotrimoxazole prophylaxis therapy (CPT) correlate positively with resolution of opportunistic infections with increase in CD4 cell counts. This is obvious in wellbeing of patients and increase in their appetite. These multivariable factors result in significant improvement in the haemoglobin values of the study patients when their baseline haemoglobin values was compared with their haemoglobin values over treatment period. Also it was found out that simultaneous administration of IPT and CPT with ART regimen optimizes efficiency and effectiveness of ART and expedite patients' wellbeing and reduce opportunistic infecting biomass.

Reference:

1UNAIDS/WHO. Report on the Global HIV/AIDS epidemic: June 1998, Geneva;:World Health Organization.

2 World Health Organization. Interim who clinical staging of HIV/AIDS and HIV/AIDS case definitions for surveillance, African region,2005.

3 World Health Organization.Hiv/Aids key facts,2018.
Available:<http://www.who.int/news-room/fact-sheets/detail/hiv-aids>.

4 Tsegaye AT, Wubshet M, Awoke T, et al. Predictors of treatment failure on second-line antiretroviral therapy among adults in Northwest Ethiopia: a multicentre retrospective follow-up study. *BMJ Open* 2016;6:e012537.

5.Seyoum A, Ndiovu P, Zewotir T.Quasi-Poisson versus negative binomial regression models in identifying factors affecting initial CD4 cell count change due to antiretroviral therapy administered to HIV-positive adults in north-west Ethiopia (Amhara region). *AIDS Res Ther* 2016;13:36.

Comment [d17]: Refer to APA Style.

Comment [d18]: Old reference , please mention within 5 to 10 years.

6. Patel D, Desai M, Shah AN, et al. Early outcome of second-line antiretroviral therapy in treatment-experienced human immunodeficiency virus positive patients. *Perspect Clin Res* 2013;4:215-20.
- 7 Dalhatu I, Onotu D, Odafe S, et al. Outcomes of Nigeria's HIV/AIDS treatment program for patients initiated on antiretroviral treatment between 2004-2012. *PLoS One* 2016,201611:e0165528.
- 8 Koethe JR, Jenkins CA, Lau B, et al. Body mass index and early CD4 T-cell recovery among adults initiating antiretroviral therapy in North America, 1998-2010. *HIV Med* 2015;16:572-7.
9. Chakravarty J, Sundar S, Chourasia A, et al. Outcome of patients on second-line antiretroviral therapy under programmatic condition in India, *BMC Infect Dis* 2015;15:517.
 - 10 Thao VP, Quang VM, Wolbers M, et al. Second-line HIV therapy outcomes and determinants of mortality at the largest HIV referral center in southern Vietnam. *Medicine*, 2015;94:e1715.
- 11 Nieman DC. Exercise, infection and immunity. *Int J Sports Med* 1994; 15 Suppl 3: S13`-S141.
12. Swain PK, Grover G. Determination of predictors associated with HIV/AIDS patients using accelerated failure time model for interval censored survival data. 2016;6:12-19.
13. Montessori V, Press N, Harris M, et al Adverse effects of antiretroviral therapy for HIV infection. *CMAJ: Canadian Medical Association Journal* 2004;170:229-38.
14. Hulsin t Veld D, Balestre E, Buyze J, et al. Determinants of weight evolution among HIV-positive patients initiating antiretroviral treatment in low-resource settings. *J Acquir Immune Defic Syndr*, 2015;70:146-54.