

AN OVERVIEW OF THE ADVANCED MRI TECHNIQUES USED FOR THE EARLY DETECTION OF STROKE AND OTHER CEREBROVASCULAR DISEASES IN EMERGENCY SITUATIONS

ABSTRACT

Stroke or cerebrovascular accidents pose a great threat to the health and mortality of individuals. Although strokes are now seen to be reversible, if they are not caught and intervened with, then they could lead to deadly consequences in the long term. In the modern world today, there are several interventions, therapeutic measures, and diagnostic facilities that help detect early changes within the brain in no time. Similarly, several advanced MRI techniques play a great role in the early detection of stroke and other cerebrovascular diseases in emergencies. These techniques include diffusion-weighted imaging (DWI), which allows for the visualization of acute ischemic lesions within minutes of symptom onset, aiding in rapid diagnosis. Perfusion-weighted imaging (PWI) provides valuable insights into tissue perfusion and viability, helping to identify areas of ischemia and guide treatment decisions. Susceptibility-weighted imaging (SWI) enhances the detection of hemorrhagic strokes and microvascular abnormalities, which is highly important for prompt intervention. Additionally, magnetic resonance angiography (MRA) offers detailed visualization of the cerebrovascular anatomy, aiding in identifying occlusions or stenosis contributing to cerebrovascular diseases. By integrating these advanced MRI techniques into emergency protocols, clinicians can swiftly and accurately diagnose cerebrovascular emergencies, enabling timely interventions and improving patient outcomes. This article will

discuss advanced MRI techniques utilized in emergency scenarios for early detection of stroke and other cerebrovascular diseases. Rapid and accurate diagnosis is crucial for timely intervention and improved patient outcomes in such critical scenarios. Conventional MRI protocols often face limitations in providing timely information due to lengthy acquisition times and limited sensitivity to acute pathologies. Furthermore, ongoing advancements in MRI technology, including ultrafast sequences and artificial intelligence-based image analysis, hold promise for further improving the efficiency and accuracy of emergency cerebrovascular imaging.

Keywords: Magnetic resonance imaging, MRI, stroke, cerebrovascular accidents, early diagnosis, advanced MRI techniques.

INTRODUCTION

Stroke stands as the second leading cause of death globally and a significant driver of disability, imposing substantial economic strains. It manifests in two primary forms: ischemic strokes, triggered by blood vessel blockages limiting brain blood supply, and hemorrhagic strokes, resulting from vessel ruptures and intracranial bleeding.(1)

Defined comprehensively by the American Heart Association/The American Stroke Association (AHA/ASA), stroke is essentially defined as, “an acute episode of focal neurological dysfunction persisting for over 24 hours”.(2)

Recognizing the urgency in stroke management is vital, as the potential for complete neurological recovery diminishes with every untreated minute. This underscores the critical “time is brain” principle, emphasizing the importance of prompt evaluation and intervention.(3)

Early intervention, tailored treatments, rehabilitative programs, and sustained lifestyle adjustments significantly improve outcomes for acute stroke patients.(4) Maximizing clinical

recovery for each individual not only enhances patient well-being but also mitigates the global impact of stroke. Through timely and targeted approaches, physicians should strive to minimize stroke's devastating effects and foster a healthier society.(5)

DIFFERENT TYPES OF CEREBROVASCULAR ACCIDENTS IN THE ELDERLY POPULATION

To understand how managing and detecting each type of cerebrovascular accident is important for saving the life of the patient, it is equally important that the physicians and clinicians know well and in detail regarding each sub-type:

LARGE VESSEL DISEASE:

Large-vessel disease covers conditions such as atherosclerosis, dissection of arteries, and artery-to-artery embolism, that eventually result in occlusion by thrombus or embolus in major arteries.
(6)

Certain syndromes are linked to blood flow deficiency in particular brain areas found on the test records.(7)Among these large vascular structures are intracranial arteries, for example, the circle of Willis and its main tributaries, and extracranial arteries which include common carotid, internal carotid, and vertebral arteries.(8)

LACUNAR STROKE:

Lexical strokes, frequently coming from small vessel diseases, are often found alongside lipohyalinosis and atherosclerosis. Lipohyalinosis consists of concentric hyaline narrowing of cerebral small arteries, leading to the occlusion of penetrating arteries.(9)

The atherosclerotic plaques of the parent arteries, which are particularly present in the ostium of branchal orifices, can also cause occlusion. Additionally, microatheromas precipitate occlusions in small penetrating arteries.(10)

CARDIOEMBOLISM:

Stroke may also have cardiogenic embolism as its cause, which can originate from diverse sources including arrhythmias, valvular heart disease, bioprosthetic and mechanical heart valves, as well as cardiomyopathy.(11)

Ischemic stroke displays certain links with several main risk factors, namely, advanced age, hypertension, diabetes, hyperlipidemia, cigarette smoking, arrhythmia, and cardiac disease.(12)

INTRACEREBRAL HEMORRHAGE:

It is the second most common type of stroke after the ICH. The ICH is commonly (usually) caused by the rupture of small arteries that happens as a consequence of hypertensive vasculopathy; CAA; coagulopathies; and other vasculopathies.(13)

Most of the VH is found in the non-lobar type of ICH, whereas CAA is usually associated with lobar

The cause of intracerebral hemorrhage (ICH) is attributed to a combination of various risk factors, including age, hypertension, cerebral amyloid angiopathy (CAA), smoking, excessive alcohol consumption, sympathomimetic drugs, anticoagulants, and antiplatelet medications.(14)

SUBARACHNOID HEMORRHAGE:

Subarachnoid hemorrhage (SAH) makes up about 5% of all stroke cases, with 85% of which is caused by aneurysm rupture. (15)

Occurrence of SAH without intervention can also be induced by the use of drugs; amphetamines and cocaine are examples; dural sinus thrombosis; ruptured AVM; rupture due to dural venous sinus thrombosis.(16)

Risk factors for subarachnoid hemorrhage (SAH) encompass smoking, hypertension, excessive alcohol consumption, advanced age, a history of another type of aneurysm or SAH, and a family history of intracranial aneurysms.(17)

THE USAGE OF ADVANCED MRI AND OTHER TECHNIQUES FOR THE EARLY DETECTION AND TREATMENT OF CEREBROVASCULAR ACCIDENTS

The process of clinical diagnosis, management, and treatment of stroke are greatly facilitated by neuroimaging, as well as prognostication. Visual diagnostics in the past decades mostly depended on electroencephalography (EEG), thermography and radioisotope techniques. (18)

Though CT and MRI were a major advance, they made it possible to obtain detailed images of the human brain.(19)

The key aim of these neuroimaging techniques is to pinpoint the affected vascular region of the brain of a stroke patient, namely, the infarct core and the penumbra. The accurate determination of the exact location of the stroke is important in minimizing the extent of a stroke through the selection of the most efficacious treatment approach.(20)

Neuroimaging techniques are broadly categorized into two types: structural and functional. Structural imaging is designed to visualize the anatomic structures of the brain so that any underlying abnormalities like tumors, clots, or bleeding can be detected.(21)

CT and MRI are two exemplars of the structural neuroimaging approach. However, functional imaging methods, such as fMRI and PET, are used to evaluate brain activity in different regions.

A COMPARISON BETWEEN CT SCANS AND MRI FOR THE DETECTION OF CEREBROVASCULAR ACCIDENTS

CT and MRI are both valuable imaging modalities for evaluating patients with acute stroke symptoms, but they have distinct strengths and limitations. (22)

CT is commonly utilized as the initial imaging modality within 24 hours of stroke onset due to its widespread availability and rapid acquisition time.(23) In CT images, blood products appear as distinct hyperintense lesions, making it effective for identifying acute hemorrhagic strokes in the hyperacute phase (between 0 and 6 hours). However, detecting hypointense lesions, indicative of ischemic stroke, in the first few hours of a stroke can be challenging with CT.(24)

In contrast, MRI generates high-resolution images that provide detailed information about the presence, size, and location of hyperacute cerebral ischemic strokes. (19)

MRI is particularly advantageous for visualizing ischemic lesions early in the stroke process. Additionally, MRI is superior to CT in detecting chronic hemorrhages and cerebral microbleeds, making it valuable for assessing the long-term effects of stroke and potential risk factors for recurrent events.(25)

Despite its advantages, MRI is susceptible to artifacts caused by body movements due to its longer scanning time compared to CT. This limitation can affect image quality and may require additional measures to minimize motion artifacts during MRI scanning.(26)

In summary, while CT is often used for the initial evaluation of acute stroke patients due to its speed and ability to detect acute hemorrhages, MRI offers superior sensitivity for detecting ischemic strokes and provides detailed information about chronic hemorrhages and microbleeds. The choice between CT and MRI depends on various factors, including the time since symptom onset, the specific clinical scenario, and the availability of imaging resources.(27)

ADVANCED AND NEW TECHNIQUES FOR THE EARLY DETECTION OF CEREBROVASCULAR ACCIDENTS

Discussed below are some of the recently introduced techniques for the early detection and then, treatment of cerebrovascular accidents, mainly in the elderly populations. It is expected that by utilizing these MRI sequences, clinicians can obtain comprehensive information about stroke pathology, including the type, location, extent, and underlying vascular involvement. (28)

This enables accurate diagnosis, risk stratification, and personalized treatment planning for stroke patients, ultimately improving clinical outcomes and prognosis.

DIFFUSION-WEIGHTED IMAGING (DWI):

DWI harnesses the remarkable sensitivity of MRI to the movement of water molecules within tissues, making it an indispensable tool for detecting acute ischemic stroke. (29)

Ischemic strokes induce cytotoxic edema, restricting the diffusion of water molecules in affected brain tissue. This restriction manifests as hyperintense signals on DWI images, providing clear evidence of ischemic injury. DWI's exceptional sensitivity and specificity make it invaluable for the early identification of ischemic strokes, enabling clinicians to swiftly initiate appropriate treatment strategies aimed at salvaging at-risk brain tissue and mitigating neurological deficits.(30)

FLUID-ATTENUATED INVERSION RECOVERY (FLAIR):

FLAIR represents a key advancement in MRI technology, offering enhanced visualization of brain tissue abnormalities, particularly in the context of stroke.

By selectively suppressing signals from cerebrospinal fluid (CSF), FLAIR imaging highlights pathological changes within the brain parenchyma with remarkable clarity. (31)

In the setting of stroke, FLAIR is adept at detecting hyperintense lesions indicative of subacute and chronic infarctions, as well as delineating regions of edema and gliosis surrounding infarcted tissue. This capability allows for precise characterization of stroke-related pathology, aiding in both diagnosis and treatment planning.(32)

GRADIENT-ECHO (GRE) OR SUSCEPTIBILITY-WEIGHTED IMAGING (SWI):

GRE or SWI sequences represent a significant advancement in MRI technology, offering unparalleled sensitivity to magnetic susceptibility variations associated with hemorrhagic stroke pathology. (33)

Hemorrhagic strokes, including intracerebral hemorrhage (ICH) and subarachnoid hemorrhage (SAH), manifest as hypointense signals on GRE or SWI images due to the presence of blood products. This distinctive appearance enables accurate and precise localization of hemorrhagic lesions within the brain, facilitating rapid diagnosis and appropriate intervention. (34)

Additionally, GRE or SWI sequences are invaluable for detecting other pathological features, such as calcifications or paramagnetic substances, further enhancing their utility in stroke imaging.

PERFUSION-WEIGHTED IMAGING (PWI):

PWI represents a cutting-edge MRI technique that provides crucial insights into cerebral perfusion dynamics, offering valuable information for the assessment and management of acute stroke patients. (35)

By measuring the passage of contrast agents through brain tissue, PWI enables the identification of areas exhibiting hypoperfusion, ischemia, and penumbra - regions of salvageable but at-risk tissue. (36)

This information is invaluable for guiding treatment decisions, such as thrombolytic therapy or endovascular intervention, by identifying viable tissue targets and optimizing patient outcomes.

MAGNETIC RESONANCE ANGIOGRAPHY (MRA):

MRA stands at the forefront of stroke imaging technology, offering unparalleled visualization of the cerebral vasculature and aiding in the identification of vascular pathologies contributing to stroke etiology. (37)

By generating detailed images of arterial stenosis, occlusions, and collateral circulation, MRA enables clinicians to pinpoint underlying vascular abnormalities, such as carotid artery stenosis or intracranial vessel pathology. (38)

This comprehensive assessment of the cerebrovascular system facilitates accurate diagnosis, risk stratification, and treatment planning, ultimately improving patient care and outcomes in the management of stroke.

CONCLUSION

Neuroimaging stands as an indispensable tool in stroke diagnosis, facilitating the swift identification of patients who may benefit from specific interventions like thrombolytic agents or surgery. Despite the continued preference for CT due to its wider accessibility and quicker scans in acute stroke assessment, recent advancements in MRI technology have bolstered its role in routine clinical stroke protocols. As MRI continues to evolve, its potential to become the primary imaging platform in stroke diagnosis and management is increasingly promising. In conclusion, MRI plays a pivotal role in both diagnosing and treating stroke, offering various techniques such as SWI, PWI, and DWI to identify and assess different aspects of the condition. Furthermore, ongoing advancements in MRI technology, including ultra-high magnetic fields, MRF, and neural network mapping, are enhancing its capabilities in predicting and managing strokes.

These innovations not only aid in deciphering the wealth of data collected but also pave the way for novel MR contrast agents, which promise to revolutionize diagnostic precision. With ongoing research, the future holds promising prospects for further refining MRI's role in stroke management and improving patient outcomes.

REFERENCES

1. Tadi P, Lui F. Acute Stroke. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 12]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK535369/>
2. Hui C, Tadi P, Patti L. Ischemic Stroke. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK499997/>
3. Musuka TD, Wilton SB, Traboulsi M, Hill MD. Diagnosis and management of acute ischemic stroke: speed is critical. *CMAJ Can Med Assoc J*. 2015 Sep 8;187(12):887–93.
4. Patil S, Rossi R, Jabrah D, Doyle K. Detection, Diagnosis and Treatment of Acute Ischemic Stroke: Current and Future Perspectives. *Front Med Technol*. 2022 Jun 24;4:748949.
5. Hurford R, Sekhar A, Hughes TAT, Muir KW. Diagnosis and management of acute ischaemic stroke. *Pract Neurol*. 2020 Aug;20(4):304–16.
6. Betrains A, Blockmans D. Diagnostic Approaches for Large Vessel Vasculitides. *Open Access Rheumatol Res Rev*. 2021 Jun 1;13:153–65.
7. Pugh D, Karabayas M, Basu N, Cid MC, Goel R, Goodyear CS, et al. Large vessel vasculitis. *Nat Rev Dis Primer*. 2022 Jan 6;7(1):93.
8. Harky A, Fok M, Balmforth D, Bashir M. Pathogenesis of large vessel vasculitis: Implications for disease classification and future therapies. *Vasc Med Lond Engl*. 2019 Feb;24(1):79–88.
9. Gore M, Bansal K, Khan Suheb MZ, Asuncion RMD. Lacunar Stroke. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK563216/>
10. Regenhardt RW, Das AS, Lo EH, Caplan LR. Advances in Lacunar Stroke Pathophysiology: A Review. *JAMA Neurol*. 2018 Oct 1;75(10):1273–81.

11. Pillai AA, Tadi P, Kanmanthareddy A. Cardioembolic Stroke. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK536990/>
12. Leary MC, Caplan LR. Cardioembolic stroke: An update on etiology, diagnosis and management. *Ann Indian Acad Neurol*. 2008 Jan;11(Suppl 1):S52–63.
13. Rajashekar D, Liang JW. Intracerebral Hemorrhage. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK553103/>
14. Tenny S, Thorell W. Intracranial Hemorrhage. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK470242/>
15. Ziu E, Khan Suheb MZ, Mesfin FB. Subarachnoid Hemorrhage. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK441958/>
16. Kairys N, M Das J, Garg M. Acute Subarachnoid Hemorrhage. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK518975/>
17. Marcolini E, Hine J. Approach to the Diagnosis and Management of Subarachnoid Hemorrhage. *West J Emerg Med*. 2019 Mar;20(2):203–11.
18. Yew KS, Cheng E. Acute Stroke Diagnosis. *Am Fam Physician*. 2009 Jul 1;80(1):33–40.
19. Bhatkar S, Mahesh KV, Sachdeva J, Goel A, Goyal MK, Takkar A, et al. Magnetic resonance imaging (MRI) versus computed tomographic scan (CT scan) of brain in evaluation of suspected cavernous sinus syndrome. *Neuroradiol J*. 2020 Dec;33(6):501–7.
20. Qin Y, Bao A, Li H, Wang X, Zhang G, Zhu J. Application value of CT and MRI in diagnosis of primary brain lymphoma. *Oncol Lett*. 2018 Jun;15(6):8500–4.
21. Abdalkader M, Siegler JE, Lee JS, Yaghi S, Qiu Z, Huo X, et al. Neuroimaging of Acute Ischemic Stroke: Multimodal Imaging Approach for Acute Endovascular Therapy. *J Stroke*. 2023 Jan;25(1):55–71.
22. Vitali P, Savoldi F, Segati F, Melazzini L, Zanardo M, Fedeli MP, et al. MRI versus CT in the detection of brain lesions in patients with infective endocarditis before or after cardiac surgery. *Neuroradiology*. 2022;64(5):905–13.
23. Bhargava R. CT Imaging in Neurocritical Care. *Indian J Crit Care Med Peer-Rev Off Publ Indian Soc Crit Care Med*. 2019 Jun;23(Suppl 2):S98–103.
24. Masdeu JC, Gadhia R, Faridar A. Brain CT and MRI: differential diagnosis of imaging findings. *Handb Clin Neurol*. 2016;136:1037–54.

25. Grover VPB, Tognarelli JM, Crossey MME, Cox IJ, Taylor-Robinson SD, McPhail MJW. Magnetic Resonance Imaging: Principles and Techniques: Lessons for Clinicians. *J Clin Exp Hepatol*. 2015 Sep;5(3):246–55.
26. Chow MSM, Wu SL, Webb SE, Gluskin K, Yew DT. Functional magnetic resonance imaging and the brain: A brief review. *World J Radiol*. 2017 Jan 28;9(1):5–9.
27. Despotović I, Goossens B, Philips W. MRI Segmentation of the Human Brain: Challenges, Methods, and Applications. *Comput Math Methods Med*. 2015;2015:450341.
28. Khaku AS, Tadi P. Cerebrovascular Disease. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK430927/>
29. Baliyan V, Das CJ, Sharma R, Gupta AK. Diffusion weighted imaging: Technique and applications. *World J Radiol*. 2016 Sep 28;8(9):785–98.
30. Chilla GS, Tan CH, Xu C, Poh CL. Diffusion weighted magnetic resonance imaging and its recent trend—a survey. *Quant Imaging Med Surg*. 2015 Jun;5(3):407–22.
31. Sati P, George IC, Shea CD, Gaitán MI, Reich DS. FLAIR*: A Combined MR Contrast Technique for Visualizing White Matter Lesions and Parenchymal Veins. *Radiology*. 2012 Dec;265(3):926–32.
32. Sohn CH, Sevick RJ, Frayne R, Chang HW, Kim SP, Kim DK. Fluid Attenuated Inversion Recovery (FLAIR) Imaging of the Normal Brain: Comparisons between Under the Conditions of 3.0 Tesla and 1.5 Tesla. *Korean J Radiol*. 2010;11(1):19–24.
33. Hargreaves B. Rapid Gradient-Echo Imaging. *J Magn Reson Imaging JMRI*. 2012 Dec;36(6):1300–13.
34. Copenhaver BR, Shin J, Warach S, Butman JA, Saver JL, Kidwell CS. Gradient echo MRI. *Neurology*. 2009 May 5;72(18):1576–81.
35. Xu W, Wang Q, Shao A, Xu B, Zhang J. The performance of MR perfusion-weighted imaging for the differentiation of high-grade glioma from primary central nervous system lymphoma: A systematic review and meta-analysis. *PLoS ONE*. 2017 Mar 16;12(3):e0173430.
36. Mannam SS, Nwagwu CD, Sumner C, Weinberg BD, Hoang KB. Perfusion-Weighted Imaging: The Use of a Novel Perfusion Scoring Criteria to Improve the Assessment of Brain Tumor Recurrence versus Treatment Effects. *Tomography*. 2023 May 23;9(3):1062–70.
37. De Leucio A, De Jesus O. MR Angiogram. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 13]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK558984/>

38. Ludwig DR, Shetty AS, Broncano J, Bhalla S, Raptis CA. Magnetic Resonance Angiography of the Thoracic Vasculature: Technique and Applications. *J Magn Reson Imaging JMRI*. 2020 Aug;52(2):325–47.

UNDER PEER REVIEW