

1
2 **A Computational Approach on**
3 **Fenofibrate drug using Degree-Based**
4 **Topological indices and M-Polynomials**
6

9
10
11
12
13 **ABSTRACT**
14

Fenofibrate is a drug approved by FDA Food and Drugs Administration used to treat patients with Hypertriglyceridemia primary hypercholesterolemia or mixed dyslipidemia. It reduces low-density lipoprotein cholesterol (LDL-c), total cholesterol, triglycerides, and apolipoprotein B and increases high-density lipoprotein cholesterol (HDL-C) in adults. The sum and multiplicative versions of several topological indices such as General Zagreb, General Randić, Arithmetic Geometric Index, Inverse sum (Indeg) Index, Symmetric Division (Deg) Index, Forgotten Indices M-polynomials of Fenofibrate are computed in this article.

15
16 *Keywords: Topological indices, Fenofibrate, M-Polynomial, Chemical graph.*
17
18

19 **1. INTRODUCTION**
20

21 All molecular graphs in this paper are finite, loopless, connected, and do not have
22 numerous edges. $A = (V, E)$ graph having vertex set V and edge set E .

23 In Mathematical chemistry the analytical and structural properties of topological
24 indices have been studied in depth over the last years. The Theoretical and practical interest
25 of topological indices lies in the fact that they have become an important tool for the study of
26 multiple practical problems in Computer science, Physics, Ecology, among others countless
27 applications of topological indices have been reported, most of them concerned with
28 exploring medicinal and pharmacological issues. A molecular graph is a graph-theoretic
29 representation of the structural formula of a chemical molecule, with edges standing in for
30 chemical bonds and vertices for atoms. Topological indices based on end-vertex degrees of
31 edges have been used over 50 years. Among them, several indices are recognized to be
32 useful tools in chemical researches. Probably the best known such descriptor is the Randić
33 connectivity index. In 1972, Trinajstić and Gutman obtained a formula concerning the total
34 energy of π electrons of molecules [1-3].

35 Fenofibrate was first synthesized in 1974, as a derivative of clofibrate, and was
36 initially offered in France. It was initially known as procetofen, and was later renamed
37 Fenofibrate to comply with World Health Organization international Nomenclature
38 guidelines. It is developed by Groupe Fournier SA of France. Fenofibrate comes as a capsule
39 , a delayed-release (long-acting) capsule, and a tablet to take by mouth. It is usually taken
40 once a day. Some Fenofibrate products (Fenoglide, Lipofen and Lofibra) should be taken

41 with a meal. Other brands (Antara, Fibracor, Tricor, Triglide and Trilipix) may be taken with or
 42 without food. Its molecular formula is $C_{20}H_{21}ClO_4$.[4]

43

44

45

46

47




Figure 1-Chemical structure of Fenofibrate

48

49

50

51

52

53

54

55

56

57

58

59




Figure 2- Molecular graph of Fenofibrate

60 2. MATERIAL AND METHODS

61

62 2.1. Additive degree -based Topological indices.

63

64

65

66

67

Topological indices are classified into different categories such as entropy-based, eigenvalue-based, spectrum-based, symmetry-based, distance-based and vertex-degree-based. Among these categories, *degree-based indices* have an outstanding position. . Assuming that the edge connecting vertices r and s is denoted by rs , the additive degree-based In mathematical chemistry, topological indices have a general form.

68

$$TI = TI(A) = \sum_{rs \in E(A)} F(d_r, d_s)$$

69

Table 1. Topological indices of a graph with additive degree basis.

70

S. No.	Topological index	Notation	Formula of Topological index
1	First Zagreb Index	$M_1(A)$	$\sum_{rs \in E(A)} (d_r + d_s)$
2	Second Zagreb Index	$M_2(A)$	$\sum_{rs \in E(A)} (d_r \cdot d_s)$
3	Third Zagreb Index	$M_3(A)$	$\sum_{rs \in E(A)} d_r - d_s $
4	Modified First Zagreb Index	$M_1^*(A)$	$\sum_{s \in V(A)} (d_s)^2$
5	Randic Index	$R(A)$	$\sum_{rs \in E(A)} \frac{1}{\sqrt{d_r \cdot d_s}}$
6	First Hyper Zagreb Index	$HM_1(A)$	$\sum_{rs \in E(A)} (d_r + d_s)^2$

7	Second Hyper Zagreb Index	$HM_2(A)$	$\sum_{rs \in E(A)} (d_r \cdot d_s)^2$
8	General First Zagreb Index	$M_1^\alpha(A)$	$\sum_{rs \in E(A)} (d_r + d_s)^\alpha$
9	General Second Zagreb Index	$M_2^\alpha(A)$	$\sum_{rs \in E(A)} (d_r \cdot d_s)^\alpha$
10	Reduced Second Zagreb Index	$RM_2(A)$	$\sum_{rs \in E(A)} (d_r - 1)(d_s - 1)$
11	Redefined First Zagreb Index	$Re M_1(A)$	$\sum_{rs \in E(A)} \frac{(d_r + d_s)}{(d_r \cdot d_s)}$
12	Redefined Second Zagreb Index	$Re M_2(A)$	$\sum_{rs \in E(A)} \frac{(d_r \cdot d_s)}{(d_r + d_s)}$
13	Redefined Third Zagreb Index	$Re M_3(A)$	$\sum_{rs \in E(A)} (d_r \cdot d_s)(d_r + d_s)$
14	Modified Second Zagreb Index	${}^m M_2(A)$	$\sum_{rs \in E(A)} \frac{1}{d_r \cdot d_s}$
15	Nano - Zagreb Index	$N_z(A)$	$\sum_{s \in V(A)} [(d_r)^2 - (d_s)^2]$
16	Sum Nano - Zagreb Index	$x_{\frac{1}{2}} N_z(A)$	$\sum_{rs \in E(A)} (d_r^2 - d_s^2)^{\frac{1}{2}}$
17	Zeroth order Randić Index	$R_{\frac{-1}{2}}^0(A)$	$\sum_{s \in V(A)} (d_s)^{\frac{-1}{2}}$
18	Zeroth Order General Randić Index	$R_\alpha^0(A)$	$\sum_{s \in V(A)} (d_s)^\alpha$
19	Reciprocal Randić Index	$RR(A)$	$\sum_{rs \in E(A)} \sqrt{d_r \cdot d_s}$
20	Reduced Reciprocal Randić Index	$RRR(A)$	$\sum_{rs \in E(A)} \sqrt{(d_r - 1) \cdot (d_s - 1)}$
21	Inverse Randić Index	$RR_\alpha(A)$	$\sum_{rs \in E(A)} \frac{1}{(d_r \cdot d_s)^\alpha}$
22	Modified Randić Index	$R'(A)$	$\sum_{rs \in E(A)} \frac{1}{\max\{d_r \cdot d_s\}}$
23	Sum connectivity Index	$SCI(A)$	$\sum_{rs \in E(A)} \frac{1}{\sqrt{d_r + d_s}}$
24	Arithmetic Geometric Index	$AG(A)$	$\sum_{rs \in E(A)} \frac{d_r + d_s}{2\sqrt{d_r \cdot d_s}}$
25	Variable Sum Index	$SEI_\alpha(A)$	$\sum_{s \in V(A)} d_s(\alpha)^{d_s}$

26	Harmonic Index	$H(A)$	$\sum_{rs \in E(A)} \frac{2}{d_r + d_s}$
27	General Harmonic Index	$H_k(A)$	$\sum_{rs \in E(A)} \left(\frac{2}{d_r + d_s} \right)^k$
28	General Ordinary Geometric Arithmetic Index	$OGA_k(A)$	$\sum_{rs \in E(A)} \left(\frac{2\sqrt{d_r \cdot d_s}}{d_r + d_s} \right)^k$
29	Ordinary Geometric Arithmetic Index	$OGA(A)$	$\sum_{rs \in E(A)} \frac{2\sqrt{d_r \cdot d_s}}{d_r + d_s}$
30	SK Index	$SK(A)$	$\sum_{rs \in E(A)} \frac{d_r + d_s}{2}$
31	SK1 Index	$SK_1(A)$	$\sum_{rs \in E(A)} \frac{d_r \cdot d_s}{2}$
32	SK2 Index	$SK_2(A)$	$\sum_{rs \in E(A)} \left(\frac{d_r + d_s}{2} \right)^2$
33	Forgotten Index	$F(A)$	$\sum_{rs \in E(A)} [(d_r)^2 + (d_s)^2]$
34	Inverse sum (Indeg) Index	$ISI(A)$	$\sum_{rs \in E(A)} \frac{d_r \cdot d_s}{d_r + d_s}$
35	Symmetric Division (Deg) Index	$SDD(A)$	$\sum_{rs \in E(A)} \left[\frac{\min\{d_r, d_s\} + \max\{d_r, d_s\}}{\max\{d_r, d_s\} \cdot \min\{d_r, d_s\}} \right]$
36	IRM Index	$IRM(A)$	$\sum_{rs \in E(A)} (d_r - d_s)^2$
37	Augmented Zagreb Index	$AZI(A)$	$\sum_{rs \in E(A)} \left(\frac{d_r \cdot d_s}{d_r + d_s - 2} \right)^3$
38	Albertson Index	$A(A)$	$\sum_{rs \in E(A)} d_r - d_s $
39	Atomic Bond Connectivity Index	$ABC(A)$	$\sum_{rs \in E(A)} \sqrt{\frac{d_r + d_s - 2}{d_r \cdot d_s}}$
40	Bell Index	$B(A)$	$\sum_{s \in V(A)} \left(d_s - \frac{2q}{p} \right)^2$
41	First Entire Zagreb Index	$M_1^e(A)$	$\sum_{s \in V(A) \cup E(A)} (d_s)^2$
42	Sombor Index	$SO(A)$	$\sum_{rs \in E(A)} \sqrt{(d_r)^2 + (d_s)^2}$

71
72

73 **2.2. Multiplicative degree -based Topological indices.**

74 Several significant multiplicative degree-based Topological indices were outlined in
 75 this section. This can be used to provide our primary results[5,6]. $NK(A)$ is the Multiplicative
 76 Topological index that Narumi and Katayama suggested. Following that, $NK(A)$ was
 77 redefined by Wang et al. as a broad multiplicative index. In order to advance the study of
 78 Topological indices, Kulli.V R, devised the multiplicative sum connectivity index($SCII(A)$)
 79 and multiplicative product connectivity index ($PCII(A)$) ratings. The generalized version of
 80 multiplicative Topological indices is given below, according to Chemical Graph Theory.

81
$$MTI = MTI(A) = \prod_{rs \in E(A)} F(d_r, d_s)$$

82 **Table2. Topological indices of a graph A based on multiplication of degrees**

S. No.	Topological index	Notation	Formula of Topological index
1	Narumi – Katayama Index	$NK(A)$	$\prod_{s \in V(A)} (d_s)$
2	First Multiplicative Zagreb Index	$\Pi_1(A)$	$\prod_{s \in V(A)} (d_s)^2$
3	General Multiplicative Index	$W_1^\alpha(A)$	$\prod_{s \in V(A)} (d_s)^\alpha$
4	First Multiplicative Generalized Zagreb Index	$MZ_1^\alpha(A)$	$\prod_{uv \in E(G)} (d_u + d_v)^\alpha$
5	Second Multiplicative Generalized Zagreb Index	$MZ_2^\alpha(A)$	$\prod_{rs \in E(A)} (d_r \cdot d_s)^\alpha$
6	Multiplicative version of First Zagreb Index	$\Pi_1^*(A)$	$\prod_{rs \in E(A)} (d_r + d_s)$
7	Second Multiplicative Zagreb Index	$\Pi_2(A)$	$\prod_{rs \in E(A)} (d_r \cdot d_s)$
8	Multiplicative First Hyper Zagreb Index	$H\Pi_1(A)$	$\prod_{rs \in E(A)} (d_r + d_s)^2$
9	Multiplicative Second Hyper Zagreb Index	$H\Pi_2(A)$	$\prod_{rs \in E(A)} (d_r \cdot d_s)^2$
10	Multiplicative Sum Connectivity Index	$SC\Pi(A)$	$\prod_{rs \in E(A)} \frac{1}{\sqrt{d_r + d_s}}$
11	Multiplicative Product Connectivity Index	$PC\Pi(A)$ (or) $R\Pi(A)$	$\prod_{rs \in E(A)} \frac{1}{\sqrt{d_r \cdot d_s}}$

12	Multiplicative Sum Connectivity F –Index	$SCF \Pi(A)$	$\prod_{rs \in E(A)} \frac{1}{\sqrt{d_r^2 + d_s^2}}$
13	Multiplicative Product Connectivity F – Index	$PG \Pi(A)$	$\prod_{rs \in E(A)} \frac{1}{\sqrt{d_r^2 \cdot d_s^2}}$
14	Multiplicative First F –Index	$F_1 \Pi(A)$	$\prod_{rs \in E(A)} [(d_r)^2 + (d_s)^2]$
15	Multiplicative Second F – Index	$F_2 \Pi(A)$	$\prod_{rs \in E(A)} [(d_r)^2 \cdot (d_s)^2]$
16	Multiplicative Atomic Bond Connectivity Index	$ABC \Pi(A)$	$\prod_{rs \in E(A)} \sqrt{\frac{d_r + d_s - 2}{d_r \cdot d_s}}$
17	Multiplicative Harmonic Index	$H \Pi(A)$	$\prod_{rs \in E(A)} \frac{2}{d_r + d_s}$
18	Multiplicative Geometric Arithmetic Index	$GA \Pi(A)$	$\prod_{rs \in E(A)} \frac{2\sqrt{d_r \cdot d_s}}{d_r + d_s}$
19	General Multiplicative Geometric Arithmetic Index	$GA^\alpha \Pi(A)$	$\prod_{rs \in E(A)} \left(\frac{2\sqrt{d_r \cdot d_s}}{d_r + d_s} \right)^\alpha$
20	Multiplicative Augmented Zagreb Index	$AZ \Pi(A)$	$\prod_{rs \in E(A)} \left(\frac{d_r \cdot d_s}{d_r + d_s - 2} \right)^3$

83
84
85
86
87
88
89
90
91
92
93
94
95
96

2.3. Zagreb, Forgotten, Harmonic, Inverse degree indices and M-Polynomials of a graph.

Fath-Tabar provided the definitions of the first, second, and third Zagreb polynomials. Harmonic polynomial are introduced by Iranmanesh et al[7]. Shuxian[8] established the modified first zagreb index. In 2015 [9], Deutsch and Klavzar created a number of new algebraic polynomials. By specifying the closed form of specific degree-based Topological indices, it serves a similar function. One of M-Polynomial's best features is the abundance of information it offers on molecular labels. A few polynomials of graph invariants are shown in Tables 3 and 4.

Table 3: Topological indices of a graph A depending on degree and their corresponding polynomials.

S. No.	Topological index	Polynomial $P(A, a)$	Derivation from $P(A, a)$
--------	-------------------	----------------------	---------------------------

1	First Zagreb	$M_1(A, a) = \sum_{rs \in E(A)} a^{d_r + d_s}$	$D_a(M_1(A; a)) _{a=1}$
2	Second Zagreb	$M_2(A, a) = \sum_{rs \in E(A)} a^{d_r \cdot d_s}$	$D_a(M_2(A; a)) _{a=1}$
3	Third Zagreb	$M_3(A, a) = \sum_{rs \in E(A)} a^{ d_r - d_s }$	$D_a(M_3(A; a)) _{a=1}$
4	Forgotten	$F(A, a) = \sum_{rs \in E(A)} a^{(d_r^2 + d_s^2)}$	$D_a(F(A; a)) _{a=1}$
5	Modified First Zagreb	$M_1^*(A, a) = \sum_{s \in V(A)} d_s a^{d_s}$	$D_a(M_1^*(A; a)) _{a=1}$
6	Harmonic	$H(A, a) = \sum_{rs \in E(A)} a^{d_r + d_s - 1}$	$2I_a(H(A; a)) _{a=1}$
7	Modified Forgotten	$F^*(A, a) = \sum_{s \in V(A)} a^{d_s^3}$	$D_a(F^*(A; a)) _{a=1}$
8	Inverse Degree	$ID(A, a) = \sum_{s \in V(A)} a^{d_s - 1}$	$I_a(H(A; a)) _{a=1}$

97
98
99

Theorem 2.1. [9] Assume that A is a simple connected graph.

100 (1) If $I(A) = \sum_{rs \in E(A)} f(dr \cdot ds)$, where $f(a, b)$ is a polynomial in a and b , then

101
$$I(A) = f(D_a, D_b)(M(A; a, b))|_{a=b=1}.$$

102 (2) If $I(A) = \sum_{rs \in E(A)} f(dr \cdot ds)$, where, $I(A)$ can be obtained from $M(A; a, b)$ using the

103 operators D_a, D_b, S_a, S_b .

104 (3) If $I(A) = \sum_{rs \in E(A)} f(dr \cdot ds)$, where $f(a, b) = \frac{a^i b^j}{(a + b + \alpha)^t}$, where a and b are

105 variables, $i, j \geq 0, t \geq 1$, then $I(A) = S_a^t Q_\alpha J D_a^i D_b^j (M(A; a, b))|_{a=b=1}$.

106 The definition of a molecular graph's M-Polynomial [10-13] is

107
$$M(A; a, b) = \sum_{\delta \leq i \leq j \Delta} m_{ij}(A) a^i b^j \text{ -----(1)}$$

108
109
110

Table 4 .Degree –based Topological indices derived from M-Polynomial of a graph A.

S. No	Topological index	$f(a, b)$	Derivation from $M(A; a, b)$
1	$M_1(A)$	$a + b$	$(D_a + D_b)(M(A; a)) _{a=b=1}$
2	$M_2(A)$	ab	$(D_a D_b)(M(A; a)) _{a=b=1}$
3	${}^m M_2(A)$	$\frac{1}{ab}$	$(S_a S_b)(M(A; a)) _{a=b=1}$

4	$SDD(A)$	$\frac{a^2 + b^2}{ab}$	$(D_a S_b + D_b S_a)(M(A; a)) _{a=b=1}$
5	$H(A)$	$\frac{2}{a+b}$	$2S_a J(M(A; a)) _{a=b=1}$
6	$ISI(A)$	$\frac{ab}{a+b}$	$S_a J D_a D_a (M(A; a)) _{a=b=1}$
7	$AZI(A)$	$\left(\frac{ab}{a+b-2}\right)^3$	$S_a^3 Q_{-2} J D_a^3 D_b^3 (M(A; a)) _{a=b=1}$
8	$F(A)$	$a^2 + b^2$	$(D_a^2 + D_b^2)(M(A; a)) _{a=b=1}$
9	$Re M_3(A)$	$ab(a+b)$	$(D_a D_b)(D_a + D_b)(M(A; a)) _{a=b=1}$
10	$ABC(A)$	$\sqrt{\frac{a+b-2}{ab}}$	$S_a^{-2} Q_{-2} J D_b^{-2} (M(A; a)) _{a=b=1}$

111
112
113
114

To calculate the corresponding Topological indices of a graph A from the $M(A; a, b)$ as stated in Table 4. Table 2 lists the equations of derivations in terms of integral, derivative, or both integral and derivative. [9], where

115
$$D_a = a \frac{\partial(f(a,b))}{\partial a}, \quad D_b = b \frac{\partial(f(a,b))}{\partial b}, \quad S_a = \int_0^a \frac{f(t,b)}{t} dt, \quad S_b = \int_0^{ba} \frac{f(a,t)}{t} dt,$$

116
$$J(f(a,b)) = f(a,a) \text{ and } Q_\alpha(f(a,b)) = x^\alpha f(a,b), \quad \alpha \neq 0$$

117
118
119

2.4. Methodolgy

120
121
122
123
124
125
126
127
128
129
130
131

One of our main conclusions is that Tis of antiviral drug structures can be derived using algebraic polynomials. Fenofibrate molecular structure can be accessed at pubchem.ncbi.nlm.nih.gov. hydrogen suppressed molecular graph of a compound can be considered, since the vertices representing the hydrogen atoms do not contribute to graph isomorphism. We employ edge partitioning, degree counting, graph theoretical tools, combinatorial computation, and analytical techniques to arrive at our main findings. Using edge partitions and the formulas listed in Tables 1 and 2, expressions of additive and multiplicative degree-based Topological indices were calculated. likewise, Tables 3 and 4 are used to generate several closed versions of M-Polynomials, Zagreb, Forgotten, Harmonic, Inverse degree polynomials, and so on. To compare the numerical findings graphically, MATLAB 2017b was used to plot the polynomials on a surface.

3. RESULTS AND DISCUSSION

132
133
134
135

This section begins with the following derivation of the ID, Zagreb, Harmonic, and Forgotten polynomials.

136 **Theorem 3.1.** For a molecular graph Fenofibrate ($C_{20}H_{21}ClO_4$) drug . Then we have

137 (1) $M_1(A, a) = a^7 + 3a^6 + 13a^5 + 9a^4.$

138 (2) $M_2(A, a) = a^{12} + 2a^9 + a^8 + 11a^6 + 6a^4 + 5a^3.$

139 (3) $M_3(A, a) = 2a^3 + 6a^2 + 12a + 6$

140 (4) $F(A, a) = a^{25} + a^{20} + 2a^{18} + 2a^{17} + 11a^{13} + 5a^{10} + 4a^8$.

141 (5) $H(A, a) = a^6 + 3a^5 + 13a^4 + 9a^3$.

142 (6) $M_1^*(A, a) = 4a^4 + 21a^3 + 20a^2 + 7a$.

143 (7) $F^*(A, a) = a^{64} + 7a^{27} + 10a^8 + 7a^3$.

144 (8) $D(A, a) = a^3 + 7a^2 + 10a + 7$.

145 **Proof:** Consider a molecular graph of Fenofibrate as A . There are 26 edges and 25 vertices
 146 in the graph A . Let J_m be the collection of the vertices with degree m , i.e.,

147 $J_m = \{s \in V(A) : d_s = m\}$. Let j_m be the number of vertices in J_m . Let $K_{n,m}$ be the set that
 148 has edges with vertices of degree n and m . i.e., $k_{n,m} = \{rs \in E(A) : d_r = n, d_s = m\}$. $k_{n,m}$
 149 denotes the number of edges in. $j_1 = 7, j_2 = 10, j_3 = 7, j_4 = 1$

$d_r, d_s : rs \in E(A)$	Number of edges
(1,3)	5
(1,4)	2
(2,2)	4
(2,4)	1
(2,3)	11
(3,3)	2
(3,4)	1

150 Applying the polynomial formulas provided in Table 3, we obtain

151 (1) $M_1(A, a) = \sum_{rs \in E(A)} k_{n,m} a^{n+m}$
 152 $= 5a^{(1+3)} + 11a^{(2+3)} + 4a^{(2+2)} + 2a^{(1+4)} + a^{(2+4)} + a^{(3+4)} + 2a^{(3+3)}$
 153 $= a^7 + 3a^6 + 13a^5 + 9a^4$

154 (2) $M_2(A, a) = \sum_{rs \in E(A)} k_{n,m} a^{n \cdot m}$
 155 $= 5a^{(1 \cdot 3)} + 11a^{(2 \cdot 2)} + 2a^{(1 \cdot 4)} + a^{(2 \cdot 4)} + a^{(3 \cdot 4)} + 2a^{(3 \cdot 3)}$
 156 $= a^{12} + 2a^9 + a^8 + 11a^6 + 6a^4 + 5a^3$

157 (3) $M_3(A, a) = \sum_{rs \in E(A)} k_{n,m} a^{|n-m|}$
 158 $= 5a^{|1-3|} + 11a^{|2-3|} + 4a^{|2-2|} + 2a^{|1-4|} + a^{|2-4|} + a^{|3-4|} + 2a^{|3-3|}$
 159 $= 2a^3 + 6a^2 + 12a + 6$

160 (4) $F(A, a) = \sum_{rs \in E(A)} k_{n,m} a^{(n^2+m^2)}$
 161 $= 5a^{(1^2+3^2)} + 11a^{(2^2+3^2)} + 4a^{(2^2+2^2)} + 2a^{(1^2+4^2)} + a^{(2^2+4^2)} + a^{(3^2+4^2)} + 2a^{(3^2+3^2)}$
 162 $= a^{25} + a^{20} + 2a^{18} + 2a^{17} + 11a^{13} + 5a^{10} + 4a^8$

163 (5) $M_1^*(A, a) = \sum_{rs \in E(A)} m a^m$
 164 $= j_1(1)a^1 + j_2(2)a^2 + j_3(3)a^3 + j_4(4)a^4$
 165 $= 4a^4 + 21a^3 + 20a^2 + 7a$

$$\begin{aligned}
166 \quad (6) \quad H(A, a) &= \sum_{rs \in E(A)} k_{n,m} a^{(n+m-1)} \\
167 & \\
168 &= 5a^{(1+3-1)} + 11a^{(2+3-1)} + 4a^{(2+2-1)} + 2a^{(1+4-1)} + a^{(2+4-1)} + a^{(3+4-1)} + 2a^{(3+3-1)} \\
169 &= a^6 + 3a^5 + 13a^4 + 9a^3
\end{aligned}$$

$$\begin{aligned}
170 \quad (7) \quad F^*(A, a) &= \sum_{rs \in E(A)} k_{n,m} a^{m^3} \\
171 &= j_1 a^{(1)^3} + j_2 a^{(2)^3} + j_3 a^{(3)^3} + j_4 a^{(4)^3} \\
172 &= a^{64} + 7a^{27} + 10a^8 + 7a^3
\end{aligned}$$

$$\begin{aligned}
173 \quad (8) \quad ID(A, a) &= \sum_{rs \in E(A)} a^{m-1} \\
174 &= j_1 a^{1-1} + j_2 a^{2-1} + j_3 a^{3-1} + j_4 a^{4-1} \\
175 &= a^3 + 7a^2 + 10a + 7.
\end{aligned}$$

176 We obtain the relevant Topological indices using the polynomials mentioned below.

177

178 **Theorem 3.2** If A is a molecular graph of Fenofibrate ($C_{20}H_{21}ClO_4$) drug then,

- 179 (i) $M_1(A) = 126$
180 (ii) $M_2(A) = 143$
181 (iii) $M_3(A) = 30$
182 (iv) $F(A) = 340$
183 (v) $M_1^*(A) = 126$
184 (vi) $H(A) = 7.74$
185 (vii) $F^*(A) = 354$
186 (viii) $ID(A) = 14.55$

187 **Proof:** We derive using Table 3 and Theorem 3.1,

$$\begin{aligned}
188 \quad (i) \quad M_1(A, a) &= D_a(M_1(A; a))|_{a=1} = \frac{d}{da} (a^7 + 3a^6 + 13a^5 + 9a^4) \\
&= 7a^6 + 18a^5 + 65a^4 + 36a^3 = 126
\end{aligned}$$

$$\begin{aligned}
189 \quad (ii) \quad M_2(A, a) &= D_a(M_2(A; a))|_{a=1} = \frac{d}{da} (a^{12} + 2a^9 + a^8 + 11a^6 + 6a^4 + 5a^3) \\
&= 12a^{11} + 18a^8 + 8a^7 + 66a^5 + 24a^3 + 15a^2 = 143
\end{aligned}$$

$$\begin{aligned}
190 \quad (iii) \quad M_3(A, a) &= D_a(M_3(A; a))|_{a=1} = \frac{d}{da} (2a^3 + 6a^2 + 12a + 6) \\
&= 6a^2 + 12a + 12 = 30
\end{aligned}$$

$$\begin{aligned}
191 \quad (iv) \quad F(A, a) &= D_a(F(A; a))|_{a=1} = \frac{d}{da} (a^{25} + a^{20} + 2a^{18} + 2a^{17} + 11a^{13} + 5a^{10} + 4a^8) \\
192 &= 25a^{24} + 20a^{19} + 36a^{17} + 34a^{16} + 143a^{12} + 50a^9 + 32a^7 = 340
\end{aligned}$$

193 (v) $M_1^*(A, a) = D_a(M_1^*(A; a))|_{a=1} = \frac{d}{da}(4a^4 + 21a^3 + 20a^2 + 7a)$
 $= 16a^3 + 63a^2 + 40a + 7 = 126$

194 (vi) $H(A, a) = 2I_a(H(A; a))|_{a=1} = 2 \int_0^a (a^6 + 3a^5 + 13a^4 + 9a^3) d_a$
 $= 2 \left(\frac{a^7}{7} + \frac{3a^6}{6} + \frac{13a^5}{5} + \frac{9a^4}{4} \right)_{a=1} = 7.74$

195 (vii) $F^*(A, a) = D_a(F^*(A; a))|_{a=1} = \frac{d}{da}(a^{64} + 7a^{27} + 10a^8 + 7a^3)$
 $= 64a^{63} + 189a^{26} + 80a^7 + 21a^2 = 354$

196 (viii) $ID(A, a) = I_a(ID(A; a))|_{a=1} = \int_0^a (a^3 + 7a^2 + 10a + 7) d_a$
 $= \left(\frac{a^4}{4} + \frac{7a^3}{3} + \frac{13a^2}{2} + 7a \right)_{a=1} = 14.55$

197

198 **Theorem 3.3.** If A is a molecular graph of Fenofibrate ($C_{20}H_{21}ClO_4$) drug then,

199 $M(A; a, b) = 5ab^3 + 2ab^4 + 4a^2b^2 + 11a^2b^3 + a^2b^4 + 2a^3b^3 + a^3b^4$.

200 **Proof:** The M -Polynomials of A are obtained as follows from equation (1).

$M(A; a, b) = \sum_{n \leq m} k_{n,m} a^n b^m$, where $n, m \in \{1, 2, 3, 4\}$

201 $= k_{1,3} a^1 b^3 + k_{1,4} a^1 b^4 + k_{2,2} a^2 b^2 + k_{2,4} a^2 b^4 + k_{2,3} a^2 b^3 + k_{3,3} a^3 b^3 + k_{3,4} a^3 b^4$
 $= 5ab^3 + 2ab^4 + 4a^2b^2 + 11a^2b^3 + a^2b^4 + 2a^3b^3 + a^3b^4$

202 The polynomials obtained in the previous theorem are now used to evaluate a few indices
 203 for the drug's molecular graph.

204

205 **Theorem 3.4.** Let A be the fenofibrate molecular graph ($C_{20}H_{21}ClO_4$). Then we have,

206 i. $M_1(A) = 126$

207 ii. $M_2(A) = 143$

208 iii. ${}^m M_2(A) = 5.415$

209 iv. $H(A) = 5.492$

210 v. $ISI(A) = 28.264$

211 vi. $AZI(A) = 186.22$

212 vii. $F(A) = 626$

213 viii. $Re M_3(A) = 886$

214 **Proof:** Let $M(A; a, b) = 5ab^3 + 2ab^4 + 4a^2b^2 + 11a^2b^3 + a^2b^4 + 2a^3b^3 + a^3b^4$

215 Using Table 4 we obtained the following results

$$\begin{aligned}
216 \quad (i) \quad M_1(A) &= (D_a + D_b)M(A; a, b) \Big|_{a=b=1} \\
&= 20ab^3 + 10ab^4 + 16a^2b^2 + 55a^2b^3 + 6a^2b^4 + 12a^3b^3 + 7a^3b^4 \Big|_{a=b=1} \\
&= 126
\end{aligned}$$

$$\begin{aligned}
217 \quad (ii) \quad M_2(A) &= (D_a D_b)M(A; a, b) \Big|_{a=b=1} \\
&= 15ab^3 + 8ab^4 + 16a^2b^2 + 66a^2b^3 + 8a^2b^4 + 18a^3b^3 + 12a^3b^4 \Big|_{a=b=1} \\
&= 143
\end{aligned}$$

$$\begin{aligned}
218 \quad (iii) \quad {}^m M_2(A) &= (S_a S_b)M(A; a, b) \Big|_{a=b=1} \\
&= \frac{5ab^3}{3} + \frac{2ab^4}{4} + \frac{4a^2b^2}{4} + \frac{11a^2b^3}{6} + \frac{a^2b^4}{8} + \frac{2a^3b^3}{9} + \frac{a^3b^4}{12} \Big|_{a=b=1} \\
&= 5.415
\end{aligned}$$

$$\begin{aligned}
219 \quad (iv) \quad H(A) &= 2S_a J(M(A; a, b)) \Big|_{a=1} \\
J(f(a, b)) &= a^7 + 3a^6 + 13a^5 + 9a^4 \\
2S_a J(M(A; a, b)) &= 2 \int_0^a \frac{a^7 + 3a^6 + 13a^5 + 9a^4}{x} d_a \\
&= 2 \left(\frac{a^7}{7} + \frac{3a^6}{6} + \frac{13a^5}{5} + \frac{9a^4}{4} \right) \Big|_{a=1} = 2(5.492) = 10.984
\end{aligned}$$

$$\begin{aligned}
220 \quad (v) \quad ISI(A) &= S_a J(D_a D_b)M(A; a, b) \Big|_{a=b=1} \\
&= 15ab^3 + 8ab^4 + 16a^2b^2 + 66a^2b^3 + 8a^2b^4 + 18a^3b^3 + 12a^3b^4 \Big|_{a=b=1} \\
&= 15a^4 + 8a^5 + 16a^4 + 66a^5 + 8a^6 + 18a^6 + 12a^7 \\
&= \left(\frac{12a^7}{7} + \frac{26a^7}{6} + \frac{74a^5}{5} + \frac{31a^4}{4} \right) \Big|_{a=1} \\
&= 28.594
\end{aligned}$$

$$\begin{aligned}
221 \quad (vi) \quad AZI(A) &= S_a {}^3 Q_{-2} J(D_a {}^3 D_b {}^3)M(A; a, b) \Big|_{a=b=1} \\
(D_a {}^3 D_b {}^3)M(A; a, b) &= 135ab^3 + 128ab^4 + 256a^2b^2 + 2376a^2b^3 + 512a^2b^4 + 1458a^3b^3 + 1728a^3b^4 \\
&= 1728a^7 + 1970b^6 + 2504b^5 + 391b^4 \\
{}^3 Q_{-2} J(D_a {}^3 D_b {}^3)M(A; a, b) &= 1728a^5 + 1970a^4 + 2504a^3 + 391a^2 \\
&= \left(\frac{1728a^5}{125} + \frac{1970a^4}{64} + \frac{2504a^3}{27} + \frac{391a^2}{8} \right) \Big|_{a=1} \\
AZI(A) &= 186.22
\end{aligned}$$

$$\begin{aligned}
222 \quad (vii) \quad F(A) &= (D_a {}^2 + D_b {}^2)M(A; a, b) \Big|_{a=b=1} \\
223 \quad &= 80ab^3 + 50ab^4 + 64a^2b^2 + 275a^2b^3 + 36a^2b^4 + 72a^3b^3 + 49a^3b^4 \Big|_{a=b=1} \\
&= 626
\end{aligned}$$

(viii) $Re M_3(A) = (D_a + D_b)(D_a D_b)M(A; a, b) |_{a=b=1}$
 224 $= 60ab^3 + 40ab^4 + 216a^2b^2 + 330a^2b^3 + 48a^2b^4 + 108a^3b^3 + 84a^3b^4 |_{a=b=1}$
 225 $= 886$

225 The multiplicative Topological indices of fenofibrate are obtained in the following manner

226

227 **Theorem 3.5.** For a chemical graph of Fenofibrate ($C_{20}H_{21}ClO_4$) drug A, we obtain,


- 228 (i) $NK(A) = 52$
 229 (ii) $\prod_1(A) = 126$
 230 (iii) $\prod_1^*(A) = 126$
 231 (iv) $H\prod_1(A) = 626$
 232 (v) $H\prod_2(A) = 907$
 233 (vi) $SC\prod(A) = 0.089$
 234 (vii) $PC\prod(A) = 0.083$
 235 (viii) $SF\prod(A) = 7.423$
 236 (ix) $PF\prod(A) = 5.418$
 237 (x) $F_1\prod(A) = 340$
 238 (xi) $F_2\prod(A) = 907$
 239 (xii) $ABC\prod(A) = 12.726$
 240 (xiii) $H(A) = 0.015$
 241 (xiv) $GAI(A) = 0.189$

242 **Proof:** By using the degrees of the vertices d_r, d_s of vertices r and s and the values


243 $j_1 = 7, j_2 = 10, j_3 = 7, j_4 = 1$

244 $k_{1,3} = 5, k_{1,4} = 2, k_{2,2} = 4, k_{2,3} = 11, k_{3,3} = 2, k_{2,4} = 1, k_{3,4} = 1,$

245 We are able to compute multiplicative Topological indices using the formulas shown in Table
 246 2.




(i)




(ii)

250
 259
 260

261
262




(iii)




(iv)

263
264

265




(v)




(vi)

277
278
279



(vii)



(viii)

281
282

283

284 Figure 3. Plotting (i) $M_1(A)$, (ii) $M_2(A)$, (iii) $M_3(A)$, (iv) $F(A)$, (v) $M_1 * (A)$ (vi)

285 $F^* A$, (vii) $H(A)$ (viii) $ID(A)$

286

287

288

289
290

291
292
293
294
295
296
297
298
299
300
301
302
303
304




Figure 4. M-Polynomial

305
306
307
308
309
310
311

The graph in Figure 3 illustrates the different Fenofibrate drug polynomials, which are mostly self-explanatory. In Figure 4, the M-polynomial of the medication is plotted in three dimensions.

312
313

4. CONCLUSION

314
315
316
317
318
319
320

Our analysis of many significant sums and the product connectivity topological index for the Fenofibrate drug's molecular graph is being provided numerically for the first time. Using graphical representations, the Zagreb, Harmonic, Forgotten, and Inverse degree polynomials were also calculated. Additionally, it was successful to commence the closed version of Fenofibrate's M-polynomials, and precise formulas were found. The calculated indices may have uses in the pharmaceutical sciences, including data mining, chemical documentation investigations, and the field of medication design.

321

322
323

REFERENCES

324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340

1. Gutman, I.; Trinajstić, N. Graph theory and molecular orbitals. Total ϕ -electron energy of alternanhydrocarbons. *Chemical Physics Letters* **1972**, *17*, 535-538, [https://doi.org/10.1016/0009-2614\(72\)85099-1](https://doi.org/10.1016/0009-2614(72)85099-1).
2. I. Gutman, E. Milovanovic, and I. Milovanovic, "Beyond the Zagreb indices," *AKCE International Journal of Graphs and Combinatorics*, vol. 17, no. 1, pp. 74–85, **2020**.
3. Nikolić, S.; Kovačević, G.; Miličević, A.; Trinajstić, N. The Zagreb Indices 30 Years After. *CroaticaChemicaActa* **2003**, *76*, 113-124.
4. V R Kulli, Niranjana Km, Venkanagouda M Goudar, Degree Based Topological Indices Of Fenofibrate. *Eur.chem.bull.* **2023**, *12*(special issue 8) 6057-6062.
5. M. Kamran, N. Salamat, R. H. Khan, U. ur Rehman Asghar, M. A. Alam, and M. K. Pandit, "Computation of M-polynomial and topological indices of phenol formaldehyde," *Journal of Chemistry*, vol. 2022, Article ID 8655347, 11 pages, **2022**.

- 341 6. Dearden, J.C. The Use of Topological Indices in QSAR and QSPR Modeling. Advance
342 in QSAR Modeling, *Challenges and Advances in Computational Chemistry and Physics*
343 **2017**, 24, 1–555, <https://doi.org/10.1007/978-3-319-56850-8>.
344
- 345 7. Iranmanesh, M.A.; Saheli, M. On the harmonic index and harmonic polynomial of
346 Caterpillars with diameter four. *Iran. J. Math. Chem.* **2014**, 5, 35–43,
347 <https://dx.doi.org/10.22052/ijmc.2015.9044>.
348
- 349 8. S. Delen, R. H. Khan, M. Kamran et al., “Ve-degree, ev-degree, and degree-based
350 topological indices of fenofibrate,” *Journal of Mathematics*, vol. 2022, Article ID 4477808,
351 6 pages, 2022..
352
- 353 9. Deutsch, E.; Kavzar, S. M-polynomial and degree-based topological indices. *Iran*
354 *Journal of Mathematical Chemistry* **2015**, 6, 93–102.
355
- 356 10. Mumtaz, H.B.; Javaid, M.; Awais, H.M; Bonyah, E. Topological Indices of Pent-
357 Heptagonal Nanosheets via M-Polynomials. *Journal of Mathematics*, **2021**, 2021, 1–13,
358 <https://doi.org/10.1155/2021/4863993>.
359
- 360 11. Irfan, M.; Rehman, H. U.; Almusawa, H.; Rasheed, S.; Baloch, I. A. M-polynomials and
361 topological indices for line graphs of chain silicate network and h-naphtalenic
362 nanotubes, *Journal of Mathematics*, **2021**, 2021, 1–11,
363 <https://doi.org/10.1155/2021/5551825>.
364
- 365 12. Raza Z.; Sukaiti, E. K. M-polynomial and degree based topological indices of some
366 nanostructures, *Symmetry*, **2020**, 12, 1–16, <https://doi.org/10.3390/sym12050831>.
367
- 368 13. Khalaf, A. J. M.; Hussain, S.; Afzal, D.; Afzal, F.; Maqbool, A. M-polynomial and
369 Topological indices of book graph, *Journal of Discrete Mathematical Sciences and*
370 *Cryptography*, **2020**, 23,