

## Studies on insect pest succession and their natural enemies in brinjal

### Abstract:

The Present investigation was carried out at College of Agriculture, Vijayapura, Karnataka to study insect - pests succession and their natural enemies on brinjal during 2021-22. Studies revealed that, four species of insect pests and one coccinellid predator were observed at different crop growth stages. The first attack on the crop appeared at one week after transplantation and continued up to till crop reached fruiting and maturity stage. Pests that found attacking on the crop were leaf hopper (*Amrasca biguttula biguttula* Ishida), aphids (*Aphis gossypii* Glover), whitefly (*Bemisia tabaci* Gennadius), shoot and fruit borer (*Leucinodes orbonalis* Guenee) and coccinellid predator, *Cheliomenes sexmaculata* (Fabricius). Brinjal shoot and fruit borer (*L. orbonalis*) was identified as a major pest among them. From the seedling stage to fruiting and maturity stage, the populations of Jassids, aphids, coccinellids were observed, white flies were observed from vegetative stage to fruiting and maturity stage, while the population of shoot and fruit borer were observed and recorded during the vegetative stage to fruiting and maturity stage of the crop. Leaf hoppers, aphids and whitefly were found to damage the crop moderately and extent of damage caused by *L. orbonalis* was created much economic loss.

**Key Words:** Damage, natural enemies, *Leucinodes orbonalis*, pest succession, *Solanum melongena*,

### Introduction

“Brinjal (*Solanum melongena* L.) belongs to the family Solanaceae is one of the most important vegetable crop which grown in South-East Asia where hot and wet climate condition prevails” (Thapa 2010). “India is the second largest producer of vegetables in the world, next to the China. Brinjal is the most consuming vegetable among the vegetarian people. It is one of the main sources of cash crop for many farmers” (Daniel Miller., 2007)

“After China, India is second in terms of area (730,4000 ha), production (12800.8 metric tonnes), and productivity (17.5 metric tons/ha). Bihar, Orissa, Karnataka, Andhra

Pradesh, Maharashtra, West Bengal, and Uttar Pradesh are the major states in India where brinjal is grown. It is grown on 7.1 lakh ha in India, with a yield of 19.1 MT/ha and a production of 135.58 lakh MT. Among the states that cultivate brinjal, Bihar produces the most, with a net sown area of 57,500 ha with the production of 12.40 lakh MT, and 21.6 MT/ha productivity. It is cultivated in Madhya Pradesh over an area of 51,35 thousand hectares, yielding 1073.63 metric tonnes”(Anon., 2018).

Brinjal is a long season crop, it has been attacked by many insect pests during different stages of the crop growth. The information about the incidence of insect pests on crop during different stages at different weather conditions is scanty. Brinjal is most popular vegetable in northern dry zone of Karnataka.

“Brinjal is also known as ‘Poor man’s vegetable’ since it is available at reasonable prices. Now a days cultivation is becoming a menace to the farmer because of attack of insect pests causing damage to the crop from seedling stage upto its maturity. Various pests are observed from seedling to harvesting stage and the loss caused by brinjal pests vary from season to season depending on environmental factors” (Anjali *et al.*, 2012).

Gangawaret *al.* (2014) investigated on “brinjal pests during *Kharif* and revealed that, a total of eight insect species were found associated with brinjal crop at different crop growth stages. The first attack on the crop appeared one week after transplantation and continued till crop harvested. Pests found attacking on the crop were jassids (*Amrascabiguttulabiguttula*), aphids (*Aphis gossypii*), whitefly (*Bemisiatabaci*), leaf roller (*Eublemmaolivacae*), shoot and fruit borer (*Leucinodesorbonalis*), epilachna beetle (*Epilachnaviginioctopunctata*), leaf webber (*Psara bipunctalis*) and grass hopper (*Chrotogonus spp.*)”. “Among them, brinjal shoot and fruit borer (*L. orbonalis*) was recorded as major pest. Jassids (*Amrascabiguttulabiguttula* Ishida), aphid (*Aphis gossypii* Glover.) and epilachna beetle (*E. viginioctopunctata*) were found to damage the crop moderately”. [17]

Nasif and Siddiquee (2020) reported on “the succession of different insect pests and among different insect pests revealed that, brinjal shoot and fruit borer was the major insect pest causing significant reduction in economic yield followed by epilachna beetle, aphid, jassids and whitefly”.

“The first attack of pests on the crop appeared in 1<sup>st</sup> week after transplantation and continued till crop harvested. Pests which were found attacking on the crop were jassids (*Amrascabiaguttulabiguttula*, Ishida), aphids (*Aphis gossypii*, Glover), white fly (*Bemisiatabaci*, Gennadius), Leaf roller (*Eublemmaolivacae*Walker), Shoot and fruit borer (*Leucinodesorbonalis*, Guenee), Epilachna beetle (*Epilachnavigintioctopunctata*, Fabricius), Leaf webber (*Psara bipunctalis*, Fabricius) and Grass hopper (*Chrotogonusspp*Blanchard). Among them, brinjal shoot and fruit borer (*L. orbonalis*) was recorded as major pest. Jassids (*A. biguttulabiguttula*), aphid (*A. gossypii*) and epilachna beetle (*E. vigintioctopunctata*) were found to damage the crop moderately. Other insect pests recorded on the crop were of less importance and extent of damage caused by them was found without much economic loss” (Soren *et al.*, 2020).

## 2. Material and methods

The studies on “Insect pest succession and their natural enemies on brinjal during 2021-22.” were conducted on brinjal hybrid ‘Super mahyco - 10’ at College of Agriculture, Vijayapura, Karnataka, India. The plot size was 46.08m<sup>2</sup> with 120 cm x 60 cm spacing. Randomly ten selected plants forming representative samples were tagged and observation on population of following insect pests of brinjal were recorded (Plate 1) at weekly interval right from transplanting till harvest of crop by adopting standard operational procedures. The extent of damage caused by various insect pests was recorded to assess the economic status of the pests. Observations were also recorded on natural enemies (Plate 2) at weekly interval from 10 randomly selected plants. The Insect- pests and natural enemies collected were preserved and identified by expert taxonomists and later their population was correlated with weather parameters.

### 2.1.1 Observations recorded

1. **White fly** – Number of Adults / Nymphs per three leaves (Each from top, middle and bottom of the plant)
2. **Hoppers** - Number of Adults / Nymphs per three leaves (Each from top, middle and bottom of the plant.)

3. **Aphids** - Number of Adults / Nymphs per three leaves (Each from top, middle and bottom of the plant.)
4. **Fruit and shoot borer** – Percentage shoot and fruit damage.
5. **Natural enemies** - Adults and grubs of predators were enumerated and expressed as a mean number per plant from ten randomly selected plants.

### 3. Results and Discussion:

#### **Leaf hopper, *Amrascabiguttulabigutulla*(Ishida) (Hemiptera: Cicadellidae)**

In current study leaf hopper occurrence was first noted on the crop at seven days after transplanting, and it persisted until the crop has reached reproductive stage (fruiting stage) *i.e* 46<sup>th</sup> SMW to 13<sup>th</sup> SMW (Nov 7<sup>th</sup> - Apr 4<sup>th</sup>) (Table 1).

Both nymphs and adults suck the sap from lower surface of leaves. The infested leaf curl upward along the margins, which turn yellowish and show burnt up patches. Fruit setting is adversely affected by the infestation. In the later stages reduction in size of leaves, conversion of floral parts into leafy structures is observed, plants become bushy, fruiting is rare are the symptoms of 'little leaf of brinjal' transmitted by leaf hopper.

Present findings are in accordance with Gangwar and Singh (2014) and Sheojat *et al.* (2022) According to their reports, leaf hopper population started to appear one week after transplanting and keep on increasing as the crop grow. According to Sahu (2019), the jassid was first noticed on the crop at 10 days after transplanting and remained until crop reached maturity.

Thokcham *et al.* (2022) noticed the leaf hopper population when the crop was 30 days old (vegetative stages), they were active from November to March of 2021, and there were heavy populations of jassid with a relative abundance of 37.5 per cent and they persisted until the crop reached maturity.

#### **Aphid, *Aphis gossypii* (Glover) (Hemiptera: Aphididae)**

The aphid population was initially noticed on the crop at seven days after transplanting and it continued to persist through the entire cropping season from 46<sup>th</sup> to

13<sup>th</sup> SMW (Nov 7<sup>th</sup> - Apr 4<sup>th</sup>) (Table 1).

The tiny pests are found in the colonies of hundreds on abaxial leaf surface and tender shoots. Both nymphs and adults suck cell sap from tender leaves, which gradually became curled and fade and ultimately dry up. Sooty moulds developed on honey dews.

The results of the present study are consistent with Ghose *et al.* (2004) where they have reported that *Aphis gossypii* as an important sucking pest of brinjal. Jiaswal (2018) observed the peak infestation of aphid during 35 days after transplanting vegetative stage and found to be continued up to 175 days after transplanting (harvesting stage). Nasif *et al.* (2020) reported that, aphid infestation was seen during November to March and population peaked during February (fruiting stage). According to Sheojet *et al.* (2022) population of aphids reached peak during the first week of January (flowering stage) and the infestation decreased from February to the last week of March (reproductive stage).

#### **Whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae)**

Whitefly activity was first seen on the crop at twenty-one days after transplanting and continued until the crop had reached fruiting and maturity stage (48<sup>th</sup> SMW to 13<sup>th</sup> SMW) (Nov 23<sup>th</sup> - Apr 4<sup>th</sup>) (Table 1).

Nymphs and adults cause severe damage to plants by feeding on sap and the affected leaves turn yellowish, curl downwards and are ultimately shed. Plants show stunted growth, Honey dew excreted by nymphs leads to formation of sooty molds which form black coating on leaves and reduces the photosynthetic activity.

The present findings are in similarity with Patial and Mehta (2008) who reported whitefly as a moderate pest. According to Gangwar and Singh (2014) whitefly is an important pest of brinjal ecosystem. Shrivastava (2016) who conducted a field study on the incidence of whiteflies and reported that pest was first noticed on a crop at 21 DAT during the *rabi* season. Jiaswal (2018) observed the population of whitefly 16 DAT and continued up to second fortnight of March (reproductive stage)

### **Shoot and fruit borer, *Leucinodes orbonalis*(Guenee) (Lepidoptera:Pyraustidae)**

The first shoot damage was observed at 36 days after transplanting *i.e* 50<sup>th</sup> SMW and continued to occur until 10<sup>th</sup> SMW (Dec 7<sup>th</sup> – Mar 7<sup>th</sup>). Fruit damage was recorded between 4<sup>th</sup> SMW and continued until 13<sup>th</sup> SMW (Jan 18<sup>th</sup> – Apr 4<sup>th</sup>) (Table 1).

The incidence of shoot and fruit borer was seen 45 days after transplanting and continued up to fruit harvesting stage. Larva bores into tender shoots resulting in drooping/drying of tips. It also bores into developing fruits and bore hole is plugged with excreta. Damaged fruits are unfit for marketing.

The present findings are in agreement with Nirmali and Saikia (2017) where they reported that *L. orbonalis* as major pest infecting during vegetative and reproductive phase. Nasif *et al.* (2020) revealed that, a significant decline in economic production was caused by the main pest, the brinjal shoot and fruit borer. Sheojat *et al.* (2022) recorded shoot and fruit borer on the crop at 35 days after transplanting (vegetative stage) and it was continued to be active until the last week of April (Fruiting stage). Thokcham *et al.* (2022) observed that, shoot and fruit borer incidence was observed throughout the year mainly during the fruiting stage from December to March, with the highest relative abundance of 49.95 percent.

### **Lady bird beetle, *Cheilomenessexmaculatus*(Fabricius) (Coleoptera: Coccinellidae)**

Lady bird beetle activity was first observed on the crop at seven days after transplanting and persisted through the fruiting and maturity stage (46<sup>th</sup>SMW to 13<sup>th</sup> SMW) (Nov 7<sup>th</sup> - Apr 4<sup>th</sup>)(Table 1).

Its population fluctuated with the population of its aphid prey.

Naik *et al.* (2008) reported the presence of predatory coccinellid beetle, *Cheilomenessexmaculatus* on brinjal. Jaiswal *et al.* (2018) noted lady bird beetle population during 10 DAT which continued up to end of the March 30<sup>th</sup> (reproductive phase). Thokcham *et al.* (2022) revealed that, coccinellid population was observed during vegetative stage of the crop and they are usually seen where the aphid infestation was high and are noticed during Nov to Feb, 2021 and they continue until the fruiting stage of the crop and these observations are in conformity with the findings of the current study.

Brinjal is cultivated round the year, therefore it is very susceptible to be damaged by many insect pests throughout the growth period especially sucking insects which are having many alternative hosts including weeds and by shoot and fruit borer which can survive both in shoots as well as fruits.

#### **4. Conclusion**

On the basis present investigation of study on insect pest succession and their natural enemies of brinjal crop ecosystem concluded that a total of four insect species and one natural enemy found associated with brinjal crop at different crop growth stages. Aphids, leaf hoppers and coccinellids were observed during seedling phase and continued until fruiting and maturity stage. Whereas, white fly and shoot and fruit borer were observed during vegetative stage and continued until fruiting and maturity stage. Among the four insect pests brinjal shoot and fruit borer as the major dominating pest at vegetative as well as fruiting and maturity stage of the crop. ladybird beetle was found to be predominating species predated the aphid at 55 days old crop. The pests are so destructive leading to reduction of brinjal production.



**1A.***Amarscabiguttulabiguttula*(Ishida)



**1B.***Bemesiatabaci*(Gennadius)



**1C. *Aphisgossypii*(Glover)**




**1D. *Leucinodesorbonalis*(Guenee) (Shootdamage)**



**1E. *Leucinodesorbonalis*(Guenee) (Fruitdamage)**

**Plate 1. population of insect pests of brinjal**



**Plate 2. Natural enemies recorded during studies on succession and population dynamics**

Table 1. Succession of major insect pests of brinjal

		Insect pest		Order	Family	Crop age (Days)	Crop stage
		Common name	Scientific name				
46	Nov7-Nov 15	Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae	7	Seedlingstage
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		
		Ladybird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
47	Nov16-Nov 22	Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae	14	Vegetative stage
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		
		Ladybird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
48	Nov23-Nov 29	Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae	21	Vegetative stage
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Ladybird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
49	Nov30-Dec 6	Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae	28	Vegetative stage
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Ladybird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
50	Dec7-Dec 13	Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae	35	Vegetative stage
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		

		Insect pest		Order	Family	Crop age (Days)	Crop stage
		Common name	Scientific name				
		Shoot and fruit borer (Shoot damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
51	Dec14-Dec20	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	42	Reproductive stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Shoot damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Ladybird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
52	Dec21-Dec27	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	49	Reproductive stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Shoot damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
1	Dec28-jan 3	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	56	Reproductive stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		

		Insect pest		Order	Family	Crop age (Days)	Crop stage
		Common name	Scientific name				
		Shoot and fruit borer (Shoot damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
2	Jan 4 –Jan 10	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	63	Reproductive stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		
		Shoot and fruit borer (Shoot damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
3	Jan 11–Jan 17	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	70	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae		
		Shoot and fruit borer (Shoot damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
4	Jan 18–Jan24	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	77	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		

		Insect pest		Order	Family	Crop age (Days)	Crop stage
		Common name	Scientific name				
		Shoot and fruit borer (Shoot and fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
5	Jan 25–Jan31	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	84	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Shoot and fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
6	Feb 1 – Feb 7	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	91	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Shoot and fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
7	Feb 7 – Feb14	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	98	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		

		Insect pest		Order	Family	Crop age (Days)	Crop stage
		Common name	Scientific name				
		Shoot and fruit borer (Shoot and fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
8	Feb 15– Feb 21	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	105	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Shoot and fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
9	Feb 22 – Feb28	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	112	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Shoot and fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
10	Mar1 – Mar 7	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	119	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		

		Insect pest		Order	Family	Crop age (Days)	Crop stage
		Common name	Scientific name				
		Shoot and fruit borer (Shoot and fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
11	Mar8– Mar14	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	126	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
12	Mar15– Mar28	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	133	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Lady bird beetle	<i>Cheilomenessexmaculata</i> (Fabricius).	Coleoptera	Coccinellidae		
13	Mar28– Apr 4	Aphid	<i>Aphis gossypii</i> (Glover)	Hemiptera	Aphididae	140	Fruiting and maturity stage
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		
		Whitefly	<i>Bemesiatabaci</i> (Gennadius)	Hemiptera	Aleyrodidae		
		Shoot and fruit borer (Fruit damage)	<i>Leucinodesorbonalis</i> (Guenn.)	Lepidoptera	Pyraustidae		
		Leaf hopper	<i>Amrascabigutuulabiguttula</i> (Ishida)	Hemiptera	Cicadellidae		

## References:

1. Anjali Mathur, Singh N P, Meena Mahesh, Singh S, 2012, Seasonal incidence and effect of abiotic factor on population dynamics of major insect pests on brinjal crop. *Journal of Environmental Research and Development*, 7(1A): 431-435.
2. Anonymous. Horticultural Statistics at a Glance, 2018, National Horticulture Board, Horticulture Statistics Division, Department of Agriculture, Cooperation & Farmers Welfare, Ministry of Agriculture & Farmers Welfare, Government of India, New Delhi.
3. Chavan SM, Kumar S, Aarwe SS, 2013, Population dynamics and development of suitable pest management module against major insect pests of tomato (*Lycopersicon esculentum*). *Journal of Applied Horticulture*, 15(2):150-155
4. Daniel Miller. Genetically engineered eggplant. Span 41., 2007
5. Gangwar R K and Singh D V, 2014, Study on insect pest succession of brinjal crop ecosystem in Western Region of Uttar Pradesh. *Indian Journal Agricultural Sciences*, 4(17): 116-119.
6. Ghosh SK, Laskar N, Basak SN, Senapati SK. Seasonal fluctuation of Bemisia tabaci Genn. On brinjal and field evaluation of some pesticides against Bemisia tabaci under terai region of West Bengal. *Environ. Eco.* 2004; 22(4):758-762
7. Jaiswal S K, Dhingra M R, Kumar Akhilesh B H and Kaushik U, 2018, Incidence of Insect Pest in Brinjal under Agro-Climatic Condition of Rewa District, Madhya Pradesh, India. *International Journal of Current & Microbiology Applied Science*, 7(06): 1241-1249.
8. Naik V C B, Rao P A, Krishnayya P V and Rao V S, 2008, Seasonal incidence and management of *Leucinodes orbonalis* (Guenee) on brinjal. *Annals of Plant Protection sciences*, 16: 329-332.
9. Nasif O and Siddiquee S, 2020, Host preference, mode of damage and succession of major insect pests of brinjal, *Annual Research and Review in Biology*, 35(8): 68-78.
10. Patial A and Mehta P K, 2008, Pest complex of brinjal and their succession under mid hills of Himachal Pradesh. *Journal of Insect Science*, 21(1): 67-71.
11. Sahu A, 2019, Study on succession of insect pests on *Solanum melongena* L. and evaluation of insecticides against *Leucinodes orbonalis* Guen. *M.Sc. (Agri.) Thesis* submitted to JNKVV Jabalpur (MP) India.

12. Sheojat N K S, Kumar N, Naveen A C and Suman S, 2022, Studies on the succession and incidence of major insect pests of Brinjal, *Solanum melongena* Linnaeus in Gwalior (MP), *The Pharma Innovation Journal*, 11(7): 1560-1565.
13. Shrivastava P, 2016, Studies on seasonal incidence of insect pest complex of brinjal and their management with chemical and bio-rational insecticides. *M.Sc. (Agri.) Thesis* submitted to JNKVV Jabalpur (MP) India.
14. Soren A, Chakravarty M K, Singh P K, Kudada N, Kumari A and Pandey C, 2020, Study on the succession of insect pests of brinjal. *Journal of Entomology and Zoology Studies*, 8: 1035-1037.
15. Thapa RB, Integrated management of brinjal fruit and shoot borer *Leucinodes orbonalis* Guen. An overview. *Journal of Agriculture and Animal Science*. 2010; 30(32):1-16
16. Thokchom S, Akoijam R, Ningombam A, Bharteey P K and Erla S, 2022, Succession of insect pests and natural enemies in brinjal in Manipur, North East India with bio-pesticide application, *Agricultural Mechanisation in Asia*, 53 (07).
17. Chaukikar K, Vaishampayan S, Marabi RS. The succession of insect pest complex on brinjal at central Narmada Valley region (Madhya Pradesh). *J. Entomol. Zool. Stud.* 2020;8:1757-61.