

Original Research Article

Variability study and DUS characterization in kodo millet (*Paspalum scrobiculatum* L.)

Comment [BA1]: Variability

Abstract :

Comment [BA2]: Abstract:

Aims: To study variability and DUS characterization for yield and its attributing traits

Comment [BA3]: Its

Study Design: Randomized Block Design

Place and Duration of Study: 49 genotypes of kodo millet, were evaluated at the Experimental Farm of Department of Genetics and Plant Breeding, B. A. College of Agriculture, Anand Agricultural University, Anand, during *kharif*, 2021.

Comment [BA4]: The Department

Methodology: The experimental material consisted of 49 different kodo millet genotypes procured from Hill Millet Research Station, Muvaliya Farm, Anand Agricultural University, Dahod, Gujarat.

Results: Qualitative traits offered considerable variability and differentiated the genotypes distinctly but genotypes did not vary much for characters *viz.*, internode pigmentation, spike curvature and spike branching. No co-linearity was found between pigmentation at leaf sheath, leaf juncture, internode or leaf blade, indicating involvement of different loci for pigmentation at these different plant parts. The genotypic and phenotypic variance were high ($>>40$) for plant height followed by fodder yield per plant, total carbohydrate and days to maturity. High GCV and PCV were recorded for total phenol, peduncle length and grain yield per plant, whereas Low GCV and PCV were recorded for all other traits including days to 50% flowering. High heritability along with high genetic advance as percent of mean was observed for grain yield per plant, flag leaf blade length, raceme length, peduncle length, total carbohydrate and total phenol. This indicated involvement of additive genetic control in expression of those traits, hence the traits can be continued for selection in segregating generations.

Comment [BA5]: Little bit need to improve the grammar in the abstract.

Conclusion : Qualitative characters differentiated the genotypes distinctly except pigmentation at leaf sheath, spike curvature and spike branching because genotypes did not vary much for this characters and estimates of GCV were lower than PCV for all the characters which indicated interaction of genotypes with environments.

Comment [BA6]: Conclusion:

Key words: Kodo millet, DUS, Heritability, Genetic advance as % of mean

1. INTRODUCTION:

Kodo millet (*Paspalum scrobiculatum* L.) is a small grained tetraploid ($2n=4x=40$) species, also known as varagu, kodo, haraka, arakalu, cow grass, rice grass, ditch millet, native paspalum, or Indian crown grass and belonging to the family Poaceae (Gramineae). The crop originates in tropical Africa, and domesticated in India 3000 years ago (Upadhyaya *et al.*, 2016).

In kodo millet the percentage of open flowers does not exceed (15-20)% and show cleistogamy condition and thereby self-pollination is the rule in kodo millet (Galinato, 1999; Yadava and Jain, 2006).

This crop is ideal for rainfed agriculture due to its ease of cultivation, low risk of pests and diseases, broad adaptability, and resistance to drought. It is an extremely resilient crop that can thrive on marginal soils also (Heuze *et al.*, 2012).

In India small millets production is highest in Rajasthan followed by Karnataka, Madhya Pradesh, and Uttar Pradesh. In India kodo millet has the highest productivity among the small millets. It occupies an area of 9.08 lakh ha with an annual production of 3.11 lakh tones and average productivity of 342 kg/ha (Anonymous, 2022). It is mainly grown in the states of Madhya Pradesh, Chhattisgarh, Gujarat, Karnataka and parts of Tamil Nadu as a major food source. Madhya Pradesh ranks first in area of kodo millet cultivation with 71.7% share in the country (Makadia *et al.*, 2012). In Gujarat small millets are mainly grown in hilly and tribal regions.

In the post-green revolution era area under millets cultivation has been declining due to the dominance of rice and wheat. But nowadays millets cultivation is increasing as millets provide regional food security in the dry and marginal lands, as well as nutritional superiority as compared to the major cereals (Kumar *et al.*, 2017). The United Nations (UN) designated 2023 as "International Year of Millets" and the intention behind this is to create awareness among peoples and increase production and productivity of small millets.

The genetic and heritable components are expressed by the relative magnitude of the genotypic coefficient of variation (GCV), whereas the environmental and genetic influences on a character are expressed by the phenotypic coefficient of variation (PCV). Heritability estimates by themselves are not very useful for selection based on phenotypic performance. Genotypic coefficient of variation (GCV) along with heritability estimates provides a better understanding of the genetic gain to be expected by phenotypic selection. Moreover, heritability along with genetic advance as percent of mean are more useful in predicting gain under selection. Genetic advance refers to the enhancement in the mean genotypic value of selected individuals over the parental population and understanding the type of gene action involved in polygenic traits expression (Rahikaben and Madhubala, 2020).

Assessment of genetic variability is necessary for any crop improvement programme and because hybridization is restricted as in self-pollinated crops so highly variable germplasm offers much scope for improvement.

Comment [BA7]: Self-pollinated.

Testing of experimental material for distinctness, uniformity and stability (DUS) is useful for identification and classification of varieties as well as for registration. Descriptors for different species have been specified by Protection of Plant Varieties and Farmers' Rights Authority in India.

2. MATERIALS AND METHODS :

A total of 49 different genotypes of kodo millet was collected from Hill millet research station, Anand Agricultural University, Dahod (Table 1).

Comment [BA8]: Were

Table 1 List of genotypes used in the present study

Sr. No	Genotypes	Sr. No	Genotypes	Sr. No	Genotypes
1	ERP-78	18	RPS-811	35	RPS-697
2	RPS-712	19	RPS-824	36	RPS-925
3	RPS-754	20	RPS-648	37	ELB-80
4	RPS-589	21	RPS-882	38	GK-4
5	RPS-745	22	RPS-974	39	RPS-861
6	ERP-49	23	RPS-727	40	RPS-510
7	RPS-560	24	ERP-77	41	GAK-3
8	RPS-903	25	RPS-689	42	RPS-533
9	RPS-685	26	RPS-866	43	ERP-55
10	RPS-1004	27	ELB-56	44	ELB-103
11	RPS-981	28	RPS-310	45	ELB-76
12	ELB-77	29	RPS-771	46	ELB-109
13	RPS-623	30	RPS-802	47	RPS-784
14	RPS-661	31	RPS-794	48	RPS-599
15	RPS-584	32	ELSB-77	49	ELSG-22
16	ELB-61	33	RPS-612		
17	ELB-89	34	RPS-877		

The material was planted in randomized complete block design with three replications at the Experimental Farm of Department of Genetics & Plant Breeding,

B. A. College of Agriculture, Anand Agricultural University, Anand, Gujarat during *kharif*, 2021 with a spacing of 30 × 10 cm. All the recommended package of practices was followed to raise a good crop.

A total of 16 different quantitative parameters, namely, number of basal tillers per plant, days to 50% flowering, flag leaf blade length (cm), flag leaf blade width (cm), peduncle length (cm), number of productive tillers per plant, panicle length (cm), plant height (cm), days to maturity, 1000 seed weight (g), grain yield per plant (g), fodder yield per plant (g), harvest index (%) including three biochemical parameters, viz., crude protein content (%), total carbohydrate content (%), total phenol content (%).

Analysis of variance (ANOVA) was carried out following Panse and Sukhatme (1978), Genotypic (σ^2_g), phenotypic (σ^2_p) and environmental variance (σ^2_e) were calculated as per the formulae suggested by Johnson *et al.* (1955). GCV and PCV were computed according to the method suggested by Burton (1952). Heritability in broad sense (H^2_b) and genetic advance (as % of mean) was calculated according to

Allard *et al.* (1960) and Johnson *et al.*, (1955), respectively. Traits were classified as having high, moderate or low genetic advance as per the method suggested by Johnson *et al.*, (1955).

3. RESULTS AND DISCUSSION:

Estimates of variance components are presented in Table 2. Genotypic variances (σ^2g) and estimates of GCV were found lower than phenotypic variances (σ^2p) and PCV respectively, for all the traits indicating influence of environment in the expression of the traits (Table 2).

Comment [BA9]: An influence

High GCV and PCV ($\gg 20.0$) were recorded for total phenol, peduncle length and grain yield per plant. Similar results were observed by Thakur (2018) and Sreeja *et al.* (2014) in kodo millet.

Moderate GCV and PCV (10.1 - 20.0) were recorded for fodder yield per plant, culm branching, flag leaf blade length, harvest index, total carbohydrate content, raceme length, panicle length, plant height and crude protein content.

Low GCV and PCV were recorded for other characters including days to 50% flowering, days to maturity, number of basal tillers per plant and number of productive tillers per plant indicating narrow range of existing variation for these traits. Similar findings in kodo millet were reported by Sao *et al.* (2017).

Highest magnitude of heritability was observed for total phenol followed by flag leaf blade length, peduncle length, 1000 seed weight and total carbohydrate content indicating that these characters are least affected by environment. The result is supported by similar findings from Thakur (2018) and Anuradha *et al.* (2020).

Comment [BA10]: The highest

Low heritability was observed for number of raceme per tiller (78.28%), number of productive tillers per plant, culm branching and least was observed for number of basal tillers per plant.

Comment [BA11]: Racemes

High heritability (H^2b) along with high genetic advance as percent of mean (GA as % of mean) was observed for grain yield per plant, flag leaf blade length, peduncle length, raceme length, total carbohydrate and total phenol, which indicated involvement of additive gene action in control of those traits, hence for selection of genotype based on the traits can be carried out.

Similar results were observed by Sreeja *et al.* (2014), Sao *et al.* (2017), Nirubana *et al.* (2017), Thakur (2018) and Anuradha *et al.* (2020) in kodo millet.

Table 2 Estimates of genetic parameters

Sr. No.	Characters	σ^2g	σ^2p	GCV %	PCV %	Heritability (%)	GA as % mean
1	Number of basal tillers per plant	0.490	0.848	5.129	6.745	57.83	8.036
2	Days to 50% flowering	12.183	14.978	4.902	5.435	81.34	9.108
3	Flag leaf blade length (cm)	18.850	19.103	17.384	17.500	98.67	35.573
4	Flag leaf blade width (cm)	0.007	0.008	8.821	9.765	81.61	16.415
5	Peduncle length (cm)	2.889	2.929	25.834	26.013	98.63	52.852
6	Number of productive tiller per	0.405	0.539	8.425	9.717	75.17	15.047

	plant						
7	Culm branching	0.298	0.462	16.554	20.608	64.53	27.393
8	Panicle length (cm)	0.844	0.886	13.489	13.820	95.26	27.121
9	Raceme length (cm)	1.011	1.051	15.847	16.159	96.18	32.014
10	Number of raceme per tiller	0.105	0.134	9.121	10.399	78.28	16.624
11	Plant height (cm)	75.148	76.980	12.288	12.437	97.62	25.011
12	Days to maturity	44.307	48.588	5.993	6.276	91.19	11.790
13	1000 seed weight (g)	0.264	0.269	10.945	11.056	98.00	22.320
14	Grain yield per plant (g)	18.912	19.787	24.257	24.812	95.58	48.852
15	Fodder yield per plant (g)	60.422	72.075	17.447	19.055	83.84	32.910
16	Harvest index (%)	22.176	25.720	16.391	17.652	86.22	31.353
17	Total carbohydrate (%)	62.693	64.028	16.124	16.295	97.91	32.868
18	Crude protein (%)	1.726	1.823	11.767	12.096	94.63	23.581
19	Total Phenol (%)	0.002	0.002	40.470	40.470	99.10	83.369

σ^2_g = Genotypic variance σ^2_p = Phenotypic variance GCV= Genotypic coefficients of variation PCV= Phenotypic coefficients of variation (H^2_b) % = Broad sense heritability % G.A as % of mean = Genetic advance as % of mean

DUS Characterization:

The performance of 49 genotypes for different 18 qualitative and 11 quantitative characters as per DUS guidelines are summarized in Table 3.

No co-linearity was found between pigmentation at leaf sheath, leaf juncture, internode or leaf blade, indicating involvement of different loci for pigmentation at these different plant parts. Much variation was observed for genotype performance for the characters viz., plant growth habit, leaf attitude, leaf juncture pigmentation, internode pigmentation, leaf blade pigmentation, panicle appearance, panicle exertion, spikelet arrangement on rachis, intensity of irregular rows in spikelet, spikelet density, appearance of glumes nerves, lodging, seed shattering, grain colour and grain shape, while, genotypes did not vary much for characters viz., internode pigmentation, spike curvature and spike branching.

Similar observation for characters, viz., Leaf juncture pigmentation, internode pigmentation, leaf blade pigmentation, spike curvature, panicle appearance, seed shattering, grain colour and grain shape was recorded by Nirubana *et al.* (2019) in kodo millet.

Considering all traits it can be suggested that genotypes, viz., RPS-824, RPS-882, RPS-612 and GK-4 were better performers in terms of yield, its component traits with early to medium flowering.

Table 3 DUS characterization

Sl. no.	Characteristics	States	Example Varieties	Stage of observations	Type of assessme
1 (+)	Plant: Growth habit	Erect	2, 3, 5, 6, 7, 8, 12, 14, 15, 16, 17, 18, 19, 21, 23, 26, 27, 32, 35, 36, 38,40, 41, 42, 43, 44, 45, 47	15	VG
		Decumbent	9,10, 29, 33, 37		
		Prostrate	1, 4, 11, 13, 20, 22, 24, 25, 28, 30,		

Comment [BA12]: In the states portion maintain spaces between the words

			31, 34, 39, 46, 48, 49		
2	Basal tillers: Number	Very low (<10)	-	26	MS
		Low (10-20)	1-49		
		Medium (20.1-30.0)	-		
		High (>30)	-		
3 (* (+)	Leaf : Attitude	Erect	1 to 10,12, 14, 15, 16, 17, 18, 20 to 25, 27, 28, 29, 30, 32, 33, 34, 36 to 41, 43, 45, 46, 48	26	VG
		Droopy	11, 13, 19, 26, 31, 35, 42, 44, 47, 49		
4 (* (+)	Days to 50%flowering	Early(<65)	14, 38	51	MG
		Medium (65-75)	1 to 13, 16 to 27, 29 to 34, 36, 37, 39, 41 to 46, 48		
		Late(75-85)	7, 15, 28, 35, 40, 47, 49		
		Very late(>85)	-		
5 (*	Leaf Sheath: Pigmentation	Absent	3, 6, 7, 8, 9, 12, 16, 17, 18, 20, 22, 23, 25, 26, 27, 28, 29, 35, 38, 39, 40, 41, 42, 48, 49	54	VS
		Present	1, 2, 4, 5, 10, 11, 13, 14, 15, 19, 21, 24, 30, 31, 32, 33, 34, 36, 37, 43, 44, 45, 46, 47		
6 (*	Leaf juncture: Pigmentation	Absent	1, 3, 6, 10, 14, 5, 16, 17, 18, 21, 22, 23, 24, 26, 27, 28, 29, 35, 37, 38, 39, 41, 43, 44, 45, 47, 48, 49	54	VS
		Present	2, 4, 5, 7, 8, 9, 11, 12, 13, 19, 20, 25, 30, 31, 32, 33, 34, 36, 40, 42, 46		
7 (*	Internode: Pigmentation	Absent	-	54	VS
		Present	1 to 49		
8	Leaf blade: Pigmentation	Absent	3, 4, 6 to 19, 21, 22, 24 to 30, 32, 34, 35, 36, 39, 40, 42, 43 to 47, 49	54	VG
		Present	1, 2, 5, 20, 23, 31, 33, 37, 38, 41, 48		
9 (+)	Flag leaf blade: Length (cm)	Short(<20.0)	4, 6, 12, 13, 15, 19, 22, 30	54	MS
		Medium (20.0-30.0)	1, 2, 3, 5, 7 to 11, 14, 16, 18, 20, 21, 23, 24, 25, 26, 27, 28, 29, 31, 32, 34, 36, 37, 40, 41, 42, 43, 45, 46, 47, 48, 49		
		Long(>30.0)	17, 33, 35, 38, 39, 44		
10 (+)	Flag leaf blade: width(cm)	Narrow(<0.5)	-	54	MS
		Medium (0.5-1.0)	1, 3 to 9, 11, 12, 13, 14, 16-19, 22, 23, 25, 27, 30, 31, 32, 36, 37, 39, 41, 42, 43 46, 47, 48, 49		
		Wide(>1.0)	2, 10, 15, 20, 21, 24, 26, 28, 29, 33, 34, 35, 38, 40, 44, 45		
11 (+)	Peduncle: Length (cm)	Short(<5.0)	1, 2, 9, 11, 12, 14, 19, 38, 39	54	MS
		Medium (5.0-10.0)	3, 4, 5, 7, 8, 10, 13, 15, 16, 17, 18, 20 to 34, 36, 37, 40, 41, 42, 43, 44, 46, 47, 49		
		Long(> 10.0)	6, 35, 45, 48		

12 (* (+)	Panicle: Appearance	Compact	-	54	VG
		Semi compact	1, 14, 19, 22, 28, 30, 40		
		Open	2 to 13, 15, 16, 17, 18, 20, 21, 23 to 27, 29, 31 to 39, 41 to 49		
13 (+)	Panicle: Exertion	Partial	4, 5, 6, 7, 9, 10, 11, 15, 16, 19, 20, 21, 26 to 29, 31, 32, 33, 35, 36, 37, 38, 40, 41, 43, 44, 45, 46, 48, 49	54	VS
		Complete	1, 2, 3, 8, 12, 13, 14, 17, 18, 22, 23, 24, 25, 30, 34, 39, 42, 47		
14 (* (+)	Spikelet: Arrangement on rachis	Regular	2, 3, 4, 5, 7 to 13, 15, 16, 17, 18, 20, 21, 23 to 27, 29, 31 to 35, 37, 38, 39, 41, 42, 43, 47, 48	67	VG
		Irregular	1, 6, 14, 19, 22, 28, 30, 36, 40, 44, 45, 46, 49		
15	Spikelet irregular rows: Intensity	Two-three	14	67	VG
		Two-four	6		
		Lower half	1, 19, 22, 28, 30, 36, 40, 44, 45, 46, 49		
16 (+)	Spike: Branching	Absent	1 to 49	67	VG
		Present	-		
17 (+)	Spike: Curvature	Straight	1 to 6, 8 to 20, 22 to 28, 30 to 39, 41 to 44, 46, 47, 48, 49	67	VG
		Curved	7, 21, 29, 40, 45		
18	Spikelet: Density	Lax	2 to 12, 14, 15, 16, 19, 20, 21, 26, 27, 29 to 36, 38, 40, 41, 42, 43, 46, 47, 48,	67	VG
		Dense	1, 13, 17, 18, 22, 23, 24, 25, 28, 37, 39, 44, 45, 49		
19 (+)	Culm: Branching	Low (<3)	2, 9, 12, 18, 19, 21, 25, 26, 30, 32, 38	67	MG
		Medium (3-7)	1, 3, 4, 5, 6, 7, 8, 10, 11, 13, 14, 15, 16, 17, 20, 22, 23, 24, 27, 28, 29, 31, 33, 34, 35, 36, 37, 39, 40, 41 to 49		
		High (>7)	-		
20 (+)	Panicle: Length (cm)	Short (<5.0)	-	77	MS
		Medium (5.1-8.0)	1, 2, 3, 4, 5, 6, 8 to 16, 17, 18-32, 34, 37 to 42, 44 to 48		
		Long (>8.0)	7, 33, 35, 36, 43, 49		
21 (+)	Raceme: Length (cm)	Short (< 5.0)	32, 44	77	MS
		Medium (5.0-10.0)	1 to 31, 33 to 43, 45 to 49		
		Long (10.1-15.0)	-		
		Very long (>15.0)	-		
22 (* (+)	Raceme: Number (Above thumb)	Low (<2)	-	77	MS
		Medium (2-4)	1 to 27, 29-32, 34-43, 45, 47, 48, 49		
		High (4-6)	28, 33, 44, 46		
		Very high (>6.0)	-		
23	Glume: Appearance of	Narrow (7 nerves)	1, 3, 4, 5, 6, 8, 9, 11 to 19, 21, 22, 24, 25, 26, 28, 30 to 34, 36, 37, 39 to 47	77	VG

(*)	Nerves	Broad (5 nerves)	2, 7, 10, 20, 23, 27, 29, 35, 38, 48, 49		
(+)					
24	Plant: Height (cm)	Semi-dwarf (<30.0)	-	83	MS
(*)		Dwarf (30-50)	-		
(+)		Medium (50.1-70.0)	1, 2, 5, 7, 8, 9, 10, 20, 21, 24, 27, 28, 30, 31, 32, 35, 36, 37, 39, 43, 44, 48		
		Tall (>70.0)	3, 4, 6, 11 to 19, 22, 23, 25, 26, 29, 33, 34, 38, 40, 41, 42, 45, 46, 47, 49		
25	Lodging	Absent	1, 2, 21, 22, 31, 35, 40, 41, 42, 45	83	VG
		Present	3 to 20, 23 to 30, 32, 33, 34, 36, 37, 38, 39, 43, 44, 46, 47, 48, 49		
26	Seed: Shattering	Absent	2, 3, 11, 27, 32, 39, 43, 47	83	VG
		Present	1, 4 to 10, 12 to 26, 28, 29, 30, 31, 33 to 38, 40, 41, 42, 44, 45, 46, 48, 49		
27 (*)	Grain: Colour	Light brown	2, 4, 5, 12, 23, 27, 31, 34, 46,	83	VG
		Brown	1,3, 6, 8, 10, 11, 13 to 22, 24, 26, 28, 29, 30, 32, 33, 36 to 42, 44, 45, 47, 48, 49		
		Dark brown	7, 9, 25, 35, 43		
28	Grain: Shape	Elliptical	1, 2, 6, 8, 9, 10, 12, 14, 16, 18, 21, 23, 26, 27, 33, 36, 38, 41, 45	95	VG
		Oval	3, 4, 5, 7, 11, 13, 15, 17, 19, 20, 22, 24, 25, 28, 29, 30, 31, 32, 34, 35, 37, 39, 40, 42, 43, 44, 46, 47, 48, 49		
29 (*)	1000- grain weight (g)	Low (<5.0)	1, 3 to 21, 23, 26 - 30, 32, 33, 34, 36, 37, 38, 39, 40, 45 - 49	95	MG
		Medium (5.0-6.0)	2, 22, 24, 25, 31, 35, 42, 43, 44		
		High (>6.0)	41		

Table 4. Decimal code for the growth stage

Stage code	General Description
15	2-4 Leaf stage
26	Vegetative
51	50 %Flowering
59	Complete flowering
67	Dough stage
77	Seed filling
83	Maturity
95	After harvest

MG: Single measurement of a group of plants or parts of plants.

MS: Measurement of a number of individual plants or parts of plants.

VG: Visual assessment by a single observation of a group of plants or plant parts.

VS: Visual assessment by observation of individual plant or parts of plants.

Comment [BA13]: several

4. CONCLUSION :

Qualitative traits offered considerable variability and differentiated the genotypes distinctly except pigmentation at leaf sheath, spike curvature and spike

branching. Magnitudes of phenotypic variances were higher as compared to genotypic variances for all the traits which suggested influence of environment. Estimates of GCV were lower than PCV for all the characters under study which indicated interaction of genotypes with environments.

Comment [BA14]: An influence

REFERENCES:

- Allard RW, John Willey and Sons Inc. Principles of Plant Breeding, New York. 1960; 85-95
- Anonymous. Department of Agriculture, Corporation and Farmers Welfare. accessed on 28 February 2022. Available at <https://agricoop.nic.in>.
- Anuradha N, Patro TSSK, Triveni U, Rao PJ, Rani YS, Priya PK . Assessment of genetic variability estimates in Kodo millet (*Paspalumscrobiculatum* L.). International Journal of Current Microbiology and Applied Sciences. 2020; 8(5):1907-1909.
- Burton GW. Quantitative inheritance in grasses. Proceedings of 6th International Grassland Congress. 1952; 1: 227-283.
- Galinato MI, Moody K, Piggin CM. Upland rice weeds of South and Southeast Asia. International Rice Research Institute, Manila, Philippines. 1999;156.
- HeuzéV, Tran G, Giger-Reverdin S. Scrobic (*Paspalumscrobiculatum*) forage and grain. 2012.
- Johnson H W, Robinson HF, Comstock RI. Estimates of genetic and environmental variability in soybean. Agronomy Journal. 1955;47:314-318.
- Kumar P, Abhinav Sao AK, Yadav SC, Sahu P. Resourceful photosynthesis system and stem reserve accumulation plays decisive role in grain yields of kodo millet (*Paspalumscrobiculatum*). International Journal of Pure and applied Bioscience. 2017;5(2): 420-426.
- Makadia JJ, Patel K S, Ahir NJ. Cost structure of minor millets grown in tribal dang district of south Gujarat. International Research Journal of Agricultural Economics and Statistics. 2012;3(1): 40-44.
- Nirubana V, Ganesamurthy K, RavikesavanR, Chitdeshwari T. Genetic variability studies for yield and yield components in kodo millet (*Paspalumscrobiculatum* L.). Electronic Journal of Plant Breeding. 2017;8(2):704-707.
- Nirubana V, Ravikesavan R, Ganesamurthy K. Characterization and clustering of kodo millet (*Paspalumscrobiculatum*L.) genotypes based on qualitative characters. Electronic Journal of Plant Breeding. 2019;10(1):101-110.
- PanseVG, Sukhatme PV. Statistical methods for agricultural workers;1978.
- RahikabenPN, Bala M. Genetic variability study for yield and its components in black gram [*Vigna mungo* (L.) Hepper]. Journal of Pharmacognosy and Phytochemistry. 2020;9(4):2061-2064.
- Sao A, Singh P, Kumar P, Pradhan A. Estimates of genetic parameters for yield and contributing traits in kodo millet (*Paspalumscrobiculatum* L.). Research Journal of Agricultural Sciences. 2017;8 (1):120-122.
- Sreeja R, Subramanian A, NirmalakumariA, KannanBapu JR. Selection criteria for culm strength in Kodo millet (*Paspalumscrobiculatum* L.) to suit mechanical harvesting. Electronic Journal of Plant Breeding. 2014;5(3):459-466.
- Thakur V. *Genetic variability and multivariate analysis in kodo millet* (Master's thesis, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur); 2018.

Upadhyaya HD, Vetriventhan M, Dwivedi SL, Singh SK, Pattanashetti SK. "Proso, barnyard, little, and kodo millets," in genetic and genomic resources for grain cereals improvement. *Frontiers in Plant Science*. 2016;321-343.

Yadava HS, Jain AK. Advances in kodo millet research. Directorate of Information and Publications of Agriculture, Indian Council of Agricultural Research. 2006;84-95.

Comment [BA15]: Overall paper need to recheck grammar and proper spaces between the words or in quotation.

UNDER PEER REVIEW