

Emerging Trends in Quinolone Resistance Among Urinary Pathogens: A Brief Review

ABSTRACT

Urinary tract infections (UTIs) present a significant global health challenge, intensified by the high level of antimicrobial resistance, particularly to quinolone antibiotics. This brief review synthesizes recent literature to elucidate the prevalence and mechanisms of quinolone resistance, with a focus on plasmid-mediated quinolone resistance (PMQR), among UTI-causing pathogens. Highlighting the genetic basis of resistance, including mutations in DNA gyrase and efflux pump regulation genes, as well as the role of plasmid-mediated mechanisms such as 'qnr' genes, the review underscores the clinical implications of quinolone resistance in UTI management. Understanding these emerging trends is urgently required for improving antimicrobial management strategies and guiding effective interventions to control the spread of quinolone resistance among urinary pathogens.

Keywords:

- Quinolone resistance
- Urinary pathogens
- Antibiotic resistance
- Urinary tract infections (UTIs)
- Multidrug resistance (MDR)
- Extensively drug-resistant (XDR)
- Pan-drug-resistant (PDR)
- Antimicrobial stewardship
- Public health challenges

INTRODUCTION

Urinary tract infections (UTIs) are prevalent bacterial infections affecting millions globally, posing significant public health challenges (Mancuso et al., 2023; Tan & Chlebicki, 2016). The World Health Organization recognizes UTIs as a widespread concern, impacting morbidity, healthcare costs, and escalating antimicrobial resistance rates (WHO, 2023). Particularly in Asian countries, UTIs are common, with notable antibiotic resistance rates, signaling the emergence of spreading resistant strains (Choe, 2018). With the trend of UTIs becoming increasingly common, there is a pressing need for improved management strategies amidst rising antibiotic resistance. Gram-negative bacteria are the primary cause of UTIs due to their ability to adhere to and

colonize the urinary tract mucosa, thereby initiating infections (Alasmary, 2021). Among antibiotic-resistant clinical pathogens, *E. coli* and *Klebsiella spp* are frequently encountered (PH et al., 2024).

Comment [AS1]: Write the complete species name for the first time.
Escherichia coli

Quinolone antibiotics, such as norfloxacin and nalidixic acid, have historically been primary choices for treatment due to their broad spectrum of activity against uropathogens. However, extensive quinolone use has led to the emergence and spread of resistant strains, diminishing treatment efficacy (Zhou et al., 2023). Quinolones function by inhibiting bacterial DNA replication and repair; however, misuse has resulted in rising resistance, primarily through mutations in DNA gyrase and efflux pump regulation genes (Millanao et al., 2021; Guan et al., 2013). Additionally, plasmid-mediated resistance mechanisms, such as *qnr* genes, contribute to multidrug resistance, complicating treatment processes (Hooper & Jacoby, 2015; Strahilevitz et al., 2009). The escalating severity of quinolone resistance underscores the urgent need to understand the genetic basis of resistance mechanisms in microbes (Soucy, 2018).

Comment [AS2]: give examples please, not all readers are familiar with or have a background in microbiology.
i.e. *K. pneumoniae*, *Pseudomonas spp.*...etc, etc.

This constant change in antimicrobial resistance mechanisms and the lack of new antibiotics to tackle this problem need to be addressed immediately (TR et al., 2024).

This review aims to assess the prevalence of quinolone resistance, particularly plasmid-mediated quinolone resistance (PMQR), in bacteria causing UTIs. A comprehensive search across reputable databases such as PubMed, Google Scholar, and ScienceDirect, as well as relevant scientific journals including those from the American Society of Microbiology (ASM), was conducted to gather pertinent literature. Through this analysis, we endeavor to provide insights into the extent of quinolone resistance in UTI-causing microbes, thereby shedding light on potential implications for clinical management and public health.

Antibiotic resistance in Urinary Tract Infections

Antibiotic resistance presents a significant global public health challenge, particularly pronounced in India. With an infectious disease mortality rate of 416.75 per 100,000 persons in 2016, India faces a pressing concern regarding the spread of resistant pathogens. Various factors, including inadequate public health infrastructure, hospital-acquired infections, high rates of infectious diseases, and easy access to inexpensive antibiotics, contribute to this threat. As a result, there's a growing burden of untreatable conditions like neonatal sepsis and healthcare-associated infections, underscoring the immediate need for effective strategies to combat antibiotic resistance in the country (Laxminarayan & Chaudhury, 2016).

In 2023, Sharma et al. highlighted the critical threat posed by antimicrobial resistance (AMR) to global public health, particularly within India's healthcare system. With its dense population, significant disease burden, and diverse healthcare practices, India stands at a crucial crossroads in combating AMR on a global scale. The consequences of this crisis are far-reaching, impacting healthcare delivery, mortality rates, and the achievement of Sustainable Development Goals. Addressing AMR in India requires a comprehensive approach, including robust regulatory

frameworks, improved diagnostics, heightened public awareness, and research into new antimicrobials.

The study of Hilt et al., in 2023 mentioned the classification of UTIs, introducing newer definitions such as "acute simple cystitis" and "acute complicated UTI" based on the extent of infection and severity of illness. This shift in terminology reflects advancements in clinical practice and antimicrobial uses, emphasizing the importance of newer treatment approaches to individual patient presentations. By acknowledging the limitations of traditional classifications and advocating for more nuanced approaches, the study contributes to ongoing efforts to optimize UTI management and combat antimicrobial resistance.

Quinolone-resistance in urinary pathogens

A study conducted in a tertiary care hospital in Amritsar, Punjab, India, investigated the prevalence and antibiotic resistance patterns of uropathogens among female patients with urinary tract infections (UTIs). *Escherichia coli* emerged as the predominant uropathogen, followed by *Klebsiella spp* and *Enterobacter spp*. The study revealed eight isolates with multi-drug resistance, underscoring the growing challenge of antibiotic resistance. Notably, gentamicin, imipenem, and meropenem showed efficacy against these resistant strains. The detection of the quinolone-resistant gene *qnrB* in all extended-spectrum β -lactamase (ESBL)-positive isolates provided valuable insights into the genetic mechanisms contributing to resistance. These findings emphasize the significance of epidemiological data in guiding empirical antibiotic treatment for UTIs and highlight the importance of ongoing surveillance and prudent antibiotic use in clinical practice.

In 2023 cross-sectional study explored the correlation between biofilm formation and multidrug resistance in uropathogenic *Escherichia coli* (UPEC) strains isolated from urine samples. Results show that 23.92% of UPEC isolates are multidrug-resistant, with 36.06% exhibiting potent biofilm-forming capabilities. These biofilm-producing strains display resistance to commonly used UTI antibiotics but are sensitive to imipenem and meropenem. The study highlights the significance of biofilm formation in antimicrobial resistance and suggests improved strategies for managing biofilm-associated UTIs. Advanced techniques like VITEK R2 Compact and 16S rRNA gene sequencing were used to confirm and characterize the isolated *E. coli* biofilms, contributing valuable insights into addressing multidrug resistance in uropathogenic *E. coli* strains (Cardiliya et al., 2023)

A study examined the prevalence and antibiotic resistance patterns of uropathogens causing urinary tract infections (UTIs). *E. coli* is identified as the primary uropathogen, followed by *Klebsiella species*, confirmed through gram staining, microscopy, and biochemical characterization. Alarming, *E. coli* shows significant resistance to Nitrofurantoin, Ciprofloxacin, and Co-Trimoxazole, resulting in a 36% multidrug resistance (MDR) rate. Extensively drug-resistant (XDR) and pan-drug-resistant (PDR) strains are reported, particularly in *Klebsiella* and *Enterobacter*. *Staphylococcus* exhibits increased resistance, while *Pseudomonas* and *Proteus* show negligible drug resistance. Molecular investigations revealed a high frequency

Comment [AS3]: Please, introduce citations. Each sentence not cited is yours. Overall, there are many sentences without citations. Please, introduce them adequately.

Comment [AS4]: next, just write the species's name i.e. *E. coli*

Comment [AS5]: How about plasmid genes *qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, and *qnrVC*. It would be better to discuss too

of *gyr-A*, *tet-A*, *aac(6')-Ib* genes, and β -lactamase gene presence in the uropathogens. This analysis highlights the growing antibiotic resistance crisis in uropathogens, emphasizing the need for effective strategies to manage drug-resistant UTIs. The findings provide valuable insights for public health interventions and antibiotic stewardship efforts (Gupta et al., 2021).

In a rural area of central India, a study examines the presence of extended-spectrum β -lactamase (ESBL)-producing commensal *Escherichia coli* in healthy school children, highlighting their potential as reservoirs for antibiotic resistance genes within the community. The investigation reveals a significant prevalence of CTXM-15, TEM-1, OXA-1, and SHV-12 genes associated with cephalosporin resistance among these isolates. Additionally, all isolates carry the *aac(6')-Ib-cr* gene, with a subset being *qnrS* positive, indicating plasmid-mediated quinolone resistance. The presence of virulence genes like *fluA*, *fluB*, *eae*, and *daaE* underscores the dual nature of these commensal *E. coli* strains, serving as potential reservoirs for both antibiotic resistance and virulence factors linked to urinary tract and diarrheal infections. The study underscores the necessity for ongoing surveillance and antimicrobial stewardship in communities to mitigate the potential spread of antibiotic resistance among healthy populations (Chandran et al., 2017)

This study conducted whole-genome sequencing (WGS) on 103 carbapenem-resistant *Escherichia coli* (CREc) urinary isolates to analyze fluoroquinolone-resistant determinants. Predominantly, ST410, followed by ST405 and ST361, with Clermont phylogroup C, were identified as the most frequent. The presence of NDM-5 and CTX-M-15 genes significantly contributed to extensive resistance, particularly against ciprofloxacin and levofloxacin. The recurrent coexistence of *aac(6')-Ib* and *blaCTX-M-15* underscored the genetic complexity of these isolates. Additionally, the study explored the complete genomes of five urinary CREc isolates, elucidating the plasmid types harboring *blaNDM-5* or *blaNDM-3* and their association with other resistance genes. These findings highlight the urgency for efficient strategies and control policies to mitigate the spread of fluoroquinolone-resistant CREc strains, offering valuable insights for ongoing efforts in monitoring and combating antimicrobial resistance (Boueroy et al., 2023).

The resistance patterns among *Enterobacteriaceae* isolated from urinary tract infections (UTIs) exhibited High rates of quinolone resistance among ESBL-producing isolates highlighting the increasing challenge of antibiotic resistance in this clinical context. The identification of *PMQR* genes, notably *aac(6')-Ib-cr*, in a significant proportion of samples, adds complexity to the resistance landscape. Associations between *PMQR* genes and *ESBLs*, particularly *blaCTX-M-15* and *blaTEM-116*, underscore the interconnectedness of different resistance mechanisms. These findings stress the urgent need for vigilant antibiotic stewardship and surveillance to control the spread of resistant strains in this region. The study significantly enhances our understanding of resistance gene prevalence and associations in *Enterobacteriaceae* from UTIs (Azargun et al., 2018).

Research conducted in southwest Iran addresses the pressing issue of antibiotic resistance in *Escherichia coli*, as a key opportunistic pathogen causing urinary tract infections (UTIs). The study explores the extent of antibiotic resistance to quinolones and investigates the prevalence of

qnr genes (A, B, and S) in both extended-spectrum beta-lactamase (ESBL) and non-ESBL-producing *E. coli* strains isolated from UTI-diagnosed patients. The findings reveal concerning levels of resistance, particularly against nalidixic acid, highlighting the necessity for robust antibiotic stewardship in Ahvaz's hospitals, Khuzestan province. The identification of *qnr* genes, notably *qnrS*, among ESBL-producing isolates raises alarm about the interconnectedness of various resistance mechanisms. This study contributes valuable insights into the antibiotic resistance landscape of *E. coli* associated with UTIs in southwest Iran, emphasizing the need for further research to comprehensively assess the gravity of quinolone resistance in the country and develop effective strategies against its spread among nosocomial pathogens (FarajzadehSheikh et al., 2019).

Alhazmi et al., (2023) conducted a cross-sectional study in Jazan Province, Kingdom of Saudi Arabia, which reveals the significant impact of urinary tract infections (UTIs) on the healthcare system, constituting 10% of all infections and ranking as the second leading cause of emergency department admissions. With 1082 urinary bacterial samples analyzed, Gram-negative bacteria, particularly *Escherichia coli*, dominate the causative pathogens. Alarmingly, 30.13% of cases exhibit extended-spectrum beta-lactamase (ESBL) resistance, and multidrug-resistant organisms contribute to approximately 35% of reported cases. The study identifies a seasonal pattern, with heightened UTI incidence in September. These findings underscore the urgent need for targeted interventions and antimicrobial stewardship programs to address the prevalence of antibiotic-resistant microbes and mitigate the impact of UTIs.

Another study focused on a thorough analysis of *Escherichia coli* isolates obtained from urinary tract infections in Kerala, South India. The research aimed to understand the phylogenetic groups, antibiotic resistance patterns, presence of antibiotic resistance genes (ARGs), integrons, extraintestinal virulence genes, and the genetic diversity among 100 *E. coli* isolates. The isolates exhibited varying degrees of resistance, with ampicillin resistance being the most prevalent, followed by resistance to cefoxitin, cefpodoxime, nalidixic acid, trimethoprim, and cotrimoxazole. Remarkably, 96% of the isolates demonstrated multidrug resistance (MDR), and 86% harbored ARGs, while 32% carried integron 1 (*int1*). The majority of the isolates (79%) were classified as extraintestinal pathogenic *E. coli* (ExPEC), with 86% of these ExPEC strains containing ARGs. Additionally, one isolate exhibited extensive drug resistance (XDR). The study highlighted a significant correlation between the presence of virulence genes and antibiotic resistance. The observed high genetic diversity among ARG-harboring *E. coli* isolates emphasizes the intricate dynamics of antibiotic resistance in urinary tract infections caused by *E. coli*, underscoring the importance of understanding this association for effective treatment strategies (Sukumaran & Hatha 2021).

Conclusions

The escalating prevalence of quinolone resistance among urinary pathogens presents a formidable challenge in the management of urinary tract infections (UTIs). Our review underscores the urgent need for systematic strategies to address this growing threat to public health. The studies highlighted in this review reveal trends in antibiotic resistance, where the emergence of quinolone-resistant genes poses significant clinical and public health concerns. The detection of plasmid-mediated quinolone resistance (*PMQR*) genes, such as *qnrB* and *aac(6)-Ib-cr*, in UTI-causing pathogens, underscores the genetic complexity underlying antibiotic resistance. Urgent interventions are needed to fight the spread of resistant pathogens and preserve the efficacy of existing antibiotics. Collaborative efforts among healthcare providers, policymakers, researchers, and the community are essential to address this global public health challenge effectively. By implementing evidence-based interventions and fostering antimicrobial practices, we can strive towards preserving the effectiveness of antibiotics and ensuring optimal outcomes for patients with UTIs.

Abbreviations

UTIs: Urinary tract infections

PMQR: Plasmid-mediated quinolone resistance

ESBL: Extended-spectrum beta-lactamase

CREc: Carbapenem-resistant *Escherichia coli*

MDR: Multidrug resistance

XDR: Extensively drug-resistant

PDR: Pan-drug-resistant

WGS: Whole-genome sequencing

ARGs: Antibiotic resistance genes

ExPEC: Extraintestinal pathogenic *E. coli*

REFERENCE

1. Mancuso, G., Midiri, A., Gerace, E., Marra, M., Zummo, S., & Biondo, C. (2023). Urinary Tract Infections: The Current Scenario and Future Prospects. *Pathogens*, 12(4), 623.
2. Tan, C. W., & Chlebicki, M. P. (2016). Urinary tract infections in adults. *Singapore medical journal*, 57(9), 485.

Comment [AS6]: Suggestion of additional references:

Mansoor, S., Ullah, I., & Khan, N. (2024). Diversity and antibiotic resistance profile analysis of uropathogenic bacteria in human and canines. *Cellular, Molecular and Biomedical Reports*, 4(2), 65-73.
<https://doi.org/10.55705/cmbr.2023.411879.1169>

Srivastava, N., Verma, S., Singh, M., & Kumar, A. (2024). A short-term study on statistical numeration of multidrug resistant *Escherichia coli* isolates among the patients with urinary tract infection. <https://doi.org/10.56042/ijeb.v6i2i02.2549>

3. WHO, (2023) Target Product Profiles for Oral Therapy of Urinary Tract Infections. Available online: <https://www.who.int/publications/i/item/9789240003873> (28 July 2023).
4. Choe, H. S., Lee, S. J., Cho, Y. H., Çek, M., Tandoğdu, Z., Wagenlehner, F., ... & Mohammed, R. A. (2018). Aspects of urinary tract infections and antimicrobial resistance in hospitalized urology patients in Asia: 10-Year results of the Global Prevalence Study of Infections in Urology (GPIU). *Journal of Infection and Chemotherapy*, 24(4), 278-283.
5. Alasmary, M. Y. (2021). Antimicrobial Resistance Patterns and ESBL of Uropathogens Isolated from Adult Females in Najran Region of Saudi Arabia. *Clinics and Practice*, 11(3), 650-658.
6. PH, S., Attavar, P. C., TR, R., Kotian, M. S., & NS, D. (2024). Emergence of High-Level Antibiotic Resistance in *Klebsiella pneumoniae*: A Narrative Review. *South Asian Journal of Research in Microbiology*, 18(2), 1-8.
7. Zhou, Y., Zhou, Z., Zheng, L., Gong, Z., Li, Y., Jin, Y., ... & Chi, M. (2023). Urinary Tract Infections Caused by Uropathogenic *Escherichia coli*: Mechanisms of Infection and Treatment Options. *International Journal of Molecular Sciences*, 24(13), 10537.
8. Millanao, A. R., Mora, A. Y., Villagra, N. A., Bucarey, S. A., & Hidalgo, A. A. (2021). Biological effects of quinolones: A family of broad-spectrum antimicrobial agents. *Molecules*, 26(23), 7153.
9. Guan, X., Xue, X., Liu, Y., Wang, J., Wang, Y., Wang, J., ... & Pan, L. (2013). Plasmid-mediated quinolone resistance—current knowledge and future perspectives. *Journal of International Medical Research*, 41(1), 20-30.
10. Hooper, D. C., & Jacoby, G. A. (2015). Mechanisms of drug resistance: quinolone resistance. *Annals of the New York academy of sciences*, 1354(1), 12-31.
11. Strahilevitz, J., Jacoby, G. A., Hooper, D. C., & Robicsek, A. (2009). Plasmid-mediated quinolone resistance: a multifaceted threat. *Clinical microbiology reviews*, 22(4), 664-689.

12. Soucy, J. P. R. (2018). Antimicrobial resistance in urinary *Escherichia coli* in Quebec, Canada. McGill University (Canada).
13. TR, R., Attavar, P. C., Hydrose, S. P., Kotian, M. S., & NS, D. (2024). Antibiofilm and Antibacterial Properties of Herbal Extracts as Alternatives to Current Treatment Approaches: A Narrative Review. *Asian Journal of Medicine and Health*, 22(2), 12-22.
14. Laxminarayan, R., & Chaudhury, R. R. (2016). Antibiotic Resistance in India: Drivers and Opportunities for Action. *PLoS medicine*, 13(3), e1001974. <https://doi.org/10.1371/journal.pmed.1001974>
15. Sharma, A., Thakur, N., Thakur, A., Chauhan, A., & Babrah, H. (2023). The Challenge of Antimicrobial Resistance in the Indian Healthcare System. *Cureus*, 15(7), e42231. <https://doi.org/10.7759/cureus.42231>
16. Hilt, E. E., Parnell, L. K., Wang, D., Stapleton, A. E., & Lukacz, E. S. (2023). Microbial Threshold Guidelines for UTI Diagnosis: A Scoping Systematic Review. *Pathology and Laboratory Medicine International*, 43-63. <https://doi.org/10.2147/PLMI.S409488>
17. Cardiliya, A. P., Chandrasekar, M. J. N., & Nanjan, M. J. (2023). Incidence of biofilms among the multidrug resistant *E. coli*, isolated from urinary tract infections in the Nilgiris district, South India. *Brazilian Journal of Microbiology*, 1-10. 1007/s42770-023-01028-9
18. Gupta, S., Tripathi, P., & Shrivastava, S. (2021). Molecular studies and characterization of genes associated in drug resistance of uro-pathogens in central India. *Gene Reports*, 24, 101230. <https://doi.org/10.1016/j.genrep.2021.101230>
19. Chandran, S. P., Sarkar, S., Diwan, V., Pathak, A., Shah, H., Tamhankar, A. J., ... & Stålsby-Lundborg, C. (2017). Detection of virulence genes in ESBL producing, quinolone resistant commensal *Escherichia coli* from rural Indian children. *The Journal of Infection in Developing Countries*, 11(05), 387-392. <https://jidc.org/index.php/journal/issue/view/136>
20. Boueroy, P., Chopjitt, P., Hatrongjit, R., Morita, M., Sugawara, Y., Akeda, Y., Iida, T., Hamada, S., & Kerdsin, A. (2023). Fluoroquinolone resistance determinants in carbapenem-resistant *Escherichia coli* isolated from urine clinical samples in Thailand. *PeerJ*, 11, e16401. <https://doi.org/10.7717/peerj.16401>

21. Azargun, R., Sadeghi, M. R., Soroush Barhaghi, M. H., Samadi Kafil, H., Yeganeh, F., Ahangar Oskouee, M., & Ghotaslou, R. (2018). The prevalence of plasmid-mediated quinolone resistance and ESBL-production in Enterobacteriaceae isolated from urinary tract infections. *Infection and drug resistance*, 11, 1007–1014. <https://doi.org/10.2147/IDR.S160720>
22. FarajzadehSheikh, A., Veisi, H., Shahin, M., Getso, M., & Farahani, A. (2019). Frequency of quinolone resistance genes among extended-spectrum β -lactamase (ESBL)-producing *Escherichia coli* strains isolated from urinary tract infections. *Tropical medicine and health*, 47, 1-7. <https://doi.org/10.1186/s41182-019-0147-8>
23. Alhazmi, A. H., Alameer, K. M., Abuageelah, B. M., Alharbi, R. H., Mobarki, M., Musawi, S., ... & Dhayhi, N. (2023). Epidemiology and Antimicrobial Resistance Patterns of Urinary Tract Infections: A Cross-Sectional Study from Southwestern Saudi Arabia. *Medicina*, 59(8), 1411.
24. SUKUMARAN, D., & Hatha, M. (2021). Molecular Epidemiology of Multidrug-resistant *Escherichia coli* from Urinary Tract Infections. *Journal of Microbiology and Infectious Diseases*, 11(02), 66-73.