

Original Research Article

Principal component analysis of Yield and its attributing traits in Recombinant Inbred Lines of Rice under submerged condition(*Oryza sativa* L.)

Abstract

Evaluation of Principle Component Analysis (PCA) of recombinant inbred lines (RILs) was done at phenotypic level under submerged conditions to reduce a large series of data into smaller number of components by looking for groups that have very strong inter-correlation in a set of variables and each component explained *per cent* variation to the total variability. The RIL population was derived from an inter-specific cross between BPT5204 and HPR14 parents. A study was conducted using 1256 Recombinant Inbred Lines submerged condition in the two seasons at College of Agriculture V.C. Farm, Mandya with nine agro-morphological traits and a principle component analysis was carried out. Out of nine principle components, four exhibited Eigenvalue more than one governing 77.74% variance and 69.86% variance in the *summer* and *kharif* seasons respectively. The highest positive Eigenvalue was observed for total number of tillers, productive tillers, non-productive tillers and fallowed by single plant yield in PC1 in the *summer* and *kharif* season respectively. The highest positive Eigenvalue was observed for five panicle weight, single panicle length, single plant yield and plant height in PC2 of *summer* and *kharif* season respectively. Indicating their pronounced effect on the overall variation in the Recombinant Inbred Lines of Rice.

Keywords: Rice, RILs, PCA, Eigenvalues, Yield.

Comment [mm1]: On abstract: please add introduction, prolems, aims, methods, results, and novelty

Comment [mm2]: Alphabetic

Introduction:

Rice, *Oryza sativa* L., is the world's most important staple crop, feeding more than half of the world's population. It is a self-pollinated, short-day, C3 plant that belongs to the

Poaceae family and originated in South East Asia (Meghana et al., 2015). It is a healthy cereal crop that provides 20 % of calories and 15 % of protein and is consumed by half of the world's population. Rice is also the most important and dependable food crop in India, feeding more than two-thirds of the population. In India, rice is grown in an area of 43 million ha with a production of 118.43 million tons and average productivity of 2.75 t ha⁻¹ (Anon, 2020). The Kharif rice production in India during 2021–22 is estimated to be 107.04 million tonnes (Mt) (Anonymous, 2021). Rice yield increase of 1.0–1.2% per year beyond 2020 will be required to feed the world's still-growing population while keeping prices low (Anonymous, 2013). Plant breeders often assess numerous traits, some of which may lack sufficient discriminatory power for germplasm evaluation, characterization, and management (Maji and Shaibu, 2012). In such cases, Principal Component Analysis (PCA) emerges as a valuable tool. PCA is a multivariate statistical technique employed to unveil patterns within datasets and streamline them by reducing redundancy. It aims to simplify and assess the interrelationships among a multitude of variables into a smaller set of components without compromising the essential information within the original dataset. According to Clifford and Stephenson (1975), the initial three principal components typically capture the most crucial variation patterns. With the above arguments an experiment was conducted to evaluate the Principal Components Analysis of the 9 traits in Recombinant Inbred Lines population was derived from an inter-specific cross between BPT5204 and HPR14 parents.

Material and methods:

Plant materials

Parents with diverse genetic background *viz.*, BPT-5204 (good grain qualities and high yield) and HPR14 (high protein content) (Hittalmani, 1990) were crossed to develop 1255 segregating lines and selections were carried out based on phenotypic parameters in RILs of rice.

Experimental site and layout

The RIL population along with parents was planted in an augmented design at college of agriculture V C Farm, Mandya, during summer 2020 and observations were recorded on individual lines and used for statistical analysis. Twenty one days seedlings were transplanted with 20cm x 10cm spacing and minimum of 15 plants maintained in each line. Recommended cultural practices for rice cultivation were carried out to ensure uniform crop stand as per the package of practices. Phenotypic data was collected for a total of 1255 RILs related to yield and yield contributing traits like days to 50% flowering (DFF) was recorded as the number of days from germination to 50% of plants with initiation of flowering based

Comment [mm3]: Please add 1-2 paragraph and explain problems and novelty this research

on visual observation of the each RIL, plant height (PH) was recorded at maturity stage (cm) from base of the plant at soil surface to the panicle tip of main tiller and averaged over 5 plants, number of tillers per plant (TN) was measured by counting total number of tillers per plant at harvesting stage and averaged over five plants, panicle length (PL) (cm) measured from the panicle neck to the tip (excluding awn) at reproductive stage, panicle weight (PW) was calculated by total weight of 5 panicles and averaged, thousand grain weight (TW) (grams) measured by weighing 1000 filled grains from each plant (Mohanty *et al.*, 2018).

Result and Discussion:

Principal component analysis (PCA) is a statistical procedure that used to identify the traits which contribute maximum to the most observed variation within a large group of genotypes and employed to reduce the complexity of the data while minimizing the variation within the data and increasing interpretability (Upadhyay *et al.*, 2022). It is efficient to finding the components useful to group the genotypes that maximize the variation and it is also helpful in reducing the number of variables with a maximum variance but does not estimate the specific effect. The PCA was performed separately for each season using data of all nine agro-morphological traits evaluated on 1256 recombinant inbred lines of rice genotypes. The resulted PCA from nine independent principal components (PCs) explained cumulative variance during summer and Kharif season during 2020.

The importance of a character towards the PCs could be seen from the corresponding eigenvalues. The results of PCA revealed that, out of the 9 PCs, first four PCs with eigenvalue of greater than 1.00 accounted for 77.74 per cent of total variability for summer season (Table 1). The PC1 accounted for 29.19 per cent of the total variation in the population during summer and the results also showed that total number of tillers (0.584), productive tillers (0.528), non-productive tillers (0.447), single plant yield (0.344) contributed positively and significantly to PC1 followed by days to fifty per cent flowering, five panicle weight and single panicle length. The PC2 accounted for 21.23 per cent of the total variation in the population during summer and the results also showed that five panicle weight (0.591), single panicle length (0.457), single plant yield (0.386), test weight (0.364) contributed positively and significantly to PC2 followed by plant height and days to fifty per cent flowering (Table2).

PC	Eigenvalue	Per centage of Variance	Cumulative per centage of variance
1	2.62	29.19	29.19

Comment [mm4]: On discussion: Please add discussion for all parameters. Add references, explain the results from the others study, then compares with your study. Then placed discusion before conclusion

2	1.91	21.23	50.43
3	1.48	16.54	66.97
4	1.06	10.77	77.74
5	0.75	8.38	86.13
6	0.52	5.79	91.92
7	0.38	4.24	96.17
8	0.32	3.55	99.73
9	0.02	0.26	100

Table 1: Estimate of eigenvalues and per centage of variation contributed by each principal component (PC) to the single plant yield and yield related components under submerged conditions during summer season.

	PC 1	PC 2	PC 3	PC 4	PC 5	PC 6	PC 7	PC 8	PC 9
DF	0.135	0.129	-0.704	-0.123	0.015	-0.139	0.537	0.382	-0.004
PH	-0.004	0.282	0.66	-0.124	-0.173	0.104	0.557	0.345	0.013
TT	0.584	-0.147	0.112	0.092	0.147	-0.052	0.091	-0.082	-0.76
NPT	0.447	-0.175	-0.025	-0.343	0.229	0.692	-0.104	0.137	0.301
PT	0.528	-0.112	0.138	0.276	0.084	-0.432	0.193	-0.229	0.575
TW	-0.132	0.364	-0.013	0.607	0.634	0.264	0.076	0.06	0.007
5 P W	0.143	0.591	-0.152	-0.112	-0.261	0.244	0.144	-0.668	-0.018
1 P L	0.083	0.457	0.102	-0.532	0.468	-0.416	-0.302	0.088	-0.001
S P Y	0.344	0.386	-0.053	0.321	-0.447	0.009	-0.477	0.446	0.028

Table 2: Contribution of each trait to the principle component during summer season 2020

DDF- Days to 50 % flowering, **PH**- Plant height (cm), **TT**- Total number of tillers, **NPT**- Number of non-productive tillers, **PT**-Number of productive tillers, **TW**- Test weight (gm), **5PW**- 5 Panicle weight (gm), **1PL**- One panicle length, **SPY**- Single plant yield.

The first four PCs with eigenvalue of greater than 1.00 accounted for 69.86 per cent of total variability for kharif season (Table 3). The PC1 accounted for 25.67 per cent of the total variation in the population during kharif and the results also showed that total number of tillers (0.642), productive tillers (0.598), non-productive tillers (0.378), single plant yield (0.201) contributed positively and significantly to PC1 followed by days to fifty per cent flowering, five panicle weight. The PC2 accounted for 20.00 per cent of the total variation in the population during kharif and the results also showed that single panicle length (0.512), plant height (0.496), five panicle weight (0.476), single plant yield (0.410) contributed positively and significantly to PC2 followed by days to fifty per cent flowering and test weight (Table 4). Recorded results are in the line of Sahu *et al.* (2016), Manohara *et al.* (2020), Christina *et al.* (2021).

Comment [mm5]: On discussion: Please add discussion for all parameters. Add references, explain the results from the others study, then compares with your study. Then placed discusion before conclusion

PC	Eigenvalue	Per centage of variance	Cumulative per centage of variance
1	2.31	25.67	25.67
2	1.80	20.00	45.68
3	1.12	12.46	58.14
4	1.05	11.72	69.86
5	0.83	9.24	79.10
6	0.71	7.90	87.01
7	0.64	7.17	94.18
8	0.52	5.78	99.97

9	0	0.02	100
---	---	------	-----

Table 3: Estimate of eigenvalues and per centage of variation contributed by each principal component (PC) to the single plant yield and yield related components under submerged conditions during Kharif season

	PC 1	PC 2	PC 3	PC 4	PC 5	PC 6	PC 7	PC 8	PC 9
DF	0.194	0.272	-0.542	-0.145	0.556	0.024	0.513	0.009	-0.001
PH	-0.043	0.496	0.456	-0.216	-0.085	-0.061	0.257	0.649	0.001
TT	0.642	-0.077	0.087	-0.031	-0.026	-0.162	-0.084	0.047	-0.733
NPT	0.378	-0.106	0.412	0.035	0.379	0.686	-0.083	-0.026	0.232
PT	0.598	-0.05	-0.05	-0.044	-0.164	-0.437	-0.067	0.062	0.64
TW	-0.028	0.026	0.131	0.894	0.292	-0.249	0.08	0.167	-0.003
5 P W	0.065	0.479	-0.415	0.126	-0.013	0.215	-0.682	0.26	0

1 PL	-0.047	0.512	0.356	-0.127	0.305	-0.319	-0.218	-0.592	-0.002
SPY	0.201	0.41	-0.044	0.311	-0.578	0.314	0.365	-0.356	0.001

Table 4: **Contribution of each trait to the principle component during kharif season 2020**

DDF- Days to 50 % flowering, **PH**- Plant height (cm), **TT**- Total number of tillers, **NPT**- Number of non-productive tillers, **PT**-Number of productive tillers, **TW**- Test weight (gm), **5PW**- 5 Panicle weight (gm), **1PL**- One panicle length, **SPY**- Single plant yield.

Conclusion:

PCA can be used to extract all of the relevant components and emphasize their contributions to overall variability, making it an excellent tool for speeding up the breeding process. Out of nine principle components, four exhibited Eigenvalue more than one governing 77.74% variance and 69.86% variance in the *summer* and *kharif* seasons respectively. The highest positive Eigenvalue was observed for total number of tillers, productive tillers, non-productive tillers and followed by single plant yield in PC1 in the *summer* and *kharif* season respectively. The highest positive Eigenvalue was observed for five panicle weight, single panicle length, single plant yield and plant height in PC2 of *summer* and *kharif* season respectively. Indicating their pronounced effect on the overall variation in the Recombinant Inbred Lines of Rice.

Reference:

Anonymous (2013). GRiSP (Global Rice Science Partnership 2013). Rice almanac, 4th edition. International Rice Research Institute, Los Baños, Philippines, p.283.

Anonymous (2021). Second Advanced Estimates of Production of Major crops for 2021-22 Press Information Bureau, Delhi.

ANONYMOUS,2020,Adikailuvarigeadunikabesayapaddathigalupackageofpracticesforfieldvr ops. Univ. Agril.Sci., Bangalore.

CHRISTINA,M.,JONES,M.R.,VERSINI,A.,MÉZINO,M.,LEMEZO,L.,AUZOUX,S.,SOULI

Comment [mm6]: Add references using limited
5 years ago

E,J.C.,POSER,C.ANDGÉRARDEAUX,E.,2021,Impactofclimate variability and extreme rainfall events on sugarcane yield gap in a tropical Island. *FieldCropsResearch*, **274**:108-112.

Clifford, H. T. and Stephenson, W. (1975). An Introduction to Numerical Classification. Academic Press, London. p. 229.

HITTALMANI, S.V., 1990,Inheritanceof quantitative traitsand exploitationof male sterilityin rice(*Oryzasativa*L.)breeding.*M.ScThesis*, Univ.Agril.Sci.,Bangalore.

Maji, A. T. and Shaibu, A. A. (2012). Application of principal component analysis for rice germplasm characterization and evaluation. *Journal of Plant Breeding and Crop Science*, 4(6): 87–93.

MANOHARA, K. K., MAHAJAN, G. R. AND SAHOO, R. N., 2020. Spectroscopy based novel spectral indices, PCA-and PLSR-coupled machine learning models for salinity stress phenotyping of rice.

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, **229**:117-121.

MEGHANA,H.S.,SHAILAJA,H.,GANDHI,R.V.ANDMEERA,N.,2015,Phenotypic screening for salt tolerance at germination and seedling stage and SSR marker validation in rice(*Oryza sativa*L.).*Mysore J. Agric. Sci.*, **49**(4): 686-692.

MOHANTY,A.,MARNDI,B.C.,SHARMA,S.ANDDAS,A.,2011,Biochemical characterization of two high protein rice cultivars from Assam rice collections.*Oryza*, **48**(2):171-174.

UPADHYAY, S., RATHI, S., CHOUDHARY, M., SNEHI, S., SINGH, V., SINGH, P.K. AND SINGH, R.K., 2022 Principal Component analysis of Yield and its attributing Traits in advanced Inbred Lines of Rice under Sodicity condition (*Oryza sativa* L.).