

**An Evaluation of Hepato-Protective Activity of Ethanolic ~~extract~~  
Extract of *Solanum nigrum* on CCL<sub>4</sub> Induced Hepatic Injured Rat**

**Abstract:**

Using herbs and herbal therapies to promote health and avoid, alleviate, or cure illness is known as herbal medicine, which is both an art form and a profession. Herbal remedies are another name for herbal medicine in certain communities. Researchers looked at the lipid profiles of *Solanum nigrum* extract by using rats in their studies. In the case of SGPT, group 6 showed statistically significant outcomes ( $p < 0.05$ ), whereas in the case of SGOT, both groups 5 and 6 showed statistically significant outcomes ( $p < 0.05$ ). In relation to the SGPT and the SGOT, this is being said. Both group 5 and group 6 showed statistically significant amounts of creatinine and urea ( $p < 0.05$ ) during the renal function test. Group 6 with a dose of 1200 mg/kg showed

statistically significant results for high-density lipoprotein (HDL) and low-density lipoprotein (LDL) ( $p < 0.05$ ). Furthermore, at dosages of 600 mg/kg and 1200 mg/kg, respectively, the triglyceride level in groups 5 and 6 was statistically significant ( $p < 0.05$ ). There were no results that were deemed statistically significant for total cholesterol.

**Keywords:** Herbal medicine, *Solanum nigrum*, HDL, LDL, Phytochemicals.

## Introduction

The majority of a person's bodily processes are controlled by the liver, the biggest glandular organ. Numerous times during the day, the whole blood supply of an individual goes through the liver. The liver is very critical for human metabolism [1]. Hepatotoxicity, the most frequent form of liver illness, is a leading cause of death and impairment in both animals and humans. Several medications can cause it [2]. Addiction to alcohol or drugs, exposure to harmful chemicals, infections with viruses or parasites, and high levels of reactive oxygen species (ROS) are among the most damaging factors that may affect liver cells ( $\text{OH}$ ,  $\text{H}_2\text{O}_2$ ,  $\text{O}_2$ ) [2]. The ability to neutralize free radicals is why ascorbic acid and the tripeptide L-glutathione (L-cysteine,

Formatted: Font: (Default) Times New Roman, 12 pt, Subscript

Formatted: Font: (Default) Times New Roman, 12 pt, Subscript

Formatted: Font: (Default) Times New Roman, 12 pt, Subscript

glycine, and L-glutamate) are taken orally as dietary supplements. Because of their anti-inflammatory, antibacterial, and immune-enhancing qualities, they are highly prized by many.

**The third.** On the flip side, they might trigger an allergic response, such as dermatitis, or gastrointestinal issues, such as gas, diarrhea, indigestion, or even difficulties breathing due to airway narrowing. There are 1.5 billion individuals worldwide who suffer from chronic liver disease (CLD). Among Americans aged 45–64 years, the prevalence of CLD has risen by 31% in recent years. [4]. Some chemical components derived from medicinal plants may have therapeutic uses, say specialists in the subject. It follows that researchers are always on the lookout for new herbal cures and other plant-based therapies to treat a wide range of illnesses. [5]. While phytotherapy is based on scientific study, herbalism is more concerned with the practical applications of medicinal plants. Plants have played a significant role in human medicine for thousands of years due to the wide diversity of chemicals they contain, many of which have medicinal characteristics [6]. The vast variety of chemical components found in medicinal plants allows them to exert a broad spectrum of pharmacological and therapeutic effects. Tanning agents, glycosides, alkaloids, saponins, polysaccharides, essential oils, terpenoids, resins, and plant lipids are all examples of such components [7-9]. Modifying plants genetically allows for the precise regulation of chemical concentrations, allowing for the desired therapeutic effect. Reverse genetics has many potential applications, one of which is to enhance the production of secondary metabolites like alkaloids [10].

Black nightshade, deadly nightshade, or *Solanum nigrum* (SN) belonging to the family Solanaceae, is a widespread weed plant that grows wild in open fields throughout Africa, the Americas, and even the driest parts of India and other parts of the world [11]. It is also known as EnabEldib in Egypt. A wide range of conditions have has been traditionally treated with *S. nigrum* in folk medicine, including: pain, fever, inflammation, cough, asthma, wound, ulcer, leprosy, skin diseases, hemorrhoids, dropsy, liver disorders, antioxidant, hepatoprotective agents, neuroprotective, cytoprotective, antimicrobial, antinociceptive, and antipyretic effects [12–19]. Some people think *S. nigrum* is a great treatment for liver problems [13]. Solasodine, tomatidenol, tigogenin, solamargine, and solasonine are the primary components derived from the aerial sections of *S. nigrum*. Other components include lutein, lycopene, crytoxanthin, vitamin C, glucose, fructose, and tomatidenol. Polysaccharides, glycoproteins, and glycoalkaloids make up the bulk of the active ingredients. Polyphenolic compounds like gallic

**Comment [MOU1]:** Incomplete sentence

**Formatted:** Font: (Default) Times New Roman, 12 pt, Highlight

**Comment [MOU2]:** After reference [2], you have put number [4]. Number [3] is missing

**Formatted:** Font: (Default) Times New Roman, 12 pt, Italic

acid, catechin, protocatechuic acid, caffeic acid, epicatechin, rutin, and naringenin are also present and are thought to be responsible for a variety of therapeutic effects [20].

We are now investigating the potential hepatoprotective benefits of *Solanum nigrum*.

## Materials and methods

### Plant Collection and Extract Preparation

*Solanum nigrum* were collected from local market of Dhaka. The material was authenticated by the National Herbarium Bangladesh. *Solanum-S. nigrum* was 1<sup>st</sup> cleaned properly and air-dried. Next, it was severely crushed. The powder was soaked in 90% ethanol and kept for 15 days. By this time the solution was vigorously shaken occasionally. Then, the extract was filtered and collected. The extracted material was dried in a rotary evaporator at a low temperature and pressure. Finally, the crude residue was subjected to the required pharmacological testing.

### Drugs and Chemicals

Carbon tetrachloride (CCl<sub>4</sub>), a well-known hepatotoxicity causing chemical, was purchased from the Sigma Aldrich. The API of typical anti-oxidant medication silymarin was received as a free gift from Incepta Pharmaceuticals Limited

### Experimental Animal Procurement, Nursing, and Grouping

A total of 100 male rats weighing between 120 and 150 grams were obtained from Jahangirnagar University in Savar, Dhaka. Each of them was housed in a climate-controlled environment (temperature 25±3°C, relative humidity 55±5%, and a 12-h light/dark cycle) at the University of Dhaka's Institute of Nutrition & Food Science (INFS). They were given a conventional food and were permitted to drink clean water. All of the animals were maintained in this habitat for at least one week prior to the research for adaption. All experimental methods followed the recommendations of the Institutional Animals Ethics Committee (IEAC).

### Animal Model Sample Size Detection

**Comment [MOU3]:** Correct as This research aims to evaluate the potential hepatoprotective benefits of *Solanum nigrum*

**Comment [MOU4]:** What is the plant part/parts that you used?

**Comment [MOU5]:** Mention the solvent you use e.g. tap water/distilled water etc

**Comment [MOU6]:** What did you use for crush? E.g. mortar and pestle/ mechanical grinder etc

**Comment [MOU7]:** What are the weight of the powder and volume of 95% ethanol? Clearly mention the ratio.

**Comment [MOU8]:** What did you use for filter? E.g. filter paper/ muslin cloth etc. If it is a filter paper give specificity of that? Mention whether it is gravitational filtration or suction filtration.

**Comment [MOU9]:** What did you collect? If you collect filtrate, specifically mention that "filtrate was collected".

**Comment [MOU10]:** Use term filtrate instead of extracted materials

**Comment [MOU11]:** Mention temperature and pressure

**Comment [MOU12]:** Give yield%

**Formatted:** Subscript

**Formatted:** Strikethrough

**Comment [MOU13]:** How did you calculate sample size? Give sample size calculation details with references

**Comment [MOU14]:** Why did you only used male rats? Why didn't you get female rats also?

**Comment [MOU15]:** Mention the species strain Are they Wistar rats or any other?

**Comment [MOU16]:** Is it an animal house at the University? If it is mention it

**Comment [MOU17]:** Specify this? Is it a standard diet?

**Comment [MOU18]:** Say clean water was provided *ad-libitum*

**Comment [MOU19]:** Use the term acclimatized

**Comment [MOU20]:** Give ERC number

There were 100 rats in all, and they were randomly divided into ten groups of ten. The rats were randomly allocated to one of the groups. We watched the rat closely each day to ensure better health status.

**Comment [MOU21]:** How did you determine the number of rats per group?

**Comment [MOU22]:** Use passive style. Also use the term observed instead of watched.

**Comment [MOU23]:** The sentences are not in formal style. Rewrite sentences

### Dose Selection and Route of Administration for Respective Study

Carbon tetrachloride ( $\text{CCl}_4$ ) is a common chemical agent used in laboratories to study a range of liver diseases, both acute and chronic. Trichloromethyl free radical ( $\text{CCl}_3$ ), a CYP2E1 isozyme-produced  $\text{CCl}_4$  metabolite, reacts with cellular lipids and proteins to form trichloromethylperoxy radical, which attacks lipids on the endoplasmic reticulum membrane faster than the trichloromethyl free radical, causing lipid peroxidation and lobular necrosis. A single oral treatment of  $\text{CCl}_4$  mixed with olive oil as a vehicle in a 1:1 ratio (3 mL/kg of rat body weight) produced hepatic damage in all animal groups except the usual control group. *Momordica charantia* extracts were administered to animals with hepatic injury as a post-treatment. The extract was administered orally in various doses.

**Formatted:** Highlight

**Formatted:** Subscript, Highlight

**Formatted:** Highlight

**Formatted:** Subscript, Highlight

**Formatted:** Highlight

**Formatted:** Subscript, Highlight

**Formatted:** Highlight

**Formatted:** Subscript, Highlight

**Formatted:** Highlight

**Comment [MOU24]:** Don't put this highlighted detail under the methodology. Include those details under the introduction or discussion.

**Formatted:** Highlight

**Formatted:** Subscript

**Comment [MOU25]:** How did you decide this dose?

**Comment [MOU26]:** Which control group? Negative control or positive control?

**Comment [MOU27]:** Why did you

**Formatted:** Highlight

**Comment [MOU28]:** At what time you gave *S. nigrum* extract? How did you select different doses? What is positive control group?

### Evaluation of Hepato-Protective Activity

For this experiment, 100 rats were randomly picked and equally divided into fourteen groups

**Table 1:** Application of treatment efficacy

Group Number	Group Specification	Treatment species	Dose treatment species (mg/kg)	Abbreviation of Groups
1	Negative	Physiological	10	N

	Control	saline	<del>ml</del> ml/kg	
2	CCl <sub>4</sub> Control	N/A	N/A	A
3	CCl <sub>4</sub> + S <sub>10</sub>	Silymarin	10	S <sub>10</sub>
4	CCl <sub>4</sub> + SN <sub>300</sub>	<i>Solanum nigrum</i>	200	SN <sub>300</sub>
5	CCl <sub>4</sub> + SN <sub>600</sub>	<i>Solanum nigrum</i>	400	SN <sub>600</sub>
6	CCl <sub>4</sub> +SN <sub>1200</sub>	<i>Solanum nigrum</i>	600	SN <sub>1200</sub>
7	S <sub>10</sub>	Silymarin	10	S <sub>10</sub>
8	SN <sub>300</sub>	<i>Solanum nigrum</i>	200	SN <sub>300</sub>
9	SN <sub>600</sub>	<i>Solanum nigrum</i>	400	SN <sub>600</sub>
10	SN <sub>1200</sub>	<i>Solanum nigrum</i>	600	SN <sub>1200</sub>

### Biological Sample Collection:

Blood was drawn from the animals via puncturing the heart and transferred to a centrifuge tube. The samples were centrifuged at 5,000 rpm for 5 minutes to create the supernatant fluid. Biochemical testing subsequently required the transfer of this fluid to an additional micro-centrifuge tube.

### Estimation of Biochemical Parameters

**Comment [MOU29]:** How did you euthanize the animal?  
What are the anesthetic reagents you used?

**Comment [MOU30]:** Use scientific terms (Cardiac punch)

**Comment [MOU31]:** What is the volume you collected?  
Give syringe and needle size

Lipid profile, kidney, and liver function tests were performed in Humaluzer 3000 using respective parameter Kit.

Comment [MOU32]: Give parameters specifically

### Statistical analysis:

All of our findings (raw data) in terms of numerical parameters were recorded and analyzed on a broadsheet using the MS Excel application. The gathered data were subjected to descriptive statistics, with the findings reported as mean SD. To evaluate statistical significance, we used the SPSS 16 software's "One-way ~~Anova~~ ANOVA test" to interpret inter-group heterogeneity in terms of several biological factors. The occurrences are considered statistically significant since the 'p' value was less than 0.05 ( $p < 0.05$ ).

### Results and discussion

The term "herbal medicine" refers to the practice of using medicinal plants for the ~~purpose of preventing and treating~~ to prevent and treat sickness. This practice encompasses a wide range of therapies, including those that are common in every culture as well as others that are standardized and tritated. In the course of this study, we evaluated the lipid profile of the herb *Solanum nigrum* in rat models to determine its effectiveness as an anti-diabetic agent.

Group 6 demonstrated statistically significant ( $p < 0.05$ ) outcomes in the case of SGPT, whereas ~~group Group 5~~ and ~~group Group 6~~ demonstrated statistically significant ( $p < 0.05$ ) results in the case of SGOT. This is with regard to both the SGPT and the SGOT. A total of two further studies came to the same conclusions [21, 22].

Comment [MOU33]: Expand the abbreviation

When conducting the kidney function test, it was observed that the levels of creatinine and urea were statistically significant ( $p < 0.05$ ) in both group 5 and group 6. Comparable findings were discovered in a number of other investigations [23, 24].

In the case of high-density lipoprotein (HDL) and low-density lipoprotein (LDL), the findings were statistically significant ( $p < 0.05$ ) in group 6 with a dosage of 1200 mg/kg. Additionally, the triglyceride level was statistically significant ( $p < 0.05$ ) in Groups 5 and 6 with doses of 600 mg/kg and 1200 mg/kg, respectively. In the case of total cholesterol, there were no findings that

were recognized as statistically significant. There were two other investigations that came to the same conclusions [25, 26].

**Table 2:** Different biochemical parameters of *Solanum nigrum*:

Groups	SGPT	SGOT	Creatinine	Urea	TC	HDL	LDL	TG
C	35.39±0.48	46.24±1.28	0.5±0.04	32.47±3.19	96.82±3.28	68.57±3.52	34.28±0.94	50.71±2.62
CCl <sub>4</sub>	99.37±6.82	106.43±4.89	2.83±0.08	98.59±0.32	174.28±5.73	42.58±4.48	88.29±5.61	103.57±8.91
CCl <sub>4</sub> + S <sub>10</sub>	62.45±4.61	57.28±5.38	1.46±0.08	56.27±5.55	141.34±6.42	59.74±4.67	68.22±6.53	74.81±4.53
CCl <sub>4</sub> +SN <sub>300</sub>	99.21±4.15	102.03±5.16	2.50±0.09	95.93±4.21	172.86±3.21	44.23±4.89	86.24±5.15	100.42±7.95
CCl <sub>4</sub> +SN <sub>600</sub>	97.36±5.57	92.10±4.53*	2.09±0.08*	89.41±3.39*	170.53±4.59	44.82±2.30	85.43±3.37	97.63±6.29*
CCl <sub>4</sub> +SN <sub>1200</sub>	94.68±4.02*	86.77±3.27*	1.63±0.06*	84.28±4.29*	169.93±5.21	46.59±3.93*	82.01±5.04*	92.52±6.82*
S <sub>10</sub>	37.10±1.39	48.32±2.36	0.6±0.05	33.16±1.93	92.32±2.10	67.57±2.91	36.82±1.06	52.59±3.63
SN <sub>300</sub>	35.60±2.08	45.57±3.13	0.8±0.02	35.21±2.06	96.53±3.16	64.83±3.56	34.53±1.81	50.78±2.32
SN <sub>600</sub>	37.37±1.28	47.48±3.36	0.5±0.08	30.28±1.50	95.90±2.03	69.20±4.01	35.62±2.80	54.62±3.18
SN <sub>1200</sub>	34.36±3.38	42.15±0.29	0.7±0.03	32.38±3.17	93.26±3.53	65.23±3.01	34.94±2.42	48.10±2.51

**Note:** The results were expressed in Mean±SEM (standard mean error) \*p< 0.05, \*\*p< 0.01, and \*\*\*p< 0.001 were considered as statistically significant. The statistical analysis followed by one-way analysis of variance (Dunnett's test) compared to the control.

## Conclusion

Within the scope of this investigation, the hepatoprotective properties of the [95%](#) ethanolic extract of *Solanum nigrum* were investigated. The results of this study support the idea that an ethanol extract from the plant *Solanum nigrum* might help protect against high cholesterol, liver damage, and kidney problems. As a result, more research is necessary in order to determine the active components in the entire extract that have the ability to reduce hyperlipidemia and diabetes. When it has been determined which chemicals are active, it is possible to conduct a comprehensive exploration.

#### References:

1. Bhawna S, Kumar SU. Hepatoprotective activity of some indigenous plants. *Int J Pharm Tech Res.* 2009 Oct;4:1330-4.
2. Nadeem M.P.C, Dandiya P.C, K.V., Pasha M., Imran D., Balani K, Vohora S.B., Hepatoprotective activity of *Solanum nigrum* fruits. *Fitoterapia.*, 1997, 68, 245:51
3. FM SS, Juliana AB, Bornila M, Puja B, Nur-Neasha D, Rafat T. An Assessment of Hepato-Protective Activity of *Psidium guajava* Fruit Extract against Hepatic Injured Rodent Model. *Asian Journal of Medical Principles and Clinical Practice.* 2023 Oct 7;6(2):240-5.
4. Islam M, Rupak AH, Nasrin N, Chowdhury MM, Sen P, Foysal AU, Uddin MJ, Ferdous J, Tahsin MR, Aktar F, Kabir S. An Evaluation of Potential Hepato-Protective Properties of *HylocereusUndatus* Fruit in Experimental Rat Model. *Biomedical Journal of Scientific & Technical Research.* 2022;43(2):34405-16.
5. Baroi JA, Hossian MR, Chowdhury MM, Dolon NN, Maliha F, Rupak MA, Lima NN, Ullah MR, Tahsin R. An assessment of anti-hyperlipidemic potentialities of ethanolic extract of *hemidesmus indicus* in high fat induced rat model. *Asian Journal of Food Research and Nutrition.* 2023 Jul 9;2(4):323-30.
6. Mim IJ, Peya FY, Chowdhury MM, Khan TR, Mandal SK, Maliha F, Alam M, Rahman T, Tashin R. An evaluation of anti-diabetic activity of ethanolic extract of *asparagus racemosus* in alloxan induced rat model. *International Journal of Advances in Nephrology Research.* 2023 Aug 2;6(1):60-8.

7. Lima NN, Dolon NN, Maliha F, Ullah MR, Humayra F, Chowdhury MM, Rupak MA, Baroi JA, Shohan FS, Tashin R. An Evaluation of Analgesic and Anti-Inflammatory Activity of *Ficus racemosa* in Rat Model.
8. Yang L, Stöckigt J. Trends for diverse production strategies of plant medicinal alkaloids. *Natural product reports* 2010;27(10):1469-1479.
9. Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *Journal of Pharmacognosy and Phytochemistry*. 2013;1(6):168-182. Rupak MA, Chowdhury MM, Shurov
10. Rupak MA, Chowdhury MM, Shurovi FS, Ferdous J, Tahsin MR, Sarif S, Hasan MM, Chowdhury JA, Kabir S, Chowdhury AA, Aktar F. An Evaluation of Analgesic and Anti-Inflammatory Activity of Ethanolic Extract of *Cynodon Dactylon* on Stressed Rodent Model. *Biomedical Journal of Scientific & Technical Research*. 2022;42(3):33550-7
11. Mbagwa FN, Nwachukwu CU, Okore OO (2007) Root anatomical studies on *Solanum macrocarpum* and *Solanum nigrum* (Solanaceae). *J Am Sci* 3(3)
12. Arulmozhi V, Krishnaveni M, Karthiashwaran K, Dhamodharan G, Mirunalini S (2010) Antioxidant and antihyperlipidemic effect of *Solanum nigrum* fruit extract on the experimental model against chronic ethanol toxicity. *Pharmacogn Mag* 6(21):41-50
13. Abdou ZA, Attia MH, Raafat MA (2007) Protective effect of citric acid and thiol compounds against cadmium and lead toxicity in experimental animals. *J Biol Chem Environ Sci* 2(2):481-497
14. Tarique AQ, Sayeed MZ, Moyad S, Ghassan S, Nahid B (2008) Effect of *Solanum nigrum* on immobilization stress induced antioxidant defence changes in rat. *Res J Biol Sci* 3(12):1426-1429
15. Elhag RAM, El-Badwi SMA, Bokhiet AO, Galall M (2011) Hepatoprotective activity of *Solanum nigrum* extracts on chemically-induced liver damage in rat. *J Vet Med Anim Health* 3(4):45-50

16. Perez RM, Perez JA, Garcia LMD, Sosa HM (1998) Neuropharmacological activity of *Solanum nigrum* fruit. *J Ethnopharmacol* 62(1):43–48
17. Prashaut KV, Shasidhara S, Kumar MM, Sridhar A (2001) Cytoprotective role of *S. nigrum* against gentamicin-induced kidney cell damage. *Fitoterapia* 72:481–486
18. Jainu M, Shayamala CS (2004) Antioxidant effect of methanolic extract of *Solanum nigrum* barriers on aspirin induced gastric mucosal injury. *Indian J ClinBiochem* 19(1):57–61
19. Rani P, Khullar N (2004) Antimicrobial evaluation of some medicinal plants for their anti-enteric potential multidrug resistant. *Phytother Res* 18:670–673
20. Sildar M, Dutta U (2008) Traditional phytotherapy among the Nath people of Assam. *EthnoMed* 2:39–45
21. Elhag RA, El Badwi SM, Bakhiet AO, Galal M. Hepatoprotective activity of *Solanum nigrum* extracts on chemically induced liver damage in rats. *Journal of Veterinary Medicine and Animal Health*. 2011 Aug;3(4):45-50.
22. Abdel-Rahim EA, Abdel-Mobdy YE, Ali RF, Mahmoud HA. Hepatoprotective effects of *Solanum nigrum* Linn fruits against cadmium chloride toxicity in albino rats. *Biological trace element research*. 2014 Sep;160:400-8.
23. Sankaran M. Protective effect of *Solanum nigrum* fruit extract on the functional status of liver and kidney against ethanol induced toxicity. *Journal of Biochemical Technology*. 2012 Feb 20;3(4):339-43.
24. Azarkish F, Hashemi K, Talebi A, Kamalinejad M, Soltani N, Pouladian N. Effect of the administration of *Solanum nigrum* fruit on prevention of diabetic nephropathy in streptozotocin-induced diabetic rats. *Pharmacognosy research*. 2017 Oct;9(4):325.
25. Chang JJ, Chung DJ, Lee YJ, Wen BH, Jao HY, Wang CJ. *Solanum nigrum* polyphenol extracts inhibit hepatic inflammation, oxidative stress, and lipogenesis in high-fat-diet-treated mice. *Journal of agricultural and food chemistry*. 2017 Oct 25;65(42):9255-65.

26. ÖNER AC, Fatmagül YU, FETHULLAH MN. Antioxidant and Antihyperlipidemic Effect of Solanum Nigrum Extract in Experimental Diabetes Model. Van Veterinary Journal. 2023 Sep 1;34(3):184-8.

UNDER PEER REVIEW