

On some recurrence relations connected with generalized Fermat numbers and some properties of divisibility for these numbers

Abstract

As a result of nice properties of Fermat numbers and their interesting applications, these numbers have recently seen a variety of developments and extensions. Within this framework, this paper contributes. The purpose of this paper is to obtain some recurrence relations connected with generalized Fermat numbers $\mathcal{F}_n = a^{2^n} + 1$ for $a, n \in \mathbb{Z}$ and $n \geq 0$ and as a result of these recurrent relations, to get some properties of divisibility for generalized Fermat numbers.

Keywords: Fermat number; recurrence relation; divisibility of integers

2010 Mathematics Subject Classification: 11B83, 11B37, 13A05.

1 Introduction

For $n \in \mathbb{Z}$ and $n \geq 0$, a Fermat number F_n is a positive integer of the form

$$F_n = 2^{2^n} + 1.$$

The first Fermat numbers are 3, 5, 17, 257, 65537, 4294967297, ... (sequence A000215 in the OEIS), see (12).

Since 1607–1665 when Fermat numbers were studied by the XVII-th century French lawyer and mathematician Pierre de Fermat, many people have been working about and around it, in many different directions and with a lot of applications in number theory, see (2), (7), (8), (9), (11). The study of the various properties of the integers is often based on their arithmetic properties such as periodicity, congruences, and divisibility, all of which can be found in (3), (4) and (5).

Many mathematicians studied different aspects of the Fermat numbers during the XIX-th and XX-th centuries.

The Fermat number F_n for $n = 0, 1, 2, \dots$ must always give a prime according to Fermat's conjecture in 1650. As a matter of fact, the first five Fermat numbers are prime numbers: $F_0 = 3, F_1 = 5, F_2 = 17, F_3 = 257, F_4 = 65537$. However, in 1732, L. Euler discovered that F_5 is a composite number:

$$F_5 = 2^{2^5} + 1 = 4294967297 = 641 \cdot 6700417,$$

where $641, 6700417 \in P$.

The set of Fermat factors starts with the numbers 3, 5, 17, 257, 641, 65537, 114689, 274177, 319489, 974849, ... (sequence A023394 in OEIS; see also sequence A050922 in OEIS), see (12).

As of 2020, 354 prime factors of Fermat numbers are known, and it is known that 310 Fermat numbers are composite. New Fermat factors are discovered each year. Subsequently, digital computers have facilitated the discovery of more factors. The distributed computing project Fermat Search is searching for new factors of Fermat numbers (6).

Calvo (1) reported some properties mainly related to the presence of common prime factors of both standard Fermat numbers, $F_n = 2^{2^n} + 1$, and generalized Fermat numbers, $\mathcal{F}_n = a^{2^n} + 1$.

Riesel (10) studied on a method to find certain common factors of numbers of the form $A_n = a^{2^n} + 1$.

Witno (13) investigated the behaviour of the numbers $E_n = (3^{2^n} + 1) / 2$ and presented an initial primality test for E_n using the knowledge that E_{n-1} is prime.

In this paper, we aim to obtain some recurrent relations connected with the generalized Fermat numbers of the form $\mathcal{F}_n = a^{2^n} + 1$ for $a, n \in \mathbb{Z}$ and $n \geq 0$, and then to investigate some properties of divisibility of the generalized Fermat numbers.

2 Recurrent relations for the generalized Fermat numbers

In this section, we now give some new recurrent relations connected with the generalized Fermat numbers of the form $\mathcal{F}_n = a^{2^n} + 1$ for $a, n \in \mathbb{Z}$ and $n \geq 0$.

In the following proposition, we obtain the generalized Fermat number with index n , using the generalized Fermat numbers with the previous index $n - 1$.

Proposition 2.1. For $n \geq 1$, we have

$$\mathcal{F}_n = (\mathcal{F}_{n-1} - 1)^2 + 1.$$

Proof. For $n \geq 1$, we have

$$\begin{aligned} \mathcal{F}_n - 1 &= a^{2^n} \\ &= a^{2^{n-1} \cdot 2} \\ &= \left(a^{2^{n-1}}\right)^2 \\ &= (\mathcal{F}_{n-1} - 1)^2. \end{aligned}$$

Thus, the proof is completed. □

In the following propositions, we obtain the generalized Fermat number with index n , using the generalized Fermat numbers with all previous indexes $0, 1, 2, \dots, n - 1$.

Proposition 2.2. For $n \geq 1$, we have

$$\mathcal{F}_n = (a - 1) \mathcal{F}_0 \mathcal{F}_1 \dots \mathcal{F}_{n-1} + 2. \tag{2.1}$$

Proof. For $n \geq 1$, we have

$$\begin{aligned}
 \mathcal{F}_n - 2 &= a^{2^n} - 1 \\
 &= a^{2^{n-1} \cdot 2} - 1 \\
 &= \left(a^{2^{n-1}}\right)^2 - 1 \\
 &= \left(a^{2^{n-1}} + 1\right) \left(a^{2^{n-1}} - 1\right) \\
 &= \left(a^{2^{n-1}} + 1\right) \left(a^{2^{n-2}} + 1\right) \left(a^{2^{n-2}} - 1\right) \\
 &= \dots \\
 &= \left(a^{2^{n-1}} + 1\right) \left(a^{2^{n-2}} + 1\right) \dots \left(a^{2^1} + 1\right) \left(a^{2^0} + 1\right) \left(a^{2^0} - 1\right) \\
 &= \mathcal{F}_{n-1} \mathcal{F}_{n-2} \dots \mathcal{F}_0 (a - 1).
 \end{aligned}$$

Thus, the proof is completed. □

Proposition 2.3. For $n \geq 2$, we have

$$\mathcal{F}_n = \mathcal{F}_{n-1} + a^{2^{n-1}} (a - 1) \mathcal{F}_{n-2} \mathcal{F}_{n-3} \dots \mathcal{F}_0.$$

Proof. For $n \geq 2$, we have

$$\begin{aligned}
 \mathcal{F}_n - \mathcal{F}_{n-1} &= a^{2^n} - a^{2^{n-1}} \\
 &= a^{2^{n-1} \cdot 2} - a^{2^{n-1}} \\
 &= a^{2^{n-1}} a^{2^{n-1}} - a^{2^{n-1}} \\
 &= a^{2^{n-1}} \left(a^{2^{n-1}} - 1\right) \\
 &= a^{2^{n-1}} \left(a^{2^{n-2} \cdot 2} - 1\right) \\
 &= a^{2^{n-1}} \left(\left(a^{2^{n-2}}\right)^2 - 1\right) \\
 &= a^{2^{n-1}} \left(a^{2^{n-2}} + 1\right) \left(a^{2^{n-2}} - 1\right) \\
 &= \dots \\
 &= a^{2^{n-1}} \left(a^{2^{n-2}} + 1\right) \left(a^{2^{n-3}} + 1\right) \dots \left(a^{2^1} + 1\right) \left(a^{2^0} + 1\right) \left(a^{2^0} - 1\right) \\
 &= a^{2^{n-1}} \mathcal{F}_{n-2} \mathcal{F}_{n-3} \dots \mathcal{F}_0 (a - 1).
 \end{aligned}$$

Thus, the proof is completed. □

Moreover, the generalized Fermat numbers satisfy the following recurrence relation.

Proposition 2.4. For $n \geq 2$, we have

$$F_n = F_{n-1}^2 - 2(F_{n-2} - 1)^2.$$

Proof. For $n \geq 2$, we have

$$\begin{aligned}
 \mathcal{F}_{n-1}^2 - 2(\mathcal{F}_{n-2} - 1)^2 &= \left(a^{2^{n-1}} + 1\right)^2 - 2 \cdot \left(a^{2^{n-2}}\right)^2 \\
 &= \left(a^{2^{n-1}}\right)^2 + 1 + 2 \cdot a^{2^{n-1}} - 2 \cdot a^{2 \cdot 2^{n-2}} \\
 &= a^{2^n} + 1 \\
 &= \mathcal{F}_n.
 \end{aligned}$$

Thus, the proof is completed. \square

3 Divisibility properties of the generalized Fermat numbers

Using the results reached before, we obtain some divisibility properties of the generalized Fermat numbers \mathcal{F}_n in this section.

Theorem 3.1. *For $m \neq n$, we have $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 1$ or $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 2$.*

Proof. In order to prove $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 1$ or $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 2$, let assume that $\gcd(\mathcal{F}_m, \mathcal{F}_n) = d$. In this case, it holds that $d \mid \mathcal{F}_n$ and $d \mid \mathcal{F}_m$. Without loss of generality, let $n > m$. Since $d \mid \mathcal{F}_n$, from (2.1) we get

$$d \mid (a - 1) \mathcal{F}_0 \mathcal{F}_1 \dots \mathcal{F}_m \dots \mathcal{F}_{n-1} + 2. \tag{3.1}$$

Also, as $d \mid \mathcal{F}_m$, then

$$d \mid (a - 1) \mathcal{F}_0 \mathcal{F}_1 \dots \mathcal{F}_m \dots \mathcal{F}_{n-1}. \tag{3.2}$$

Using relations in (3.1) and (3.2), we get $d \mid 2$. Thus, it is proven that $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 1$ or $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 2$. \square

Corollary 3.2. *Let a in the n th generalized Fermat number $\mathcal{F}_n = a^{2^n} + 1$ be an even integer. Then, the numbers \mathcal{F}_m and \mathcal{F}_n have no common prime divisors, i.e., for $m \neq n$, we have $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 1$.*

Proof. In order to prove this formula, let assume that $\gcd(\mathcal{F}_m, \mathcal{F}_n) = d, d > 1$. Since every integer $n \geq 2$ has a prime factor, there exists a prime number p , such that $p \mid d$. In this case, it holds that $p \mid \mathcal{F}_n$, and $p \mid \mathcal{F}_m$. Therefore, we have that $p \mid 2$, i.e., $p = 2$. Since a is an even integer, any Fermat number is an odd integer. Thus, it is not divisible by 2. It gives a contradiction. Consequently, we obtain $\gcd(\mathcal{F}_m, \mathcal{F}_n) = 1$ for $n \neq m$. \square

This fact allow as to give one more proof of infiniteness of the set of prime numbers.

Theorem 3.3. *There are infinity many prime numbers.*

Proof. Let a in the n th generalized Fermat number $\mathcal{F}_n = a^{2^n} + 1$ be an even positive integer. If we consider the set $\{\mathcal{F}_0, \mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_n, \dots\}$ of all the generalized Fermat numbers, then obviously we obtain that this set is infinite. If we denote by p_n the smallest prime divisor of each \mathcal{F}_n , then from Corollary 3.2 we get $p_n \neq p_m$ for $n \neq m$. Therefore, all elements of the set $P = \{p_0, p_1, p_2, \dots, p_n, \dots\}$ are different odd prime numbers and the set P is an infinite subset of the set of primes. Thus, we obtain that the set of all primes is infinite. \square

References

- [1] Calvo, I. J. (2000). A note on factors of generalized Fermat numbers. *Applied Mathematics Letters*, 13(6), 1-5.
- [2] Carmichael, R. D. (1919). Fermat Numbers $F_n = 2^{2^n} + 1$. *The American Mathematical Monthly*, 26(4), 137-146.
- [3] Conway, J. H., Guy, R. (1998). *The book of numbers*. Springer Science Business Media.
- [4] Davenport, H. (1999). *The higher arithmetic: An introduction to the theory of numbers*. Cambridge University Press.
- [5] Deza, E. (2021). *Mersenne numbers and Fermat numbers (Vol. 1)*. World Scientific.
- [6] Keller, W. (2020). Prime Factors of Fermat Numbers, <http://www.prothsearch.com/fermat.html>, 29.12.2020.
- [7] Koshy, T. (2002). *Elementary number theory with applications*. Academic press.
- [8] Luca, F. (2001). Fermat numbers in the Pascal triangle. *Divulgaciones Matemáticas*, 9(2), 189-194.
- [9] Maddux, F. C. (2005). *Fermat numbers: historical view with applications related to fermat primes*. San Jose State University.
- [10] Riesel, H. (1969). Common prime factors of the numbers $A_n = a^{2^{t_n}} + 1$, *BIT* 9, 264-269.
- [11] Robinson, R. M. (1954). Mersenne and Fermat numbers. *Proceedings of the American Mathematical Society*, 5(5), 842-846.
- [12] Sloane N.J.A. at all On-line Encyclopedia of Integer Sequences, <http://oeis.org/>29.12.2020.
- [13] Witno, A. (2010). On Generalized fermat numbers $3^{2^n} + 1$. *Applied Math. Information sciences*, 4(3) (2010), 307-313.