

Original Research Article

Mineral and labile organic nitrogen fractions in soil profile and their response to FYM and inorganic fertilization in different growth stages of rice crop

ABSTRACT

Nitrogen is the key element among the major nutrients in crop production. The mineralizable soil organic nitrogen is the main contributor of soil N supply. A better understanding of soil organic nitrogen dynamics in agro-ecosystems is needed to improve N management. The present investigation was conducted in Regional Agricultural Research Station (RARS), Assam Agricultural University, Titabar, Jorhat during 2017-18. In this study, the impact of FYM and inorganic fertilization on soil mineral nitrogen (NO_3^- -N and NH_4^+ -N) dynamics and labile organic nitrogen fractions viz. microbial biomass N (MBN), particulate organic N (PON) and water-extractable organic N (WEON) at three growth stages of rice viz. active tillering, flowering and physiological maturity stages and nitrogen stock were assessed. Six treatments viz. control (no fertilization), 100% NPK, 100% NPK + FYM 5 t ha^{-1} , 50% NPK, 50% NPK + 50% N through FYM and FYM 10 t ha^{-1} were tested in randomized block design with four replications. The results showed that NO_3^- -N and NH_4^+ -N were found to be significantly higher in 100% NPK + FYM 5 t ha^{-1} at the three growth stages of rice. The labile organic nitrogen fractions were significantly higher in FYM 10 t ha^{-1} . All these variables were decreased with increase in crop growth stages. Integrated use of inorganic fertilizer and FYM recorded the highest nitrogen stock. Stepwise regression analysis indicated that NH_4^+ -N was the main contributor to nitrogen stocks at the three growth stages of rice of the studied soils.

Keywords: Mineral nitrogen; Labile organic nitrogen fractions; N stock; rice growth stages

1. INTRODUCTION

Of all the essential nutrients, nitrogen (N) has the most pronounced effect on plant growth and development. But most of the agricultural soils are deficient in this nutrient (Ayeni 2011). Since a bulk of total soil N is present in organic form, the quantity of inorganic N is usually small and easily depleted and hence cannot support plant growth for a long time. Soil organic nitrogen (SON) is the main contributor of soil N supply (Haynes 2005, Sharifi *et al.* 2008). Short-term changes in total SON are difficult to detect, therefore it is partitioned into various pools of differing turnover rates (Chen *et al.* 2018). There is a lack of knowledge concerning the involvement of the labile organic N pool in N cycle and its significance to soil N supply. The individual fractions of labile organic N viz., microbial biomass N (MBN), particulate organic N (PON) and water-extractable organic N (WEON) have not been studied in different agricultural systems, across soils and climatic conditions. Hence, a better understanding of their interrelationships, underlying mechanisms and N transformation

through different fractions is needed. The study on various N fractions at crop growth stages has received less attention and little relevant information in paddy soil. Keeping this in view, the research was carried out with the objective to evaluate the impact of FYM and inorganic fertilizer on mineral N dynamics, labile organic N fractions and N stock during three growth stages of rice.

2. MATERIALS AND METHODS

The investigation was conducted at Regional Agricultural Research Station (RARS), Assam Agricultural University, Titabor, Jorhat in 2017-18 during *sali* season in an *Inceptisol*. The climatic condition of the region is characterized by a subtropical environment with hot humid summer and relatively dry and cool winter with an annual mean temperature of 24.5°C and a mean annual precipitation of 1,146 mm. The soil is sandy clay loam texture with organic carbon 1.1%, pH 5.4, CEC 12.5 cmol (p+) kg⁻¹, available N 495.0 kg ha⁻¹, P 22.2 kg ha⁻¹ and K 112.0 kg ha⁻¹ at initiation of the experiment. The rice variety Gitesh was used in the present investigation which is suitable for staggered planting in *sali* season of Assam with aged seedlings. Six treatments viz., control (no fertilization), 100% NPK, 100% NPK + FYM 5 t ha⁻¹, 50% NPK, 50% NPK + 50% N through FYM and FYM 10 t ha⁻¹ were tested in Randomized Block Design with four replications. Inorganic NPK fertilizers were applied at the rate of 40 kg ha⁻¹ N, 20 kg ha⁻¹ P and 20 kg ha⁻¹ K as urea, single super phosphate and muriate of potash, respectively. All fertilizers were applied as basal except urea, in which two thirds of urea was applied as basal fertilization and one third as top dressing. The available N, P and K in FYM was 0.45, 0.24 and 0.52%, respectively. Composite soil samples were collected from surface (0-20 cm) in three growth stages of rice viz. active tillering, flowering and physiological maturity stage. For determination of N stock, soil samples were collected after harvest of rice. The composite soil sample was divided into two parts. The first part of soil samples was processed (<0.25 mm sieve) for determination of mineral N (NO₃⁻-N and NH₄⁺-N), PON and WEON and the other part was stored in refrigerator for MBN analysis. NO₃⁻-N and NH₄⁺-N was estimated by Colorimetric methods outlined by Baruah and Barthakur (1997). The MBN was estimated by using chloroform fumigation extraction according to the method presented by Vance *et al.* (1987). The MBN was calculated as follows (Brookes *et al.* 1985):

$$\text{Microbial biomass N (mg kg}^{-1}\text{)} = \text{EN/kEN}$$

Where, EN = [(extractable N from fumigated soil) – (extractable N from non-fumigated soil)]; kEN = 0.54, represents the efficiency of extraction of MBN.

The PON was determined by the procedure outlined by Cambardella and Elliot (1992). The WEON was extracted using the method presented by Haney *et al.* (2012). Soil N stock was calculated after harvest of rice according to the method described by Ellert and Bettany (1995).

N stock (Mg ha⁻¹) = pb (g cm⁻³) x total N (kg ha⁻¹) x soil depth (cm), Where, pb = bulk density

All the observations were statistically analyzed by using the statistical methods described by Panse and Sukhatme (1989).

3. RESULTS AND DISCUSSION

3.1 Mineral N dynamics at different growth stages of rice under FYM and inorganic fertilization

Application of 100% NPK+FYM 5tha⁻¹ recorded significantly the highest NO₃⁻-N and NH₄⁺-N at the three growth stages of rice (Table 1). The increase in NO₃⁻-N and NH₄⁺-N in integrated application of FYM 5 tha⁻¹ with 100% NPK might be attributed to the addition of higher amount of root biomass which enhanced the microbial activity and nitrification process (Reddy *et al.* 2003, Bharti 2013, Rout *et al.* 2017).

The mineral N fractions of the soils decreased with development of the rice growth stages (Table 1). The decrease in NO₃⁻-N from active tillering to physiological maturity stage might be due to continuous N uptake by rice plant throughout the growing period (Meng *et al.* 2014, Manivannan and Sriramachandrasekharan 2016). The decrease in NH₄⁺-N with development of the growth stages might be because of transformation of a large proportion of basal fertilizers into NH₄⁺-N, but its uptake by rice plant was lower at the active tillering stage leading to higher NH₄⁺-N content in the soils (Zhang *et al.* 2009). The decrease in NH₄⁺-N with time might also be attributed to preferential uptake of NH₄⁺-N by rice crop.

Table 1. Mineral N dynamics at different growth stages of rice

Treatments	NO ₃ ⁻ -N (mg kg ⁻¹)			NH ₄ ⁺ -N (mg kg ⁻¹)		
	Active tillering	Flowering	Physiological maturity	Active tillering	Flowering	Physiological maturity
Control (no fertilization)	6.11 ^f	5.26 ^f	4.37 ^f	12.96 ^f	11.44 ^f	10.24 ^e
100% NPK	9.85 ^d	8.48 ^d	7.18 ^d	16.83 ^d	16.03 ^d	15.26 ^c
100% NPK+ FYM 5 t ha ⁻¹	15.26 ^a	13.47 ^a	12.28 ^a	22.13 ^a	21.23 ^a	20.17 ^a
50% NPK	8.16 ^e	6.20 ^e	6.08 ^e	14.35 ^e	13.47 ^e	12.11 ^d
50% NPK+ 50% (FYM) N	10.85 ^c	9.11 ^c	8.14 ^c	17.23 ^c	16.73 ^c	15.43 ^c
FYM 10 t ha ⁻¹	13.81 ^b	11.02 ^b	10.87 ^b	20.09 ^b	19.15 ^b	18.08 ^b
SEm±	0.14	0.15	0.15	0.21	0.19	0.23
CD (P= 0.05)	0.39	0.43	0.39	0.61	0.52	0.64

Means in a same column followed by different letter (s) are significantly different at P≤0.05

3.2 Labile soil organic N fractions at different growth stages of rice under FYM and inorganic fertilization

The fractions of labile soil organic N viz., MBN, PON and WEON were found to be the highest under FYM 10 t ha⁻¹ at the three growth stages of rice (Table 2). The organic manure FYM might have created a suitable condition for microbial growth as it is a good substratum for microbial activity. Many studies had reported that organic manures increased MBN of croplands and then increased MBN contributed to plant growth by enhancing the nutrient source and sink function (Singh *et al.* 2007, Cerny *et al.* 2008). The significant positive effect of FYM 10 t ha⁻¹ in PON than other treatments might be due to increase of the root and microbial biomass which are the main sources of PON (Manna *et al.* 2007, Naresh *et al.* 2017). FYM might also contribute to the PON fraction in soil because of significant quantities of N in FYM were retained in soil particulate fractions (Whalen *et al.* 2003). The results of present investigation revealed that use of FYM 10 t ha⁻¹ recorded the highest WEON at the three growth stages of rice. FYM could be used by soil microorganisms as a source of energy and be mineralized to inorganic N. Application of FYM could enhance soil microbial activity and organic matter decomposition and this might be attributed to higher WEON in the soils (Chantigny 2003). The lower WEON in the inorganic fertilization might be due to the utilization of WEON by the microbial biomass at higher rates (Whalen *et al.* 1999).

The MBN, PON and WEON were decreased with development of the rice growth stages (Table 2). The higher MBN at active tillering stage could be attributed to greater microbial activity caused by greater root exudation (Aulakh *et al.* 2001) and optimum moisture availability at this early stage of the crop (Islam and Borthakur 2016). This gradually declined when the crop was attained physiological maturity stage. Tamilselvi *et al.* (2015) observed higher MBN during vegetative stage of maize that was declining till the crop harvest. PON is an important source for mineralizable N. With the development of crop growth stages, organic matter of the soils was decreasing and therefore, PON might trend to decrease towards the later growth stages of rice. The higher WEON at active tillering stage might be attributed to the fact that, this fraction of labile organic N could be utilized by soil microorganisms at very early stages of crop growth and immediately converted to inorganic N. As a result, mineral N was higher at the early growth stage of rice. With time, rice plant prefers to uptake mineral N and thereby, WEON decreased in the later crop growth stages.

Table 2. Labile soil organic N fractions at different growth stages of rice

Treatments	MBN (mg kg ⁻¹)			PON (g kg ⁻¹)			WEON (mg kg ⁻¹)		
	Active tillering	Flowering	Physiological maturity	Active tillering	Flowering	Physiological maturity	Active tillering	Flowering	Physiological maturity
Control(no fertilization)	17.64 ^f	16.97 ^f	15.64 ^f	0.23 ^f	0.19 ^f	0.17 ^c	6.65 ^f	5.99 ^f	4.90 ^e
100% NPK	22.95 ^d	21.55 ^d	20.97 ^d	0.67 ^d	0.59 ^d	0.53 ^c	10.74 ^d	9.95 ^d	8.97 ^c
100% NPK+ FYM 5 t ha ⁻¹	26.30 ^b	25.91 ^b	24.56 ^b	0.97 ^b	0.92 ^b	0.90 ^a	11.95 ^b	11.46 ^b	10.58 ^a
50% NPK	20.16 ^e	19.76 ^e	18.94 ^e	0.27 ^e	0.24 ^e	0.23 ^d	7.59 ^e	6.70 ^e	5.88 ^d
50% NPK+ 50% (FYM) N	24.36 ^c	23.39 ^c	22.94 ^c	0.73 ^c	0.68 ^c	0.55 ^b	11.18 ^c	10.80 ^c	9.85 ^b
FYM 10 t ha ⁻¹	28.87 ^a	28.13 ^a	27.67 ^a	0.99 ^a	0.94 ^a	0.90 ^a	12.50 ^a	11.94 ^a	10.90 ^a
SEm±	0.23	0.17	0.18	0.01	0.01	0.01	0.10	0.13	0.13
CD (P= 0.05)	0.65	0.49	0.52	0.04	0.03	0.02	0.28	0.37	0.37
Means in a same column followed by different letter (s) are significantly different at P≤0.05									

3.3 Soil N stock

The highest N stock was observed 2.05 Mg ha⁻¹ in 100% NPK+ FYM 5 t ha⁻¹ which was significantly higher than the other fertilization treatments (Table 3). Gami *et al.* (2009) also reported significantly higher N stock in combined application of inorganic fertilizer and FYM than only inorganic NPK fertilized soils in Nepal.

Table 3 Soil N stock under FYM and inorganic fertilizer

Treatments	N stock (Mg ha ⁻¹)
Control (no fertilization)	1.22 ^f
100% NPK	1.66 ^c
100% NPK+ FYM 5 t ha ⁻¹	2.05 ^a
50% NPK	1.31 ^e
50% NPK+ 50% (FYM) N	1.47 ^d
FYM 10 t ha ⁻¹	1.81 ^b
SEm±	0.05
CD (P= 0.05)	0.14

Means in a same column followed by different letter (s) are significantly different at P≤0.05

3.4 Multiple regression analysis of different N fractions with N stock

The contributions of different fractions of N, alone or in combination, towards the variability of N stock were computed using stepwise multiple regression analysis (Table 4). At active tillering stage, NH₄⁺-N contributed 92.8% variation to N stock. All the fractions of N jointly contributed 94.6% variation to N stock. At flowering stage, NH₄⁺-N contributed to 92.2% variation to N stock. The R² increased from 0.922 to 0.940 when other N fractions were added. NH₄⁺-N contributed to 92.3% variation to N stock and R² was increased from 0.923 to 0.941 at physiological maturity stage. The other fractions of N contributed only 1.8% variation towards the N stock at this stage. Therefore, it is indicated that NH₄⁺-N was the main contributor to the N stock at the three growth stages of rice.

Table 4. Multiple regression analysis of different N fractions with N stock

Parameter	Regression Equation	R ²
Active tillering stage		
N stock	= 0.199 + 0.080* NH ₄ ⁺ -N	0.928

N stock	= 0.215 + 0.104* NH ₄ ⁺ -N - 0.042* WEON	0.937
N stock	= 0.277 + 0.096* NH ₄ ⁺ -N + 0.006* MBN -0.047 * WEON	0.942
N stock	= 0.343 + 0.033* NO ₃ ⁻ -N + 0.077* NH ₄ ⁺ -N + 0.010* MBN - 0.067* WEON	0.945
N stock	= 0.338 + 0.036* NO ₃ ⁻ -N + 0.075* NH ₄ ⁺ -N +0.011* MBN - 0.037* PON - 0.067* WEON	0.946

Flowering stage

N stock	= 0.317 + 0.077* NH ₄ ⁺ -N	0.922
N stock	= 0.345 + 0.103* NH ₄ ⁺ -N - 0.048* WEON	0.935
N stock	= 0.381 + 0.097* NH ₄ ⁺ -N + 0.004* MBN -0.049* WEON	0.937
N stock	= 0.398 + 0.014* NO ₃ ⁻ -N + 0.088* NH ₄ ⁺ -N + 0.007* MBN - 0.056* WEON	0.938
N stock	= 0.402 + 0.012* NO ₃ ⁻ -N + 0.089* NH ₄ ⁺ -N + 0.006* MBN + 0.0243* PON - 0.057* WEON	0.940

Physiological maturity stage

N stock	= 0.361 + 0.079* NH ₄ ⁺ -N	0.923
N stock	= 0.384 + 0.099* NH ₄ ⁺ -N - 0.038* WEON	0.931
N stock	= 0.419 + 0.092* NH ₄ ⁺ -N + 0.004* MBN -0.040* WEON	0.934
N stock	= 0.459 + 0.032* NO ₃ ⁻ -N + 0.078* NH ₄ ⁺ -N	0.938
N stock	= 0.444 + 0.034* NO ₃ ⁻ -N + 0.078* NH ₄ ⁺ -N + 0.012* MBN - 0.062* PON - 0.065* WEON	0.941

4. CONCLUSION

From the above said findings, it can be concluded that the study on mineral N dynamics, labile soil organic N fractions and N stock under intensive cropping system enables optimal N fertilization management practice and improving N supplying capacity of the soils. The present findings indicated that NH₄⁺-N has the major contribution towards N stocks of the studied soils.

REFERENCES

1. Aulakh M S, Wassmann R, Bueno C, Kreuzwieser J and Rennenberg H. 2001. Characterization of root exudates at different growth stages of ten rice (*Oryzasativa* L.) cultivars. *Plant Biology*. 2001; 3: 139-48.
2. Ayeni LS. Integrated plant nutrient management: A panacea for sustainable crop production in Nigeria. *International Journal of Soil Science*. 2011;6: 19-24.
3. Baruah TC, Barthakur HP. A Text Book of Soil Analysis. Vikas Publishing House Pvt. Ltd., New Delhi. 1997; 282.
4. Bharti B. Dynamics of organic and inorganic fractions of nitrogen in an acid alfisol after long-term addition of chemical fertilizers and amendments in maize-wheat cropping system. *International Journal of Agricultural Science and Research*. 2013; 3(3): 179-84.
5. Brookes PC, Landman A, Pruden G and Jenkinson DS. Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method for measuring microbial biomass nitrogen in soil. *Soil Biology Biochemistry*. 1985; 6: 837-42.
6. Cambardella CA, Elliott ET. Particulate soil organic matter changes across a grassland cultivation sequence. *Soil Science Society of America Journal*. 1992;56: 777-83.
7. Cerny J, Balik J, Kulhanek M and Nedved V. The changes in microbial biomass C and N in long-term field experiments. *Plant, Soil and Environment*. 2008; 54(5): 212-18.
8. Chantigny MH. Dissolved and water-extractable organic matter in soils: A review on the influence of land use and management practices. *Geodema*. 2003; 113: 357-80.
9. Chen Y, Xu X, Jiao X, Sui Y, Liu X, Zhang J, Zhou K and Zhang J. Responses of labile organic nitrogen fractions and enzyme activities in eroded mollisols after 8-year manure amendment. *Scientific Reports*. 2018; 8:1-9.
10. Ellert BH, Bettany JR. Calculation of organic matter and nutrients stored in soils under contrasting management regimes. *Canadian Journal of Soil Science*. 1995; 75: 529-38.
11. Gami SK, Lauren JG and Duxbury JM. Soil organic carbon and nitrogen stocks in Nepal long-term soil fertility experiments. *Soil Tillage Research*. 2009; 106: 95-03.
12. Haney RL, Franzluebbers AJ, Jin VL, Johnson MV, Haney EB, White MJ and Harmel RD. Soil Organic C:N vs. Water-Extractable Organic C:N. *Open Journal of Soil Science*. 2012; 2: 269-74.
13. Haynes RJ. Labile organic matter fractions as central components of the quality of agricultural soils: an overview. *Advances in Agronomy*. 2005; 85: 221-68.
14. Islam NF, Borthakur. Effect of different growth stages on rice crop on soil microbial and enzyme activities. *Tropical Plant Research*. 2016; 3(1): 40-47.
15. Manivannan R, Sriramachandrasekharan MV. Dynamics of inorganic fractions of nitrogen in an Ustifluvents soil on incorporation of organic manures and mineral nitrogen in rice. *International Journal of Development Research*. 2016; 6(9): 9333-38.

16. Manna MC, Swarup A, Wanjari RH and Ravankar HN. Long-term effects of NPK fertiliser and manure on soil fertility and a sorghum–wheat farming system. *Australian Journal of Experimental Agriculture*.2007; 47(6): 700-11.
17. Meng F, Olesen JE, Sun X and Wu W. Inorganic nitrogen leaching from organic and conventional rice production on a newly claimed calcicustoll in Central Asia. *PLoS One*.2014; 9(5): 1-10.
18. Naresh RK, Kumar A, Bhaskar S, Dhaliwal SS, Vivek, Kumar S, Kumar S and Gupta RK. Organic matter fractions and soil carbon sequestration after 15- years of integrated nutrient management and tillage systems in an annual double cropping system in northern India. *Journal of Pharmacognosy and Phytochemistry*.2017; 6(6): 670-83.
19. Panse VC, Sukhatme PV. *Statistical Methods for Agricultural Workers*, 4th edn, ICAR Publication, New Delhi, India. 1989; 34-35.
20. Reddy GB, Parawali G and Magaji SO. Influence of inorganic and organic nutrient sources on nitrogen under different depth. *Journal of American Society of Soil* .2003; 59(1): 544-56.
21. Rout PP, Chandrasekaran N, Rulmozhiselvan K and Padhan D. Effect of long-term fertilization on soil nitrogen dynamics and balance in an irrigated inceptisol under finger millet-hybrid maize cropping sequence. *Agriculture Update*.2017; 12: 1475-83.
22. Sharifi M, Zebarth BJ, Burton DL, Grant CA, Bittman S, Drury CF, McConkey BG and Ziadi N. Response of potentially mineralizable soil nitrogen and indices of nitrogen availability to tillage system. *Soil Science Society of America Journal*.2008; 72: 1124-31.
23. Singh S, Ghoshal N and Singh KP. Variation in soil microbial biomass and crop roots due to differing resource quality inputs in a tropical dryland agroecosystem. *Soil Biology and Biochemistry*.2007; 39: 76-86.
24. Tamilselvi SM, Chinnadurai C, Ilamurugu K, Arulmozhiselvan K and Balachandar D. Effect of long-term nutrient managements on biological and biochemical properties of semi-arid tropical Alfisol during maize crop development stages. *Ecological Indicators*.2015; 48: 76-87.
25. Vance ED, Brookes PC and Jenkinson DS. An extraction method for measuring soil microbial biomass C. *Soil Biology Biochemistry*.1987; 19: 703-07.
26. Whalen JK, Hu Q and Liu A. Manure applications improve aggregate stability in conventional and no-tillage systems. *Soil Science Society of America Journal*.2003; 67: 1842-47.
27. Whalen JK, Parmelee RW, McCartney DA and Vanarsdale JL. Movement of N from decomposing earthworm tissue to soil, microbial and plant N pools. *Soil Biology and Biochemistry*.1999; 31: 487-92.
28. Zhang J, Qin J, Yao W, Bi L, Lai T and Yu X. Effect of long-term application of manure and mineral fertilizers on nitrogen mineralization and microbial biomass in paddy soil during rice growth stages. *Plant, Soil and Environment*.2009; 55(3): 101-09.