

# EFFICACY OF AZOXYSTROBIN 7.5% + PROPICONAZOLE 12.5% SE AGAINST SHEATH BLIGHT IN RICE

## Abstract

Among the fungal diseases causing significant yield loss in rice, sheath blight caused by *Rhizoctonia solani* Kuhn is the most important ranked the second most important disease worldwide after blast and a serious threat in rice growing areas of the world and causes more economic yield losses. The field experiment was conducted with seven treatments and replicated three times in RBD design at ARS, Gangavati to know the effect of Azoxystrobin 7.5% + Propiconazole 12.5% SE against sheath blight in rice. The variety BPT-5204 was sown in plot size of 5 X 5 m<sup>2</sup> with all regular agronomic practices followed as per the standard package of practice of University of Agricultural Sciences, Raichur. It has been found that the fungicide Azoxystrobin 7.5% + Propiconazole 12.5% SE at different doses evaluated was effective in reducing the severity of rice sheath blight and thereby increased the rice grain yield. The treatment Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha recorded minimum sheath blight of 13.70 and 16.11 per cent PDI with yield of 64.60 q/ha and 63.20 q/ha compared to control treatment 48.70 and 57.96 per cent PDI with grain yield of 54.10 q/ha and 52.40 q/ha during the first and second season, respectively. In the current study, along with reducing sheath blight severity, fungicide treatments minimized grain yield losses in 2 years. Field trial conducted clearly indicated that Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 and 625 ml/ha dose can effectively control sheath blight of rice. Azoxystrobin 7.5% + Propiconazole 12.5% w/v SE @ 500 ml/ha was at par with higher dose and resulted better yield than other treatments.

**Keywords:** Azoxystrobin 7.5% + Propiconazole 12.5% SE, Fungicide, Rice, Sheath blight.

## **Introduction**

There are about 50 different biotic factors that can cause potential yield loss in rice including fungi, bacteria, viruses, nematodes and insects. Of the disease-causing organisms, fungal pathogens impose a greater challenge in sustaining rice production (Webster and Gunnell, 1992). Among the fungal diseases causing significant yield loss in rice, sheath blight is the most important ranked the second most important rice disease worldwide after blast (Pan *et al.*, 1999; Groth, 2005; Zhou and Jo, 2014). Sheath blight of rice caused by *Rhizoctonia solani* Kuhn is an important disease of rice and a serious threat in rice growing areas of the world and causes more economic yield losses (Ou, 1985; Savary *et al.*, 2000; Savary *et al.*, 2006), ranging from 20 to 50 per cent depending on the severity of infection

(Groth and Bond, 2007; Margani and Widadi, 2018) and also 5-10 per cent yield loss in subtropical low land paddy cultivars of Asia (Savary *et al.*, 2000). The yield losses ranging from 4 to 50 per cent have been reported depending on the crop stage at the time of infection, severity of the disease and environmental conditions (Singh *et al.*, 2004; Zheng *et al.*, 2013; Bhunkal *et al.*, 2015). The potential losses due to sheath blight disease are reported to be 50 to 54.3 per cent alone in India (Roy, 1993; Rajan, 1987) and this disease is particularly most prevalent in intensive rice cultivation system due to excess use of nitrogenous fertilizers.

In the absence of effective host plant resistance against sheath blight pathogen in rice, the management of sheath blight disease is mainly carried out through the use of chemicals (Naik *et al.*, 2017). Fungicide based management of sheath blight disease is successful at field level in majority of the cases (Kandhari *et al.*, 2003; Kandhari and Gupta, 2003; Groth and Bond 2006; Bhuvaneshwari and Raju, 2012; Kumar *et al.*, 2013). Foliar spray and seed treatment are the most popular method of fungicidal application against *R. solani*. Even though both systemic and non-systemic fungicides are used for chemical management, systemic fungicides offer better management of this disease (Naik *et al.*, 2017). Timely application of selective fungicides between panicle differentiation and heading stage offers effective protection against this disease. Periodical monitoring of the rice field and application of fungicides at the initial stages of infection especially at booting stage is recommended for managing sheath blight in susceptible varieties (Singh *et al.*, 2016; Uppala and Zhou, 2018). Most of the fungicides like benomyl, carbendazim, chloroneb, captafol, mancozeb, zineb, edifenphos, Iprobenphos, thiophanate, carboxin etc. have been found effective for the control of the disease under field conditions (Dash and Panda, 1984; Kannaiyan and Prasad, 1984; Singh and Sinha, 2004). Out of these benomyl, carbendazim, edifenphos and Iprobenphos were the most effective chemicals (Roy, 1993). For many years, strobilurin fungicides have been the backbone for management of rice sheath blight. It has been opined that *R. solani*, would not develop fungicide resistance or would be slow to develop resistance (Robinson, 2013). Presently, the Strobilurin group of systemic fungicides are the most preferred chemical group to manage sheath blight disease in rice (Yellareddygar *et al.*, 2014). Azoxystrobin from this group is very effective for not only controlling the disease but also found to enhance yield as well (Groth and Bond, 2007). The combinatory chemical formulation such as Azoxystrobin 18.2% + Difenoconazole 11.4% (Bhuvaneshwari and Raju, 2012; Kumar *et al.*, 2018), Trifloxystrobin 25% + Tebuconazole 50% (Shahid *et al.*, 2014; Rashid *et al.*, 2020), etc., are recommended to manage the disease. Recently, many combination fungicides such as kresoxim methyl 40% + hexaconazole 8%, azoxystrobin

18.2% + difenoconazole 11.4% SC, trifloxystrobin 25% + tebuconazole 50% 75 WG, have been shown to control the sheath blight disease under field condition (Kumar and Veerabhadraswamy, 2014; Bag, 2009) Continuous use of same group fungicides having same mode of action will lead to the development of resistant strain of same fungi and hence, it is necessary to search for a new molecule with different mode of action. Hence, the present study was under taken to know the field efficacy of a combination fungicide Azoxystrobin 7.5% + Propiconazole 12.5% SE against sheath blight in rice.

## **MATERIALS AND METHODS**

The field experiment was conducted with seven treatments and replicated three times in RBD design at ARS, Gangavati to know the effect of Azoxystrobin 7.5% + Propiconazole 12.5% SE against sheath blight in Rice. The variety BPT-5204 was sown in plot size of 5 X 5 m<sup>2</sup> with all regular agronomic practices followed as per the standard package of practice of University of Agricultural Sciences, Raichur. The evaluation of the fungicide was done along with standard checks and untreated control against the incidences of sheath blight disease in rice. Treatments details were as follows - T1 - Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 375 ml / ha; T2 - Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500ml / ha; T3 - Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml / ha; T4 - Azoxystrobin 23 % SC @ 500 ml / ha; T5 - Propiconazole 25% EC @ 500 ml / ha; T6 – Validamycin 3% L @ 2000 ml / ha; T7 - Untreated control.

The fungicides were applied as foliar spray treatment in the replicated plots just after the appearance of sheath blight disease in the main field at 45DAT. The plots were inspected regularly to see the disease development and further two more spray were applied at an interval of 15 days on 60 and 75 Days after Transplanting (DAT). Observations were recorded on disease severity in each treatment after three sprays as per the standard method The incidence of disease were recorded on leaves as Per cent Disease Index (PDI) on the basis of scoring of the diseases as per the disease rating scale of Standard Evaluation System (SES) for rice (IRRI, 2002). In the present study, observations for disease incidence were recorded from the randomly selected twenty clumps / hills per plot for recording the disease severity in each replicated plots of the treatments. The observations were recorded on intensity of diseases were observed in each replicated plot for each treatment on 10<sup>th</sup> day after each spray. After 10 days of last spray, the final scoring of the disease incidence was recorded. Further, the scored data were converted into Per cent Disease Index (PDI) of plants using formula given by Wheeler (1969)

$$\text{PDI} = \frac{\text{Number of leaves observed}}{\text{Maximum disease rating value}} \times \dots$$

In order to record the yield, crop was harvested plot-wise from the individual replicated plots and average paddy yield was recorded and converted into q/ha.

The original PDI values were suitably transformed into arcsine transformed values and subjected to statistical analysis for drawing conclusions

### **EXPERIMENTAL RESULTS AND DISCUSSION:**

It has been found that Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500- 625 ml/ha reduced the sheath blight infection more than rest of the treatments and improved the rice yield. The Azoxystrobin 7.5% + Propiconazole 12.5% SE was found to be effective in reducing the severity of the sheath blight disease. In the 1<sup>st</sup> season, the treatment Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha recorded least PDI of sheath blight disease incidence (13.70%) and was significantly superior over control treatment (48.70%). The same treatment was significantly on par with Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 ml/ha treatment with PDI of 14.07 per cent. All other treatments were inferior to these two treatments though significantly superior to the untreated check. Among the various treatments, Azoxystrobin 23% SC @ 500ml/ha was found least effective with higher PDI of sheath blight (26.11%) (Table-1). During the 2<sup>nd</sup> season, the Azoxystrobin 7.5% + Propiconazole 12.5% SE was found to be effective in reducing the severity of the sheath blight disease. The treatment Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha recorded least PDI of sheath blight disease incidence (16.11%) and was significantly superior over control treatment (57.96%). The same treatment was significantly on par with Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 ml/ha treatment with PDI of 16.48 per cent. All other treatments were inferior to these two treatments though significantly superior to the untreated check (57.96%). Among the various treatments, Azoxystrobin 23% SC @ 500ml/ha was found least effective with higher sheath blight (30.56%) (Table 2). These data were in accordance with Phelp and Soto (1993) and Jones *et al.* (1987). These finding are in full agreement with the previous reports of fungicide combinations such as azoxystrobin + difenconazole (Bhuvaneswari and Krishnam Raju, 2012), kresoxim methyl + hexaconazole, fluxapyroxad + epixiconazole (Kumar and Veerabhadraswamy, 2014), trifloxystrobin + tebuconazole (Bag and Saha, 2009; Bag, 2009; Visalakshmi *et al.*, 2016), flutolanil + azoxystrobin, thiophonate methyl + azoxystrobin, tebuconazole + azoxystrobin, propiconazole + azoxystrobin (Jin *et al.*, 2013) reported the better efficacy against sheath blight. Various reviews confirmed that strobilurin compounds

found to be effective in controlling many diseases like sheath blight (Bag 2009; Bag *et al.*, 2016).

### **Yield:**

Due to varying degree of reduction in sheath blight disease incidences the difference in the yield level between treated and untreated plots was very much significant. During the 1<sup>st</sup> season, the maximum grain yield of 64.60 q/ha was recorded with Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha followed by the same fungicide @ 500ml/ha which recorded 63.50 q/ha was statistically superior and comparatively effective than rest of the treatments. The treatments Propiconazole 25% EC, Validamycin 3% L, and Azoxystrobin 23% SC recorded the yield of 61.00, 60.70 and 59.40 q/ha, respectively. The lowest grain yield (54.10 q/ha) was recorded in untreated control (Table 1). During the 2<sup>nd</sup> season, the maximum grain yield of 63.20 q/ha was recorded with Azoxystrobin 7.5% + Propiconazole 12.5% w/v SE @ 625 ml/ha followed by the same fungicide @ 500ml/ha which recorded 62.20 q/ha was statistically superior and comparatively effective than rest of the treatments tested. The treatments Propiconazole 25% EC, Validamycin 3% L and Azoxystrobin 23% SC recorded the yield of 59.00, 58.70 and 57.60 q/ha, respectively. The lowest grain yield was recorded with 52.40 q/ha in untreated control (Table 2). In the current study, along with reducing sheath blight severity, fungicide treatments minimized grain yield losses in 2 years. The results are in agreement with previous studies that reported the efficacy of azoxystrobin (Groth, 2005), azoxystrobin and flutolanil (Groth and Bond, 2007) against sheath blight and in minimizing yield losses. Application of fungicides has been reported to enhance the crop yield due to reduction in disease load (Biswas and Bag, 2010; Bag, 2011). The present results are in conformity with those of previous reports (Bhuvaneshwari and Raju, 2012; Bag *et al.*, 2016) reported that fungicides application increases the yield of rice, mainly due to reduced disease severity of sheath blight.

### **CONCLUSION:**

The fungicide Azoxystrobin 7.5% + Propiconazole 12.5% SE at different doses were evaluated at Agriculture Research Station, Gangavati and was found effective in reducing the severity of rice sheath blight and thereby increased the rice grain yield. Field trial conducted clearly indicated that Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 and 625 ml/ha dose can effectively control sheath blight of rice. Azoxystrobin 7.5% + Propiconazole 12.5% w/v SE @ 500 ml/ha was at par with higher dose and resulted better yield than other treatments.

### **REFERENCES:**

- Bag, M. K., 2009, Efficacy of a new fungicide 'Trifloxystrobin 25% + Tebuconazole 50%' 75WG against Sheath Blight (*Rhizoctonia solani* Kühn) of Rice. *Journal of Crop and Weed*, 5(1): 224-226.
- Bag, M. K. and Saha, S., 2009, Fungitoxic effect of Nativo 75WG (trifloxystrobin 25% + tebuconazole 50%) on grain discolouration (GD) disease of rice in West Bengal. *Pestology*, 33: 47-49.
- Bag, M. K., 2011, Performance of a new generation fungicide Metominostrobin 20SC against sheath blight disease of rice. *J. Mycopathol. Res.*, 49(1): 167-169.
- Bag, M. K., Yadav, M. and Mukherjee, A. K., 2016, Bioefficacy of Strobilurin based fungicides against rice sheath blight disease. *Transcriptomics*, 4: 128.
- Bhunkal, N., Singh, Ram and Mehta, N., 2015, Assessment of losses and identification of slow blighting genotypes against sheath blight of rice. *J. Mycol. Pl. Pathol.*, 45: 285-292.
- Bhuvanewari, V. and Raju, K. S., 2012, Efficacy of new combination fungicide against rice sheath blight caused by *Rhizoctonia solani* (Kühn). *J. Rice Res.*, 5: 57-61.
- Biswas, A. and Bag, M. K., 2010, Strobilurins in management of sheath blight disease of rice: A review. *Pestol.*, 34: 23-26.
- Dash, S. C. and Panda, S., 1984, Chemical control of rice sheath blight disease. *Indian Phytopath.*, 37: 79-82.
- Groth, D. E., 2005, Azoxystrobin rate and timing effects on rice sheath blight incidence and severity and rice grain and milling yields. *Plant Dis.*, 89: 1171-1174.
- Groth, D. E. and Bond, J. A., 2006, Initiation of rice sheath blight epidemics and effect of application timing of azoxystrobin on disease incidence, severity, yield, and milling quality. *Plant Disease*, 90: 1073-1076.
- Groth, D. E. and Bond, J. A., 2007, Effect of cultivars and fungicides on rice sheath blight, yield, and quality. *Plant Dis.*, 91: 1647-1650.
- International Rice Research Institute, (2002). Standard Evaluation System for Rice, 5th edn., Nov. 2002, INGER, Genetic Resources Center, IRRI, P.O. Box 933, 1099, Manila, Philippines, 56 pp.
- Jin, C., Wu, R., Hu, H., Jifeng, W. and Xiaoxia, W., 2013, Control effect of 6 types of azoxystrobin mixture on rice sheath blight. *Chin. Agric. Sci. Bull.*, Available at: [http://en.cnki.com.cn/Article\\_en/CJFDTOTAL-ZNTB201318031.htm](http://en.cnki.com.cn/Article_en/CJFDTOTAL-ZNTB201318031.htm), Accessed date: 22 April 2016.

- Jones, R. E., Belmar, S. B. and Jeger, M. J., 1987, Evaluation of benomyl and propiconazole for controlling sheath blight of rice caused by *Rhizoctonia solani*. *Plant Disease*, 71: 222-229
- Kandhari, J. and Gupta, R. L., 2003, Efficacy of fungicides and resistance inducing chemicals against sheath blight of rice. *Journal of Mycological Research*, 41: 67-69
- Kandhari, J., Gupta, R. L. and Kandari, J., 2003, Efficacy of fungicides and resistance inducing chemicals against sheath blight of rice. *J. Mycological Res.*, 41
- Kannaiyan, S. and Prasad, N. N., 1984, Effect of foliar spray of certain fungicides on the control of sheath blight of rice. *Madras Agric. J.*, 71: 111-114.
- Kumar, P., Ahlawat, S., Chauhan, R., Kumar, A., Singh, R. and Kumar, A., 2018, *In vitro* and field efficacy of fungicides against sheath blight of rice and post-harvest fungicide residue in soil, husk, and brown rice using gas chromatography-tandem mass spectrometry. *Environ. Monit. Assess.*, 190: 503.
- Kumar, P. M. K., Gowda, S. D. K., Rishikant, M., Kumar, K. N., Gowda, P. K. T. and Vishwanath, K., 2013, Impact of fungicides on rice production in India. In: Fungicides – showcases of integrated plant disease management from around the world, pp. 77-98.
- Prasanna Kumar, M. K., Veerabhadraswamy, A. L., 2014, Appraise a combination of fungicides against blast and sheath blight diseases of paddy (*Oryzae sativa* L). *J. Exp. Biol. Agri. Sci.*, 2: 49–57.
- Margani, R. and Widadi, S., 2018, Utilizing *Bacillus* to inhibit the growth and infection by sheath blight pathogen, *Rhizoctonia solani* in rice. *IOP Conf. Ser. Earth Environ. Sci.* 142:012070.
- Naik, R. G., Jayalakshmi, K., and Naik, T. B., 2017, Efficacy of fungicides on the management of sheath blight of rice. *Int. J. Curr. Microbiol. Appl. Sci.*, 6: 611–614.
- Ou, S. H., 1985, Rice Diseases, 2<sup>nd</sup> Edn. Commonwealth Mycological Institute, Surrey.
- Pan, X., Zou, J., Chen, Z., Lu, J., Yu, H., Li, H., 1999, Tagging major quantitative trait loci for sheath blight resistance in a rice variety, Jasmine 85. *Chin. Sci. Bull.*, 44: 1783–1789.
- Phelp, R. H. and Soto, A., 1993, Rice diseases at CARONI and approaches to the control In Proc. Monitoring tour and workshop on Integrated Pest Management in rice in Caribbean. held in Guyana and Trinidad and Tobago. 7-11 October 1991
- Rajan, C. P. D., 1987, Estimation of yield losses due to sheath blight of rice. *Indian Phytopathol.*, 40:174-177.

- Rashid, M. M., Bhuiyan, M. R., Dilzahan, H. A., Hamid, M. A., Hasan, N., Khan, M. A. I., 2020, Biological control of rice sheath blight disease (*Rhizoctonia solani*) using bio-pesticides and bio-control agents. *Bangladesh Rice J.*, 24: 47-58.
- Robinson, E., 2013, Resistant sheath blight changing practices in Louisiana. Available at: <http://deltafarmpress.com/rice/resistant-sheath-blight-changing-practices-louisiana>, Accessed date: 24 July 2015.
- Roy, A.K., 1993, Sheath blight of rice in India. *Indian Phytopath.*, 46: 97-205.
- Savary, S., Teng, P. S., Willocquet, L. and Nutter. F. W. Jr., 2006, Quantification and modeling of crop losses: A review of purposes. *Annual Rev. Phytopathol.*, 44: 89-112.
- Savary, S., Willocquet, L., Elazegui, F. A., Castilla, N. and Teng, P. S., 2000, Rice pest constraints in tropical Asia: Quantification and yield loss due to rice pests in a range of production situations. *Plant Dis.*, 84: 357-369.
- Shahid, A. A., Shahbaz, M. and Ali, M., 2014, A comparative study of the commercially available fungicides to control sheath blight of rice in Lahore. *J. Plant Pathol. Microb.*, 5: 2157-7471.
- Singh, R. and Sinha, A. P., 2004, Comparative efficacy of local bioagents, commercial bioformulation and fungicide for the management of sheath blight of rice, under glass house conditions. *Indian Phytopath.*, 57: 494-496.
- Singh, R., Sunder, S. and Kumar, P., 2016, Sheath blight of rice: current status and perspectives. *Indian Phytopathol.*, 69: 340-351.
- Singh, S. K., Shukla, V., Singh, H. and Sinha, A. P., 2004, Current status and impact of sheath blight in rice (*Oryza sativa* L.) - a review. *Agric. Rev.*, 25(4): 289-297.
- Uppala, S. and Zhou, X. G., 2018, Field efficacy of fungicides for management of sheath blight and narrow brown leaf spot of rice. *Crop Prot.*, 104: 72-77.
- Visalakshmi, V., Raju, M. R. B., Rao, A. U., Kumar, K. M. and Satyanarayana, N. H., 2016, Compatibility and efficacy of insecticide and fungicide combinations on major pests and sheath blight of paddy. *Nat. Env. Poll. Tech.*, 15: 233-235.
- Webster, R. K. and Gunnell, P. S., 1992, *Compendium of Rice Diseases*. St Paul, MN: American Phytopathological Society, viii-62.
- Wheeler B. E. J., 1969, *An introduction to plant diseases*. John Wiley and Sons Ltd; London, UK, p. 254.
- Yellareddygari, S. K. R., Reddy, M. S., Klopper, J. W., Lawrence, K. S. and Fadamiro, H., 2014, Rice sheath blight: a review of disease and pathogen management approaches. *J. Plant Pathol. Microbiol.*, 5: 1000241.

- Zheng, A., Lin, R., Zhang, D., Qin, P., Xu, L., Ai, P., Ding, L., Wang, Y., Chen, Y. and Liu, Y., 2013, The evolution and pathogenic mechanisms of the rice sheath blight pathogen. *Nat. Commun.*, 4: 1424.
- Zhou, X. G. and Jo, Y. K., 2014, Disease management. In: Way, M. O., McCauley, G. M., Zhou, X. G., Wilson, L. T. and Brandy, M., (Eds.), 2014, *Texas Rice Production Guidelines*. Texas Rice Research Foundation, pp. 44–57.

UNDER PEER REVIEW

**Table 1. Effect of Azoxystrobin 7.5% +Propiconazole 12.5% SE on sheath blight of rice during *Kharif* - 2011 (1<sup>st</sup> season)**

Sl. No.	Treatment	Dose a.i. (g/ha)	Dose Formulation (ml/ha)	Sheath blight intensity on rice				Grain Yield (q/ha)
				Initial score	10 days after I spray	10 days after II spray	Terminal score (10 days after III spray)	
1	Azoxystrobin 7.5% + Propiconazole 12.5% SE	28.1+46.9	375	4.63 (12.43)*	12.78 (20.94)	17.22 (24.52)	25.19 (30.12)	59.40
2	Azoxystrobin 7.5% + Propiconazole 12.5% SE	37.5+62.5	500	4.07 (11.64)	6.67 (14.96)	12.41 (20.62)	14.07 (22.03)	63.50
3	Azoxystrobin 7.5% + Propiconazole 12.5% SE	46.9+78.1	625	4.44 (12.17)	6.11 (14.31)	11.85 (20.14)	13.70 (21.73)	64.60
4	Azoxystrobin 23% SC	125	500	4.26 (11.91)	13.15 (21.26)	20.37 (26.83)	26.11 (30.73)	59.10
5	Propiconazole 25% EC	125	500	4.07 (11.64)	8.96 (17.42)	14.15 (22.10)	18.37 (25.38)	61.00
6	Validamycin 3% L	60	2000	4.26 (11.91)	10.56 (18.96)	16.30 (23.81)	20.74 (27.09)	60.70
7	Control	-	-	4.81 (12.68)	26.30 (30.85)	33.52 (35.38)	48.70 (44.26)	54.10
	CD at 5% level			NS	2.40	1.38	3.30	2.48

\* Figures in the parentheses represent arcsine transformed values

**Table 2. Effect of Azoxystrobin 7.5% +Propiconazole 12.5% SE on sheath blight of rice during *Kharif* - 2012 (2<sup>nd</sup> season)**

Sl. No.	Treatment	Dose a.i. (g/ha)	Dose Formulation (ml/ha)	Sheath blight intensity on rice				Grain Yield (q/ha)
				Initial score	10 days after I spray	10 days after II spray	Terminal score (10 days after III spray)	
1	Azoxystrobin 7.5% + Propiconazole 12.5% SE	28.1+46.9	375	5.19 (13.16)*	15.37 (23.08)	19.81 (26.43)	29.07 (32.63)	57.50
2	Azoxystrobin 7.5% + Propiconazole 12.5% SE	37.5+62.5	500	5.37 (13.40)	9.63 (18.08)	15.93 (23.52)	16.48 (23.66)	62.20
3	Azoxystrobin 7.5% + Propiconazole 12.5% SE	46.9+78.1	625	5.00 (12.92)	8.70 (17.16)	15.19 (22.93)	16.11 (23.66)	63.20
4	Azoxystrobin 23% SC	125	500	5.56 (13.63)	18.52 (25.49)	25.37 (30.24)	30.56 (33.56)	57.60
5	Propiconazole 25% EC	125	500	5.37 (13.40)	11.93 (20.21)	18.85 (25.73)	22.37 (28.23)	59.00
6	Validamycin 3% L	60	2000	5.74 (13.86)	14.81 (22.64)	20.00 (26.57)	24.81 (29.88)	58.70
7	Control	-	-	5.74 (13.86)	29.81 (33.10)	34.44 (35.94)	57.96 (49.58)	52.40
	CD at 5% level			NS	2.12	2.18	3.24	3.00

\*Figures in the parentheses represent arcsine transformed values