

## **Effect of Bio control agents in the management of citrus nematode (*T.semipenetrans*) on Khasi mandarin seedling.**

### **ABSTRACT**

Studies on the effect of *Purpureocillium lilacinum*, *Glomus fasciculatum*, *Trichoderma harzianum*, vermicompost and carbofuran 3G in the management of *Tylenchulus semipenetrans* on Khasi mandarin under pot condition showed that all the treatments were effective in increasing plant growth parameters of Khasi mandarin. All the treatments significantly decreased final nematode population in soil and root over control. The treatment with vermicompost @ 15g/plant + *T. harzianum* 10g/plant found to be most effective in plant growth parameters of Khasi mandarin. The treatment with carbofuran 3G @ 4g/plant was found to be best in suppressing the final nematode population in soil and root followed by the treatment with *G. fasciculatum* @ 150 spores/plant + *P. lilacinus* @ 10g/plant.

**Key word:** *Purpureocillium lilacinum*, *Glomus fasciculatum*, *Trichoderma harzianum*, vermicompost, Khasi mandarin.

### **INTRODUCTION**

Citrus is one of the most important fruit crops particularly in the tropical and sub tropical areas of the world. It includes different citrus fruit crops like oranges, lemons, grape and limes. In Assam lemon, Sweet lemon (Mousambi) and orange occupied an area of 16058, 385 and 12680 ha with the production of 160339, 6056 and 179168 metric tonne respectively (Anon. 2023). In Dibrugarh district, it covers 468 ha of Assam lemon with production of 4067 metric tonne and oranges 75 ha with the production of 966 metric tonne. (Anon., 2015). The citrus nematode, *Tylenchulus semipenetrans* (Cobb) has been recognized as one of the greatest threat throughout the world which limits production of citrus under a wide range of environmental and edaphic conditions (Parvez *et al.*, 2003). The pest is cosmopolitan in distribution and has been reported to caused losses up to 43.3% on *C. aurantium* (Garcia Teran *et al.*, 1987).

In the State of California, *T. semipenetrans* was found to cause 10-60% reduction on growth and 10-20% in yields of lemon and oranges (Baines and Clarke, 1952; Baines, 1954; Baines and Martin, 1957).

Emphasis is now focused on the biological method of nematode control, as the nematicides may have toxic effect on non-target soil microfauna and flora, resulting in ecological imbalance. Among various biocontrol agents, *Purpureocillium lilacinum*, *Glomus fasciculatum* and *Trichoderma harzianum* play an important role in nematode management.

Among the nematode destroying fungi, there is an important group of fungi, which are capable of rapidly colonizing nematode reproductive structures and thus causing destruction of females, cysts and eggs. The common genera of such fungi are *Cylindrocarpan*,

*Exophiala*, *Fusarium*, *Gliocaladium*, *Purpureocillium*, *Phoma*, *Torula*, *Verticillium*, etc. The fungus *P.lilacinus*, a common soil hyphomycete has demonstrated tremendous potential as biocontrol agent of nematode (Jatala, 1986).

VAM fungi and plant parasitic nematodes are commonly found inhabiting the rhizosphere of the same plant, each having a characteristic but opposite effect on plant vigour. VAM fungi may stimulate plant growth, whereas the obligate plant parasitic nematodes usually suppress plant growth. It has been observed that the plants colonized with mycorrhizal fungi grow well in spite of the presence of damaging level of plant parasitic nematodes and this may be due to improved plant nutrition especially phosphorus (Hussey and Roncadori, 1982).

Ravichandra (2008) reported that *Trichoderma* sp. is one of the most commonly used and a popular bioagents against nematodes. Mechanisms involved in antagonism of *Trichoderma* sp. against nematode management are mycoparasitism, antibiosis, competition, tolerance to biotic and abiotic stresses through an enhanced root development, solubilisation and sequestration of inorganic plant nutrients and induce resistance.

Application of vermicompost has a positive effect on plant nutrition, photosynthesis, the chlorophyll content of the leaves and improves the nutrient content of the different plant components and improve the plant quality and act as a deterrent to pests and diseases. (Theunissen *et al.*, 2010). Domínguez *et al.* (2010) investigated the potentiality of vermicompost in the suppression of plant pathogens, insects, mites and plant parasitic nematodes. He reported that the addition of vermicompost to soils increases microbial diversity, providing a wider range of microorganisms that can act as biocontrol agents against different plant pathogens including nematodes.

Very little work has so far been done on citrus crop in Assam, except work Moznur (1997) on management in Khasi mandarin (*Citrus reticulata* Blanco), Deka (2000) on pathogenecity and management on *Citrus jambhiri*. Therefore present investigation is an attempt to study the management of citrus nematode, *T. semipenetrans* on Khasi mandarin.

## MATERIALS AND METHODS

The pot experiment was carried out in the net house of Department of Nematology to study the effect of *Purpureocillium lilacinum*, *Glomus fasciculatum*, *Trichoderma harzianum*, vermicompost and carbofuran 3G in the management of citrus nematode, *Tylenchulus semipenetrans* on khasi mandarin. Khasi mandarin (*Citrus reticulata*) seedlings were planted in 3kg capacity pot filled with sterilized soil and inoculated with *T. semipenetrans* @ 1 nematode per gram of soil. The pots were arranged in completely randomized design (CRD) with 5 replications for each treatment. Vermicompost and carbofuran 3G were applied 15 days and 3 days before planting respectively. The formulation of *Purpureocillium lilacinum* and *Trichoderma Harzianum* was applied at the time of planting according to the treatments. Infested soil of *G. fasciculatum* was applied at the time of planting.

The treatments were :

T1 = *Purpureocillium lilacinum* @ 20g/plant;

T2 = *Glomus fasciculatum* @ 300 spores/plant;

T3 = Vermicompost @ 15g/plant + *Trichoderma harzianum* 10g/plant;

T4 = *Glomus fasciculatum* @ 150 spores/plant + *Purpureocillium lilacinum* @ 10g/plant ;

T5 = Vermicompost @ 15g/plant + *Purpureocillium lilacinum* @ 20g/plant;

T6 = Carbofuran 3G @ 4 g/plant (W/W).

T7 = Control (Nematode alone).

Observations were made on length of shoot and root, fresh weight of shoot and root, final nematode population in soil and root after six month of nematode inoculation.

## RESULT AND DISCUSSION

In the present investigation, the result revealed that all the treatments were found to be effective in increasing the plant growth parameters of khasi mandarin seedlings. Maximum increase in plant growth parameters viz., shoot length (38.46cm), fresh weight of shoot (14.30g) and root length (31.44cm) were recorded in the treatment with vermicompost @ 15g/plant + *Trichoderma harzianum* @ 10g/plant (Table 1). The results were in agreement with Theunissen *et al.* (2010) who observed that vermicompost contains plant nutrients including N, P, K, Ca, Mg, S, Fe, Mn, Zn, Cu and B, and the uptake of which has a positive effect on plant nutrition, photosynthesis, the chlorophyll content of the leaves and improves the nutrient content of the different plant components (roots, shoots and the fruits). High percentage of humic acids in vermicompost contributes to plant health, as it promotes the synthesis of phenolic compounds such as anthocyanins and flavonoids which may improve the plant quality and act as a deterrent to pests and diseases. The increased growth parameter of plants observed in the present study may also be due to the use of *Trichoderma* as it has the ability to release plant growth promoting substances during its multiplication in the rhizosphere like indole 3-acetic acid, indole 3-acetaldehyde, indole 3-ethanol which may play role in mediating plant growth promoter (Cornejo *et al.*, 2009). Weeder *et al.* (2008) also reported that *Trichoderma* is highly rhizosphere competent i.e., able to colonize on roots as it develops, thus promote plant growth. Bio control agents *Pseudomonas fluorescens*, *Trichoderma viride*, *T. harzianum* and *Pochonia chlamydosporia* are effective to reduce the citrus nematode population in nine - year old citrus, *Citrus limonia* L. garden and to increase the fruit yield of *Citrus limonia* L. (Deepa *et al.*, 2011).

The minimum final nematode population in soil (78) was recorded in the treatment with carbofuran 3G @ 4 g/plant followed by the treatment with *G. fasciculatum* @ 150 spores/plant + *P. lilacinus* 10g/plant (110). All the treatments were effective in decreasing nematode population in soil over control. (Table-2)

In regards to the final nematode population in root, minimum nematode population in (34.00) was recorded in the treatment with carbofuran 3G @ 4 g/plant followed by the treatment with *G. fasciculatum* @ 150 spores/plant + *P. lilacinus* 10g/plant (44.00). All the treatments were effective in decreasing nematode population in root over control.

Khan (2008) reported that the hyphal network of VAM increases the absorption area of root system and therefore aids in water uptake, absorption and translocation of nutrients particularly phosphorus. The plants heavily colonized with AM fungi are able to grow well in spite of the presence of damaging levels of nematodes. The favourable effect of mycorrhizae in decreasing the disease intensity in nematode affected plants has been demonstrated in various crops. Baghel *et al.* (1990) reported that VAM inoculation stimulated the growth of *Citrus jambhiri* seedlings, while the citrus nematode, *Tylenchulus semipenetrans* decreased growth. When the 2 organisms were inoculated simultaneously the adverse effect of the

nematode partly nullified by the fungus and limited the development of the nematode. Reddy *etal.* (1995) observed that the inoculation of endomycorrhizal fungi *G. fasciculatum* along with soil amended with neem cake had reduced the population of *T. semipenetrans* on acid lime. Esser and El-Gholl (1993) reported that vegetative hyphae of *P. lilacinus* enter the gelatinous matrix of root knot nematode, or grow into the vulva or open cyst nematode. When it enters inside a penetration peg forms and grow in to the eggs shell. Penetrated eggs swell and buckle. Moznur (1997) studied the efficacy of *Paecilomyceslilacinus* against *T. semipenetrans* on khasi mandarin at different levels showed increased plant growth as well as reduced nematode population effectively with the increase in the level of *P. lilacinus*. Walode*etal.* (2008) observed that integration of *Paecilomyceslilacinus*(4g/kg soil), *Trichodermaharzianum*(4gm/kg soil) and *Glomusfasciculatum* (500spores/kg soil)was effective in increasing plant growth parameters and in reducing citrus nematode population both in soil and roots of rough lemon . Mahanta *et al.*(2016)found that application of *P.lilacinus* infected grain cause significant reduction in nematode population in soil and root over control .The minimum nematode population was recorded in the treatment with *P.lilacinus* infected grain@30gm/tree basin .Seenivasan *et.el* (2021) observed the individual and combined effect of neem seed kernel extract(NSKE) and liquid Purpureocilliumlilacinum formulation delivered through drip irrigation system on acid lime trees naturally infested with *Tylenchulussemipenetrans*.

Both bioproducts were applied at 4l/ha,but NSKE alone delivered twice at 30 days interval and their effect was compared with carbofuran 3G at 100g/tree as spot application. Results revealedthat both NSKE and *P.lilacinum*tested,significantly suppressed *T.semepenetrans* density in soil and acid lime roots. This combined treatment reduced *T.semepenetrans* density in soil by 70.0-88.8% and in roots by 73,6-79.2%. The integrated approach caused 26.3-33.6% more fruit yield with higher(1:3:2-1:4:2) cost benifit ratio.

Kumar(2021)reported that various microbes and soil-dwelling predatory mites (Acari: Mesostigmata) have been evaluated as augmentative biological control agents against citrus parasitic nematodes.They include fungi (Trichoderma spp., Purpureocilliumlilacinum, Pochoniachlamydosporia, and mycorrhizae, Glomus spp.) and bacteria (Bacillus spp., Pseudomonas fluorescens, Streptomyces avermitilis, and Pasteuria spp.), that have been applied against parasitic nematodes in different citrus cultivars. Integration of bionematicides with other management tools as well as the conservation and enhancement of indigenous antagonists, will further enhance the management of citrus parasitic nematodes.

Non-chemical practices which could substantially help in preventing and controlling *Tylenchulussemipenetrans*.This approach employs preventive techniques such as include eradication of infected plants, crop husbandry, sanitation, soil disinfection and solarization, incorporation of organic soil amendments, and crop rotation and scions, and unfavorable horticultural features (Reza Ghaderi & Manouchehr Hosseinvand, 2022)

## CONCLUSION

In the present investigation the treatment with with vermicompost @ 15g/plant + *Trichoderma harzianum*@ 10g/plant was found to be best in increasing plant height, root length, fresh weight of shoot and root. All the treatments were found to be effective in increasing the plant growth parameters and reducing the nematode population in soil and root over control. Among the biocontrol treatments *G. fasciculatum*@ 150 spores/plant + *P. lilacinus*@ 10g/plant was found to be effective in suppressing the nematode population in soil as well as in root.

UNDER PEER REVIEW

**Table 1: Effect of Bio control agentson the growth parameters of Khasi mandarin**

Sl No.	Treatments	Shoot length (cm)	Root length (cm)	Fresh shoot weight (g)	Fresh root weight (g)
1	T <sub>1</sub>	28.32 <sup>d</sup>	24.90 <sup>c</sup>	9.80 <sup>d</sup>	3.54 <sup>c</sup>
2	T <sub>2</sub>	33.56 <sup>c</sup>	27.78 <sup>b</sup>	11.02 <sup>bcd</sup>	5.04 <sup>b</sup>
3	T <sub>3</sub>	38.46 <sup>a</sup>	31.44 <sup>a</sup>	14.30 <sup>a</sup>	5.42 <sup>a</sup>
4	T <sub>4</sub>	35.16 <sup>bc</sup>	27.32 <sup>b</sup>	11.92 <sup>bc</sup>	5.28 <sup>b</sup>
5	T <sub>5</sub>	37.16 <sup>ab</sup>	30.36 <sup>a</sup>	12.36 <sup>b</sup>	5.36 <sup>a</sup>
6	T <sub>6</sub>	30.14 <sup>d</sup>	23.64 <sup>c</sup>	10.26 <sup>cd</sup>	2.98 <sup>c</sup>
7	T <sub>7</sub>	18.18 <sup>e</sup>	16.14 <sup>d</sup>	5.68 <sup>e</sup>	2.00 <sup>d</sup>
	S.Ed(±)	1.14	0.99	0.79	0.25
	CD at 5%	2.34	2.03	1.63	0.52

Note: Values of shoot and root length, fresh weight of shoot and root within parentheses are square root ( $\sqrt{x+0.5}$ ) transformed data

Mean followed by the same letter in the superscript(s) are not significantly different

**Table 2: Effect of Bio control agentson development of *T.semipenetrans* on Khasi mandarin**

Sl No.	Treatments	Soil nematode population (200 cc)	%Decrease over control	Total Nematode population per root system (2g)	%Decrease over control
1	T <sub>1</sub>	138 (12.21) <sup>bc</sup>	63.87	58 (8.07) <sup>bc</sup>	60.8
2	T <sub>2</sub>	130 (11.87) <sup>bc</sup>	65.96	56 (7.95) <sup>bc</sup>	62.16
3	T <sub>3</sub>	146 (12.55) <sup>b</sup>	61.78	60 (8.19) <sup>b</sup>	59.45
4	T <sub>4</sub>	110 (10.94) <sup>c</sup>	71.2	44 (7.13) <sup>cd</sup>	70.27
5	T <sub>5</sub>	128 (11.76) <sup>bc</sup>	66.49	52 (7.68) <sup>bc</sup>	64.86
6	T <sub>6</sub>	78 (9.28) <sup>c</sup>	80	34 (6.31) <sup>d</sup>	77.02
7	T <sub>7</sub>	382 (20.01) <sup>a</sup>	-	148 (12.64) <sup>a</sup>	-
	S.Ed(±)	0.63		0.44	
	CD at 5%	1.30		0.91	

Note: Values of final nematode population within parentheses are square root ( $\sqrt{x+0.5}$ ) transformed data  
Mean followed by the same letter in the superscript(s) are not significantly different

## REFERENCES

1. Anonymous, (2015). District wise area, production and average yield of some horticultural crops of Assam during the year 2015-16. Directorate of Horticulture & FP, Assam, Khanapara, Guwahati-22.
2. Anonymous (2023). Statistical Handbook of Assam- 2022-23. Table 5.15 PP. 69.
3. Baines, R. C. and Clarke. O. F. (1952). Citrus root nematode .Calif. Agric 6(9):13.
4. Baines, R.C. and Martin, J.P. (1957).Control of citrusnematode and Phytophthora sp. By Vapam. Plant Dis. Repr.41(5): 405-414.5.Baines, R.C.; Martin, J.P.; De Wolfe, T.A.; Boswell, S.B. and Garber, M.J. (1962). Effect of high doses of D-D on soil organisms and the growth and yield of lemon tress.Phytopathology52:723
- 6.Christie, J.R. and Perry, V.G. (1951). Removing nematodes from soil. Proc. Helminth Soc. Wash.18: 106-108.
7. Cornejo, Macias, R.Z.; Penages, C. and Buciol, J. (2009), Trichoderma virens, a plant beneficial fungus. enhances biomass production and promotes lateral root growth through an auxin dependent mechanism in arabidopsis. Plant physiol. 149: 1579-1592
8. Crozzoli, R.; Lamberti, F.; Greco, N. and Rivas, D. (1998). Plant parasitic nematode associated with citrus in venezeulea. Nematol Medit.26: 31-58
9. Deka, R. (2000), Pathogenicity and management of *Tylenchulussemipenetrans* (Cobb, 1913) on Citrus jambhiri. UnpublishedM.Sc.(Agri) Thesis, Assam Agril Univ., Jorhat.
10. Deepa, S.P.; Subramanian S. Ramakrishnan S. (2011). Biomangement of citrus nematode Tylenchulussemipenetrans Cobb on lemon, citrus limonia L. Journal of biopesticides., 4(2): 205-207
11. Dominguez, J.; Gomez-Brandon, M. and Lazcano, C. (2010). Bio-pesticide properties of vermicompost. Acta Zoologica Mexicana 26(2): 373-383
12. Duncan, L.W. and E. Cohn (1990). Nematode parasites of citrus", In: Luc, R., R.A. Sikora and J. Bridge (Eds), "Plant Parasitic Nematodes in subtropical Agriculture. CAB Int'l. pp. 321-346
13. Garcia- Teran, O.; Duarte, E. and Jimenez, Y.E. (1987). Relation between the population density of Tylenchulussemipenetransand the yield of "Valencia" oranges (Citrus sinensis). Ciencia y Tecniaen la Agricultura, protection de plants 10(1): 63-71
14. Hussey, R.S. and Roncadori, R.W. (1982). Vesicular arbuscular mycorrhizal may limit nematode activity and improve growth. *Plant Dis.* 66: 9-14
15. Jatala, P. (1986). Biological control of plant parasitic nematodes. *Ann. Rev. Phytopathol.* 24: 453-489
16. Khan, M.R. (2008). Plant Nematodes. Oxford & IBH publishing Co. pvt. Ltd., New Delhi, pp. 263-268
17. Kiran Kumar (2021) Recent advances in the biological control of citrus nematode: A review, DOI: 10.1016/j.biocontrol.2021.104593.
18. Moznur, S. (1997). Management of citrus nematode (Tylenchulussemipenetrans Cobb,1993) on Khasi Mandarin (*Citrus reticulata* Blanco). UnpublishedM.sc.(Agri) Thesis, Assam Agril Univ., Jorhat.pp- 52-55.
19. Mahanta, Bornali; Borbora,A.CandKonwar,Binita(2016)Effect of Paecilomyceslilacinus on Tylenchulussemipenetrans population in Khasimandarin .Indian J.Nematol.46(1)55-56

20. Ravichandra, N.G. (2008) Nematode management. In: Plant nematology. I.K. International Publishing House Pvt. Ltd., New Delhi, pp.550-551.
- 22.(Reza Ghaderi & Manouchehr Hosseinvand, (2022). Non- chemical Management of the Citrus Nematode, *Tylenchulus semipenetrans* (Nematoda: Tylenchulidae). Part of the Sustainability in Plant and Crop Protection book series (SUPP, volume 18).
21. Theunissen, J.; Ndakidemi, P.A. and Laubscher, C.P. (2010). Potential of vermicompost produced from plant waste on the growth and nutrient status in vegetable production. *Int. J. of the Physical Sciences*. 5(13); 1964-1973
22. Walode, N.B.; Sinha. A.K. and Neog, P.P.(2008). Biological control of citrus nematode *Tylenchulus semipenetrans* on Citrus jambhiri. *Indian J. Nematol.* 38(2): 244-255

UNDER PEER REVIEW