

AN EXPERIMENTAL STUDY TO ASSESS THE EFFECTIVENESS OF NORMAL SALINE FLUSH VERSUS HEPARIN FLUSH IN MAINTANING THE PATENCY OF PERIPHERAL INTRAVENOUS CANNULA

ABSTRACT

An experimental study to assess the effectiveness of normal saline flush versus heparin flush in maintaining the patency of peripheral intravenous cannula in the patients receiving intermittent intravenous medication.

The conceptual framework which was adopted for this study was based on Ernestine Wiedenbach's helping art of clinical nursing theory (1964) known as prescriptive theory in a modified form.

The research approach adopted was a true experimental approach and the research design was multiple treatment, pre-test, post-test control group design. The independent variable for the study were normal saline flush and heparin flush & the dependent variables were patency of peripheral IV cannula.

The present study was conducted at medical unit of Sanjeevan Hospital Dariyaganj New Delhi.

The sample for the present study comprises of all the patients who were 18 years and above of age with peripheral intravenous cannula receiving intermittent IV medication.

The data was analysed using descriptive and inferential statistics in terms of frequency, percentage and chi square.

Patency of IV cannula was conducted in three experimental groups.

The findings of the study have implications for nursing practice. Flushing of IV cannula is very effective in maintaining the patency of peripheral IV cannula. Saline flushing and heparin flush are equally effective in maintaining the patency of peripheral IV cannula. Normal saline flush not only reduces the patient's

discomfort but also reduces the health care cost of patients.

A similar study can be done on patient receiving chemotherapy to see effect of saline flushing versus heparin flush on the severity of occurrence of chemotherapy induced phlebitis.

Keywords: Heparin, Saline, Cannula, Thrombosis

INTRODUCTION

Intravenous Catheterization is the most common invasive procedure among patients admitted to hospital. The insertion and care of peripheral intravenous catheters is a routine task for nurses in all kinds of care.

The non-potency of I.V. cannula is found to be the major cause of other local and systemic complications i.e. thrombosis and blood stream infections. Careful adherence to guidelines and procedures can minimize these risks.[1]

About 20-30% of patients receiving peripheral intravenous therapy develop local complications such as infiltration, thrombophlebitis and infection.[2]

Maintenance of patency of the IV catheters is essential as resisting a catheter may produce discomfort to patient and increased cost.[3]

Heparin sodium used to be the traditionally used medication as anticoagulant in those catheters in order to prevent clotting, minimise the incidence of phlebitis as it enhances antithrombin III activity and compensates coagulation factors XII, and X.[4]

Hephzibha, conducted a systemic review of the study on heparin versus normal saline flush solution.[5]

The hospital protocols for flushing IV line vary from no routine IV cannula flushing, flushing with 0.9% NS solution to the use of heparin solution in India there are lots of differences in maintaining the IV line patency, even in the same hospital.[6]

This study will improve the clinical practice and patient care in all the clinical areas of hospitals as the peripheral IV cannula is the most common invasive procedure done for the hospitalized patients.

An experimental study to assess and evaluate the effectiveness of normal saline flush versus heparin flush in maintaining the patency of peripheral intravenous cannula in the patients receiving intermittent intravenous medication in a selected hospital of New Delhi.

METHODS AND METHODOLOGY

INDEPENDENT VARIABLE

- i. Normal saline flush
- ii. Heparin flush

DEPENDENT VARIABLE

- i. Patency of cannula
- ii. Resistance to flush

In the present study population comprises of all the patients who are 18 years and above of age with peripheral intravenous cannula receiving intermittent IV medication.

SAMPLE CRITERIA

InclusiveCriteria

1. Patients who were 18 years and above of age.
2. Both male and female
3. Patient with peripheral IV cannula receiving intermittent IV medication 2 times a day.
4. Patient with peripheral IV cannula insertion not more than 36 hours.

Exclusive Criteria

1. Patients below 18 years of age
2. Patient who have active bleeding
3. Patient who are on anticoagulant therapy
4. Patient who had history of hypersensitivity to any anticoagulant therapy
5. Patient who were on chemotherapy
6. Patients who bleeding disorders
7. Patients who were on intravenous fluids

SAMPLE SIZE

Pilot study: 9 sample subjects, 3 each in experimental group 1, experimental group 2, and control

1.1	Male	9	45	11	55	8	40	28	47
1.2	Female	11	55	9	45	12	60	32	53
2	AGEINYEAR				0				
2.1	18-28	3	15	2	10	4	20	9	15
2.2	29-38	5	25	10	50	9	45	24	40
2.3	39-48	5	25	3	15	3	15	11	18
2.4	49-58	5	25	2	10	2	10	9	15
2.5	59 and above	2	10	3	15	2	10	7	12
3	DIAGNOSIS								
3.1	Respiratory diseases	3	15	2	10	2	10	7	11
3.2	cardiac diseases	1	5	2	10	2	10	5	8
3.3	Neurological diseases	2	10	2	10	2	10	6	10
3.4	Renal Diseases	2	10	2	10	3	15	7	12

3.5	Gastro enterological Diseases	2	10	2	10	3	15	7	12
3.6	Endocrinological Diseases	2	10	3	15	4	20	9	15
3.7	Others(Typhoid,Fever,HIV, Acute,Gastritis, Anaemia, Herpeszosteretc.	8	40	7	35	4	20	19	32
4	PREVIOUS HOSPITALIZATION (last one year)								
4.1	Yes	4	20	3	15	6	30	13	22
4.2	No	16	80	17	85	14	70	47	78
5	DURATION OF PREVIOUS HOSPITALIZATION								
5.1	Less than 5days	5	25	4	20	6	30	15	25
5.2	6-10 days	8	40	9	45	7	35	24	40
5.3	11-15 days	4	20	3	15	4	20	11	18
5.4	More than 15 days	3	15	4	20	3	15	10	17

Figure1: Pie diagram showing percentage distribution of patients as per their gender




TABLE-2. Frequency and percentage distribution of varioius clinical variables of the patients

N=60

S.no	CLINICAL VARIABLE	Experimental Group 1 (saline)n=20		Experimental Group 2 (heparin)n=20		Control groupn =20		Total'f	Total %
		f	%	f	%	f	%		
CALCULATION RELATED FACTOR									
1	IV CANNULA INSERTED BY								
1.1	Doctor	14	70	13	65	8	40	35	58
1.2	Nurse	6	30	7	35	12	60	25	42
2	SIZE OF CANNULA								
2.1	18G	2	10	3	15	4	20	9	15
2.2	20G	12	60	10	50	9	45	31	52
2.3	22G	6	30	7	35	7	35	20	33
3	DURATION OF IV CANNULA INSERTION								

3.1	Upto12 hrs.	6	30	7	35	6	30	19	32
3.2	13-24 hrs	10	50	9	45	8	40	27	45
3.3	24-36 hrs	4	20	4	20	6	30	14	23

PATIENT RELATED FACTOR

4	SITE OF CANNULA								
---	------------------------	--	--	--	--	--	--	--	--

4.1	Dorsum	7	35	7	35	6	30	20	33
4.2	Wrist	7	35	8	40	7	35	22	37
4.3	Forarm	4	20	4	20	5	25	13	22
4.4	Antecubital region	2	10	1	5	2	10	5	8

5	AVOIDANCE OF PREVIOUS SITE								
---	-----------------------------------	--	--	--	--	--	--	--	--

5.1	Yes	20	100	19	95	18	90	57	95
5.2	No	0	0	1	5	2	10	3	5

6	ACTIVITY OF PATIENT								
---	----------------------------	--	--	--	--	--	--	--	--

6.1	Completely ambulatory	0	0	0	0	2	10	2	3
6.2	Walk occasionally	9	45	12	60	10	50	31	52
6.3	Restricted activity	11	55	8	40	8	40	27	45

TREATMENT RELATED FACTORS

7	DURATION OF IV THERAPY								
---	-------------------------------	--	--	--	--	--	--	--	--

7.1	Upto12 hrs.	6	30	5	25	8	40	19	32
7.2	13-24 hrs	10	50	11	55	9	45	30	50
7.3	25-36 hrs	4	20	4	20	3	15	11	18

8	IV MEDICATION ADMINISTRATION								
---	-------------------------------------	--	--	--	--	--	--	--	--

8.1	Antibiotics through IV push	8	40	7	35	9	45	24	40
-----	-----------------------------	---	----	---	----	---	----	----	----

	Antibiotics through IV push + antibiotics through infusion	4	20	3	15	5	25	12	20
8.3	Antibiotics through IV push Corticosteroids	3	15	4	20	4	20	11	18
8.4	Antibiotics through IVpush+antibiotics through infusion+corticosteroids	5	25	6	30	2	10	13	22

UNDER PEER REVIEW

Data presented in **TABLE-2** ,reveals that

1. Majority of IV cannula's that is 58% were inserted by doctors and 42% were inserted by nurses
2. Data of IV cannula related factors shows that maximum patients were having cannula of 20G size that is 52%, followed by 22 G (33%) and minimum number of patients with 18 G size cannula(15%) only.
3. Data as regard to duration of cannulation and duration of therapy shows that, that the majority of patients had IV cannula insertion and duration of IV therapy between 13-24 hours(45%), followed by 32% up to 12 hours and 23% upto 25-36 hours.
4. As regard to the patient related factor, the data shows that the site of IV cannula insertion in most of the patient was wrist 37%, followed by 33% at dorsum, 22% on the fore arm and 8% at the antecubital region.
5. The data shows that the most of the patients were walking occasionally(52%) followed by 45% of patients with restricted activity and only 3% were completely ambulatory.

6. Data further shows that the majority of the patients were on antibiotics with IV push only 40%, 22% Antibiotics through IV push+antibiotics through infusion+corticosteroids, 20% Antibiotics through IV push + antibiotic through infusion and followed by 20% Antibiotics through IV push+antibiotic through infusion and 18% Antibiotics through IV push Corticosteroids.




Figure 2: Bar graph showing percentage distribution of the patients as per the size of IV cannula.




Figure 3: Conical graph showing percentage of patients as per the duration of IV cannula insertion.




Figure 4: Cylindrical graph showing percentage distribution of patients as per the site of IV cannula.

COMPARISION OF PATENCY OF IV CANNULA

This section describe the finding related to frequency and percentage of post intervention potency of peripheral IV cannula of patients receiving intermittent IV medication through peripheral IV cannula on 3 days of post intervention observation.[11,12]

TABLE-3

Frequency and percentage of post intervention patency of IV cannula in experimental groups and control group

N=60

Group	DAY 1(At 12 Hrs.)				DAY 2(At 36 Hours)				DAY3(At 60 Hours)			
	Patent		Non-Patent		Patent		Non-Patent		Patent		Non-Patent	
	f'	%	f'	%	f'	%	f'	%	f'	%	f'	%
1. Exp Group 1(Saline flush)n=20	20	100	0	0	20	100	0	0	18	90	2	10
2. Exp Group 2(Heparin flush)n=20	20	100	0	0	20	100	0	0	19	95	1	5
3. Control group (No flush) n=20	20	100	0	0	16	80	4	20	12	60	8	40

The data given in table shows that-

- Patency of IV cannula was found to be 100% among all the groups on day1
- Patency of IV cannula in experimental group1 was 100%, experimental group 2 was 100% and control group it was 80% on day2.
- Patency of IV cannula in experimental group 1 was 90 %, experimental group 2 was 95% and ,whereas it was 60% in control group on day3.

❖ The data of all 3 days at different time show that increase in time duration of peripheral IV cannulation results more number of blockage (non patency) in IV cannula of control group as compared to experimental group1 and experimental group2 in which IV cannula were flushed.

Figure 5
Bar graph showing the percentage distribution of patency of IV cannula in experimental groups and control group patients on day1.




Figure 6: Bar graph showing the percentage distribution of patency of IV cannula in experimental groups and control group patients on day 2





Figure 7: Bar graph showing the percentage distribution of patency of IV cannula in experimental groups and control group patients on day 3.

TABLE-4 Association between the post intervention patency of IV cannula of Experimental groups and Control group on day1,day2,and day3

N=60

DAY AND TIME	Experimental group1(saline flush)n=20		Ex. Gp. 2 (heparin flush)n=20		Control gp.=20		Degree of freedom	Chi sq. value	Chi square value obtained
	Patent	NonPatent	Patent	Non Patent	Patent	Non patent			
DAY1 at 12hr	20	0	20	0	20	0	2	5.99	4.1379 NS
DAY2 at 36hr	20	0	20	0	16	4			2.0339 NS
DAY3 at 60	18	2	19	1	12	8			10.00* Sig.

*significant at 0.05 level

NS-Non significant

The data given in theTable-4 revealed that:

1. There was significant association between experimental group1,experimental group2,and control group in maintaining the patency of IV cannula on day 1. As the obtained Chi-Square value 4.1379 was less than the table value.[13]
2. There was significant association between experimental group 1, experimental group 2, and control group in maintaining the patency of IV cannula on day 2.as the obtained Chi-Square value 2.0339 was less than the table value.

3. There was no significant association between experimental group 1, experimental group 2, and control group in maintaining the patency of IV cannula on day3.as the obtained Chi-Square value was more than the table value.[14]

- ❖ The data given in table 4 shows that with each day, the Chi-Square value is increasing due to loss of patency in control group. So, as the number of cannula blockage is increased in control group on 3rd day there was a significant difference found between experimental group and control group as evident from obtained Chi-Square value (10.00) which is more than the table value (5.99) this shows that initially the patency of IV cannula was same in experimental group and control group but with the increase in the time duration there was a difference found in the IV cannula patency in these groups,
- ❖ In order to determine the significant association between the patency of peripheral intravenous cannula line in experimental group 1 and experimental group 2, Chi-Square was calculated and these findings are presented in **Table 5**

TABLE-5 Association between the post intervention patency of IV cannula of Experimental group1and Experimental group 2 on day 1, day 2, and day3

N=60

DAY AND TIME	Experimental group1(saline flush)n=20		Experimental group2(heparin flush)n=20		Degree of freedom	Chi sq. value	Chi sq. Value
	Patent	Non Patent	Patent	Non Patent			
DAY1 (at12hrs)	20	0	20	0	2	5.99	4.13 NS
DAY2 (at36hrs)	20	0	20	0			4.13 NS
DAY3 (at60hrs)	18	2	19	1			2.034 NS

*Significantat0.05level

NS-Nonsignificant

The data given in table-5 revealed that:

1. There was a significant association found between experimental group 1 and experimental group 2 in maintaining the patency of IV cannula on day 1. The data obtained Chi-Square value (4.13) was less than the table value.

2. There was a significant association found between experimental group 1 and experimental group 2 in maintaining the patency of IV cannula on day 2. The data obtained Chi-Square value (4.13) was less than the table value.[15,16,17]

3. There was a significant association found between experimental group 1 and experimental group 2 in maintaining the patency of IV cannula on day 3. The data obtained Chi-Square value (2.034) was less than the table value.

- ❖ This data is given in table 5 shows that experimental group 1 (saline flush) and experimental group 2 (heparin flush) are showing similar result in terms of maintaining the patency of IV cannula of patient receiving intermittent IV medication. In all 3 days there was no statistically significant difference found between experimental group 1 and experimental group 2 as evident from obtained Chi-Square value which is very less than the table value (5.99) in all 3 days, so the association between experimental group 1 and 2 is true association and not by chance.

DISCUSSION OF THE FINDING

The present study was aimed to assess the effectiveness of normal saline flush versus heparin [18,19] flush in maintaining the patency of peripheral intravenous cannula in patients receiving intermittent IV medication in selected hospital of New Delhi.

In this section, the major findings of the present study have been discussed with reference to the results obtained by other investigators in the same aspect.

The finding of the present study showed that there is significant association between flushing of IV cannula and patency of peripheral IV cannula. [20,21] This finding is consistent with the findings of the research studies conducted by Choudhary, V.S [5] Vinoli. S (2007) and Kaur Maninder. et al. [19]

They reported that intermittent flush found to be highly effective and best method to be used in clinical studies to standardize the patency of cannula. [23]

The findings of the present study showed that there is an association between normal saline flush and heparin flush (10 unit dose) in maintaining the patency of peripheral intravenous cannula.

This finding is consistent with the finding of the research studies conducted by Kathryn M., et al.

(2003) and Harahan & Berends (2000) where they found that both normal saline flush and heparin flush (10 unit dose) are equally effective in the maintaining the patency of peripheral IV cannula

CONCLUSION

On the basis of the findings of the present study, the following conclusion can be drawn:

1. There was a significant association between patency of peripheral IV cannula and flushing of IVcannula.
2. There was a significant association between experimental group 1 (saline flush) and experimental group2(heparin flush) in maintaining the patency of peripheral IV cannula.
3. Maintenance of patency of peripheral IV cannula was depend on the type of IV medication used and it is independent of all other factors like size of IV cannula, site of IV cannula, duration of insertion ofIVcannulaand duration ofIV therapy.[24]

Present study has shown that the majority of patients in control group had blockage of peripheralIV cannula. Further the study has also identified thatflushing of peripheral IV cannula with saline flush and heparin flush has shown equal effect in maintaining the patency of IV cannula. The study emphasize the need to follow a standard protocol for flushing the peripheralIVcannulatoprevent blockage.

Present research indicates that saline flush and heparin flush give similar result in maintaining the patency of IV cannula, so the practice of using heparin as a flushing agent can be changed to normal saline flush which is safer and cheaper as compared to the heparin flush.

An exploratory study can be done to investigate the incidence of intravenous local vascular complications and to evaluate some important related factors.

REFERENCES

1. Ashton, J. et al. (1990) “ Effect of heparin versus saline solution on intermittent infusion device irrigation” *Heart and lung journal*, 19(6), 608-612
2. Barker, P .et al. (2004). “Randomized Clinical Trial of elective re-siting of intravenous cannula” *Annals R Coil Surg England*, 86(4), 281-283
3. Barret, P.J, Lester, R.L (1990) “Heparin versus saline flushing solution in a small community hospital”, *Hospital Pharmacy* 25(2), 115-118
4. Bregenzer, T., Conen, D, et al (1988) “Is routine replacement of peripheral intravenous catheters necessary?” *Arce of internal medicine*, 158(2), 151-156.
1. Campbell, S.G, et al (2005), “How often should intravenous catheter in ambulatory patients be flushed “, *Journals of Infusion Nursing*, 28(6), 399-404
5. Hephzibha, A (2010), a systemic review of the study on heparin versus normal saline solution I.
6. Patidar, A. 2013, The hospital protocols for flushing IV line vary from no routine IV flushing with 0.9% NS solution to the use of heparin solution in India.
7. Choudhry, V.S (2013), “Impact of Intermittent Saline Flushing in the Patency of Peripheral Venous catheter”, *International Journal of Nursing Education*, 5(2), 89-96
8. Cook, L., et al. (2011) “Heparinized saline vs normal saline for maintaining of intravenous access in neonates: an evidence-based practice change”. *Advanced Neonatal Care*. 11(3) 208-15.
9. Couzigou, C., et al. (2005). “Short Peripheral venous catheter: Effects of evidence based guidelines on insertion, maintenance and outcomes in the University Hospital”. *Journal of Hospital infections way and means*”, *Journal of Intravenous*

Nursing,19(4),175-

10. Crow ,S . (1996), “prevention of intravascular infections way and means”, *Journal of Intravenous Nursing*,19(4), 175-181

11. Dennis,M.M(2001).“Component IV management,*Nurse Manager*,32(8),21-23

12. Fujita, T., Suzuki, T.,et al. (2006), Normal saline flushing for maintenance of peripheral intravenous ,*Journal of Clinical Nursing*, 15(1) 103-104

13. Goldberg,M.,Sankaran.etal.(1999)“Maintaining Patency of Peripheral Infusion Devices with Saline and Heparinised Saline”.*Neonatal intensive care*,12(1),18-22

14. Goode,C.J.,Titler,M.etal(1991),“A meta-analysis of effects of heparin flush and saline flush:quality and cost implication”*Nursing Research*. 40(6), 324-30
Journal, 7(1), 22-33

15. Gupta,A.,Walia,I.etal(1988)“Epidemiology of Infiltration and Phlebitis”,*Nursing and Midwifery Research*

16. Heilskov J.,Kleiber C.,etal.(1988)“A randomized trial of heparin and saline for maintaining intravenous locks in neonates”,*Journal of the Society of pediatric Nurses*,3(3),111-116

17. Ingram,P.(2005).“Peripheral Intravenous therapy:Key risks and implications for practice”.*Nursing Standard*, 19(46), 55-64.

18. Kathryn,M.Niesen,etal(2003)“The effects of heparin versus normal saline for maintenance of peripheral intravenous locks in pregnant women”, *Journals of Obstetric Gynecologic & Neonatal Nursing*, 32(4), 503-508.

19 Kaur, Maninder.,et al. (October 2006) “ patency of intravenous cannula with ‘Normal Saline Lock’ for patients on intermittent medication’, *Nursing and Midwifery Research Journal*,2(4),132-138

20 Mitsiou,Maria. Et al. (2008) “ finding the evidence for keeping the patency in peripheral intermittent intravenous devices” *Health science journal*2(3), 121-128

21 Mok,E.,Kwong,T.k.,&Chan,MF.(2007).“A randomized controlled trial for maintaining peripheral intravenous lock in children”, *International Journal of Nursing Practice*,13(1), 33-45

22 Patidar, B.A (2014) “ Comparative Efficiency of Heparin saline and Normal saline

flushfor maintaining patency of peripheral intravenous lines”, *International journal of healthsciencesand research*, 4(3), 159-166

23.Rickard, C.M et al. (2010) “Routine resite of peripheral intravenous deviced every 3 days did not reduce complication compared with clinically indicated resite:a randomized controlled trail”*BMC Medicine*; 53(8)

24.Schultz, A.A Drew, D., et al (2002) “Comparison of normal saline and heparinized saline for patency of IV locks in neonates”. *Applied Nursing Research*. 15(1),28-34

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW

UNDER PEER REVIEW