

Original Research Article

INFLUENCE OF INTEGRATED NUTRIENT MANAGEMENT PRACTICES ON SOIL FERTILITY AND YIELD OF FODDER MAIZE

ABSTRACT

Field experiments were conducted during *rabiseason* of 2019 and 2020 at the Department of Agronomy, Madras Veterinary College, Chennai to study the effect of fodder maize (*Zea mays* L.) by adopting different organic manures and fertilizer levels. An experiment was laid out in split plot design. Main plots consisted of different organic manure treatments viz., No manure, farm yard manure (25 t ha⁻¹), vermicompost (12 t ha⁻¹) and poultry manure (12 t ha⁻¹). Sub plots consisted of varying level of fertilizer treatments viz., No fertilizer, 125% RDF, 100% RDF, 75% RDF. Recommended dose of fertilizer (RDF) for fodder maize is 60 kg N ha⁻¹: 40 kg P₂O₅ ha⁻¹: 20 kg K₂O ha⁻¹. The organic manures were applied as per the N equivalent basis to inorganic nutrient recommendation. The results revealed that all the soil parameters, green and dry fodder yield were significantly affected with the application of organic manures and fertilizer levels. Organic sources of nutrients tended to improve soil physico-chemical properties viz., bulk density, water holding capacity, porosity and organic carbon. The highest organic carbon content was noticed with the application of poultry manure (0.50 %) followed by poultry manure. The post harvest available soil nutrients was high in fodder maize grown with poultry manure along with 125% recommended dose of inorganic fertilizer to fodder maize recorded higher soil available nitrogen (109.48 kg ha⁻¹), phosphorus (15.46 kg ha⁻¹) and potassium (118.65 kg ha⁻¹). Significantly higher green and dry fodder yield of maize was recorded with application of poultry manure along with 75 % recommended dose of fertilizer.

Keywords: *Fodder maize, Soil pH, Organic carbon, Soil available nutrients, Fodder yield*

1. INTRODUCTION

Nowadays, crop production in the world is mainly affected by over use of land and poor soil fertility. It is a major threat to food security due to ever increasing growth rate of human population every year [1]. Land available for agricultural purposes is shrinking and demands for food to feed the growing population is increasing every year. Hence, the available cultivable lands should be used effectively with proper soil fertility management.

In order to increase crop yields and to meet our public needs, the inputs in the form of fertilizers, pesticides and weedicides along with intensive irrigation practices helped to achieve the target to a certain stage. However, decrease in crop yield, low availability of soil microorganisms and poor soil fertility took place despite the application of fertilizer. Moreover, ground water, air, human and animal health will also be affected by these chemicals directly and indirectly [2].

To minimize this adverse effect of chemical fertilizers, organic manures are being promoted now, giving rise to the concept of organic agriculture. Organic manures include Farm Yard Manure, Vermicompost, Poultry Manure, Sheep Manure and Goat Manure etc., Organic fertilizers have long since been known to improve physical and chemical properties, increase the humus content and biological properties of soil that help in flourishing of beneficial macro- and microorganisms [3]. Organic amendments increase soil humus, soil carbon and nitrogen content, which results in enhanced soil fertility and crop productivity and it is also eco-friendly and cost-effective.

In India, maize is cultivated as an important food crop next to rice and wheat. It can be cultivated throughout the year due to its wide adaptability and acceptability. Maize is used as food grain, feed and fodder, preparation of starch etc. Maize green forage, particularly when it contains the stalks, leaves and ears, is an energy-rich feed for ruminant livestock [4] (Maize is quick growing, yields high biomass, and is highly palatable. It contains sufficient quantities of protein and minerals and possesses high digestibility as compared to other non-legume fodders [5].

To meet the needs of ever increasing livestock population, the production and productivity of fodder crops to be increased. The area under fodder cultivation has been decreasing. Hence, the available land will be effectively used for increasing fodder production throughout the year. Though maize is a year round crop it will not produce more yield without proper nutrient management. Maize is an exhaustive crop and requires more amount nutrients for its growth and productivity. Only scanty information is available on integrated nutrient management of fodder maize. Hence, the present study was undertaken to increase maize fodder productivity and sustain the soil fertility by integrated application of organic manures and inorganic fertilizers to fodder maize.

2. MATERIALS AND METHODS

Field experiment was carried out at the Department of Agronomy, Madras Veterinary College, Chennai during the *Rabi* season of 2019 and 2020. The soil of the field was red sandy loam in texture classified under the order *Alfisol*s belonging to Typic Ustropept. pH was slightly alkaline (7.43) having an electrical conductivity of 0.28 dSm⁻¹. The organic carbon content was 0.26 per cent. The nutrient status of the soil was low in available Nitrogen (128.5 kg ha⁻¹), medium in available Phosphorus (25.3

kg ha⁻¹), and medium in available potassium (226.3 kg ha⁻¹). African tall variety is the maize variety used for this experiment.

The experiment was laid out in RCBD with four replications. The treatments comprised of different organic sources of nutrients such as No manure (M₁), Farm Yard Manure (M₂), Vermicompost (M₃) and Poultry Manure (M₄). Different level of fertilizers are used such as, No fertilizer (F₁), 125% RDF (F₂), 100% RDF (F₃) and 75% RDF (F₄). The recommended dose of fertilizer (RDF) used for fodder maize is 60: 40: 20 Kg NPK ha⁻¹, respectively. Organic manures were applied equivalent to recommended dose of nitrogen fertilizer basis.

2.1. SOIL PHYSICAL PROPERTIES

Bulk density of the soil was recorded by Keen's cup method developed by [6] for the soil samples collected after the harvest of the crop and expressed in g cm⁻³. Maximum water holding capacity of the soil was recorded by Keen's cup method developed by [6]. It was recorded after harvest of crops from each plot and expressed in percentage. Per cent of pore space of soil was recorded by Keen's cup method developed by [6]. It was also recorded after harvest of crops from each plot and expressed in percentage.

2.2. SOIL CHEMICAL PROPERTIES

Mechanical composition of the experimental field soil was analyzed as per the procedure suggested by [6] for the soil samples collected after the harvest crop. Soil pH and EC were estimated by using pH meter and Conductivity Bridge, respectively, in a soil: water suspension of 1: 2.5 ratio [7]. The soil was also analyzed for soil organic carbon [8].

The soil samples collected from 0-15 cm depth after the harvest of the crop were air dried, sieved through 2 mm mesh and used for available N, P, and K estimation. The available soil nitrogen was estimated by the method proposed by [9] and expressed in kg ha⁻¹. Available soil phosphorus was estimated by the procedure outlined by [10] and expressed in kg ha⁻¹. Available soil potassium was estimated by the method proposed by [11] and expressed in kg ha⁻¹.

2.3. FODDER YIELD

The maize plants were cut or harvested above the ground level and taken fresh weight by using weighing balance and recorded the yield in all plots having different treatments and expressed to t ha⁻¹ which gave green fodder yield of maize. For computing the dry fodder yield, the plant samples were dried using hot air oven at 65°C for 48 hours. The dried plant samples were weighed by using weighing balance and recorded dry fodder yield of maize (t ha⁻¹). All the parameters were analyzed using ANOVA mentioned by [12] F-value was tested at 5 %significance level.

3. RESULTS AND DISCUSSION

3.1. SOIL PHYSICAL PROPERTIES

Application of organic manures tended to improve soil physical properties viz, bulk density, water holding capacity and porosity of soil compared to initial status. Application of poultry manures

resulted in lower bulk density (1.38 g cm^{-3}) and higher water holding capacity (41.82 %) and porosity (43.90 %) after the harvest of fodder maize as compared to other organic manures (Table 2). This might be due to adding of poultry manure may also reduce compactness of soils or increased the looseness of soil resulting in increased soil volume as compared to that of other organic manures.

Table 2. Influence of bulky organic nutrients and fertilizers on physical parameters of soil after harvest of fodder maize (Pooled data)

Treatments	Bulk density (g cm^{-3})					Maximum water holding capacity (%)					Porosity (%)				
	F ₁	F ₂	F ₃	F ₄	Mean	F ₁	F ₂	F ₃	F ₄	Mean	F ₁	F ₂	F ₃	F ₄	Mean
M ₁	1.54	1.53	1.52	1.41	1.50	33.43	35.15	34.87	36.06	34.88	36.28	37.25	36.97	38.16	37.17
M ₂	1.54	1.49	1.45	1.48	1.49	33.32	36.04	35.93	36.89	35.54	35.45	37.37	38.06	39.02	37.48
M ₃	1.52	1.49	1.42	1.35	1.44	38.18	37.94	38.43	39.16	38.43	39.32	40.17	38.41	42.05	39.99
M ₄	1.41	1.39	1.38	1.35	1.38	40.23	41.65	42.15	43.24	41.82	42.52	43.94	44.44	44.70	43.90
Mean	1.50	1.47	1.44	1.40		36.29	37.69	37.85	38.84		38.39	39.68	39.47	40.98	
	M	F	M at F			M	F	M at F			M	F	M at F		
SE	0.035	0.035	0.069			1.039	0.743	1.486			1.164	0.837	1.674		
CD (p=0.05)	0.085	0.071	NS			2.54	1.53	NS			2.85	1.73	NS		

The reduced bulk density with the organic manures application was due to the improvement of soil aggregation, soil structural improvement and increased porosity [13]. Application of organic manures increased moisture content of the soil better than the chemical fertilizers due to its high organic carbon content and addition of organic matter of the soil. Organic matter has the ability to retain appreciable amounts of soil moisture as suggested by [14]. Similar results were also reported by [15].

With regard to fertilizers, application of 125% RDF recorded higher bulk density (1.39 g cm^{-3}), low water holding capacity (41.65 %) and porosity (43.94 %) than 75 % RDF. This might be attributed to deterioration of soil structure by higher dose of inorganic fertilizer application [16]).

3.2. SOIL CHEMICAL PROPERTIES

3.2.1. SOIL PH AND EC

The INM practice, organic manures and recommended NPK fertilizers application did not showed any significant difference on soil pH at post harvest stage of the soil (Table 3). However, slight decrease in pH was observed due to use of poultry manure which could have been due to their acidic nature. With regard to EC, 75% RDF recorded significantly lower EC followed by 100% RDF. These

results are in agreement with the findings of [17]. Treatments without any fertilizer recorded higher EC when compared to all other treatments.

Table 3. Influence of bulky organic nutrients and fertilizers on chemical parameters of soil after harvest of fodder maize (Pooled data)

Treatments	pH					EC (ds m ⁻¹)					Organic carbon (%)				
	F ₁	F ₂	F ₃	F ₄	Mean	F ₁	F ₂	F ₃	F ₄	Mean	F ₁	F ₂	F ₃	F ₄	Mean
M ₁	7.52	7.47	7.45	7.42	7.46	0.293	0.280	0.280	0.270	0.281	0.32	0.33	0.34	0.38	0.34
M ₂	7.43	7.45	7.44	7.35	7.42	0.280	0.280	0.270	0.270	0.275	0.42	0.43	0.44	0.46	0.44
M ₃	7.57	7.40	7.38	7.37	7.43	0.275	0.270	0.260	0.268	0.268	0.45	0.43	0.45	0.46	0.45
M ₄	7.32	7.38	7.35	7.29	7.34	0.280	0.270	0.260	0.253	0.266	0.46	0.49	0.52	0.54	0.50
Mean	7.46	7.43	7.41	7.36		0.282	0.275	0.268	0.265		0.41	0.42	0.44	0.46	
	M	F	M at F			M	F	M at F			M	F	M at F		
SE	0.140	0.15	0.308			0.008	0.006	0.011			0.01	0.001	0.017		
CD (p=0.05)	NS	NS	NS			NS	0.012	NS			0.029	0.018	NS		

3.2.2. SOIL ORGANIC CARBON

At post harvest stage, higher organic carbon content was recorded with application of poultry manure (0.50%) followed by vermicompost application (0.45%). This might be due to slow mineralization could lead to high organic carbon accumulation in soil. Invariably, all the organic treatments registered with the organic carbon content ranged from 0.44 to 0.50%. The findings are in agreement with [18].

The 125% RDF recorded lower organic carbon (0.49%) when compared to 100 % RDF and 75% RDF, respectively. The control (No manure - M₁) and (No fertilizer- F₁) registered with lower organic carbon content in this study. This might be due to application of inorganic NPK fertilizer fails to sequester SOC, only the application of organic manure alone has showed effective for sequestering soil organic carbon [19].

3.2.3. POST HARVEST SOIL AVAILABLE NUTRIENTS

The post harvest available soil nutrients was significantly influenced by organic manures and inorganic fertilizers during both the years of study. The treatments differed for the available nutrient status of the soil (Table 4).

Table 4. Influence of bulky organic nutrients and fertilizers on post-harvest available soil nutrients (Pooled data)

Treatments	N (Nitrogen Kg ha ⁻¹)					P (Phosphorus Kg ha ⁻¹)					K (Potassium Kg ha ⁻¹)				
	F ₁	F ₂	F ₃	F ₄	Mean	F ₁	F ₂	F ₃	F ₄	Mean	F ₁	F ₂	F ₃	F ₄	Mean
M ₁	70.6	80.7	78.7	72.4	75.6	7.7	9.0	8.9	8.2	8.5	74.7	85.5	83.7	79.9	80.9
M ₂	84.1	91.3	89.7	85.5	87.6	9.6	10.9	10.1	9.9	10.1	91.5	98.7	95.6	92.5	94.6
M ₃	83.6	100.5	97.6	94.6	94.1	9.3	12.5	12.0	11.0	11.2	90.6	107.9	104.7	100.8	101.0

M₄	80.7	109.5	104.3	102.7	99.3	9.0	15.5	13.5	15.5	13.4	85.5	118.6	115.5	110.7	107.5
Mean	79.8	95.5	92.5	88.8		8.9	12.0	11.1	11.1		85.6	102.6	99.9	95.9	
	M	F	M at F			M	F	M at F			M	F	M at F		
SE	2.38	1.85	3.69			0.30	0.21	0.43			2.58	1.98	3.97		
CD (p=0.05)	5.82	3.81	7.62			0.73	0.44	0.88			6.31	4.09	8.19		

Among the organic manures, application of poultry manure registered higher soil available N (99.28 kg ha⁻¹), soil available P (13.35 kg ha⁻¹) and soil available potassium (107.53 kg ha⁻¹) followed by application of vermicompost to the fodder maize crop (Table 3). Higher nutrient availability in the organic manures such as poultry manure might be due to continuous and slow release of nutrients from organic manure and increased biomass and accumulated soil organic matter. Similar findings were also reported by [20]. Application of farm yard manure for the fodder maize resulted in lower quantity of soil available nutrients followed by no application of organic manures to the fodder maize.

The soil available nutrients were lesser under the 75% recommended dose of fertilizer followed by no application of inorganic fertilizer. This might be due to lower amount of residual nutrient in low dose of inorganic fertilizer applied field. Inorganic fertilizers cause immediate release of nutrients, which will be utilized by the crop or might have lost the environment through leaching or identification process. Similar results were also noted by [21]. The soil available nutrients were higher with the application of 125% recommended dose of fertilizer followed by the application of 100% recommended dose of inorganic fertilizer.

The interaction between organic manures and inorganic fertilizers was found significant. The treatment combination of poultry manure along with 125% recommended dose of inorganic fertilizer to fodder maize recorded higher soil available nitrogen (109.48 kg ha⁻¹), phosphorus (15.46 kg ha⁻¹) and potassium (118.65 kg ha⁻¹) followed by application of poultry manure along with 100% recommended dose of inorganic fertilizer to fodder maize crop. This could be due to slow release of nutrients from organic manures and excess quantity of inorganic fertilizers was not fully utilized by the maize crop and leads to accumulation of nutrients in the soil after harvest of the crop. The least soil available nutrients were recorded under the control plots.

3.3 GREEN AND DRY FODDER YIELD OF MAIZE

Among organic manures, application of poultry manure (equivalent to 50 kg N or 12 t ha⁻¹) produced higher green fodder yield (21.17 t ha⁻¹ at 45 DAS and 28.22 t ha⁻¹ at 60 DAS). Maize dry fodder yield (3.54 t ha⁻¹ at 45 DAS and 4.66 t ha⁻¹) (Table 5) also recorded higher under the treatment of poultry manure application which is followed by application of Farm Yard Manure at 45 and 60 DAS. This could be ascribed to the higher nutrient composition in poultry manure (Table 1) coupled with slow and steady rate of nutrient release in to soil solution was responsible for better absorption of nutrients by fodder maize in different crop growth stages starting from vegetative to grain formation stage. This is in conformity with the findings of [22].

With regard to fertilizer levels, higher green fodder yield was recorded under the 75% RDF (19.84, 27.07 t ha⁻¹ at 45 and 60 DAS, respectively). Higher yield of maize dry fodder was registered (3.20 and 4.30 at 45 and 60 DAS, respectively) in 75% RDF followed by 100% RDF. The increase

levels of fertilizer (125% RDF) have not increased the green fodder and dry fodder yield. Primary nutrients are the important nutrients, which enhanced the plant growth parameters such as plant height, number of leaves, dry matter accumulation etc. and other yield parameters of fodder maize. Hence, lower dose of fertilizer will be enough to increase the plant growth and thus increase the green fodder yield. The findings were similar to the previous findings of [23].

The treatment combination of bulky organic nutrients and different fertilizers were significant. Green and dry fodder yield were recorded higher under the poultry manure + 75% RDF (31.8 t ha⁻¹ and 5.20 t ha⁻¹ at 60 DAS, respectively), when compared to this, the treatment of poultry manure + 100% RDF recorded low green and dry fodder yield. The reason for increased green and dry fodder yield might be due to slow and steady release of nutrients from poultry manure and effective utilization of applied nutrients by fodder maize. These combined nutrients enhanced the maize plant height and dry matter accumulation at different stages and thus increased green and dry fodder yield.

In this study, the applied poultry manure had higher amount of all nutrients and it improved the physical, chemical and biological properties of the soil. So, applied nutrients of 75% RDF were effectively utilized by the maize crop. These results were similar to the findings of [24] in maize crop.

Nitrogen nutrient can be supplied by many commercial fertilizers like urea and ammonium nitrate, which are commonly used by the farmers. In case of bulky organic nutrients, which also supply all essential nutrients to the crop growth and thus improves soil fertility which will finally increase the crop growth and yield in a sustainable way [25].

Table 5. Influence of bulky organic nutrients and fertilizers on green and dry fodder yield (t ha⁻¹) of fodder maize (Pooled data)

Treatments	Green fodder yield (t ha ⁻¹)										Dry fodder yield (t ha ⁻¹)									
	45 DAS					60 DAS					45 DAS					60 DAS				
	F ₁	F ₂	F ₃	F ₄	Me an	F ₁	F ₂	F ₃	F ₄	Me an	F ₁	F ₂	F ₃	F ₄	Me an	F ₁	F ₂	F ₃	F ₄	Me an
M ₁	16.1	17.3	17.3	16.3	16.8	23.3	24.4	24.3	23.6	23.9	1.9	2.6	2.5	2.2	2.3	3.1	3.7	3.6	3.4	3.4
M ₂	16.8	19.4	19.6	19.8	18.9	23.9	26.5	26.9	27.1	26.1	2.4	3.2	3.4	3.4	3.1	3.5	4.4	4.5	4.5	4.2
M ₃	18.5	17.8	18.2	18.7	18.3	25.2	24.9	25.2	25.8	25.3	2.5	2.8	3.0	3.0	2.8	3.6	3.9	4.1	4.2	4.0
M ₄	17.4	20.5	22.2	24.5	21.2	24.5	27.4	29.2	31.8	28.2	2.6	3.6	3.8	4.1	3.5	3.7	4.7	5.0	5.2	4.7
Mea n	17.2	18.8	19.3	19.8		24.2	25.8	26.4	27.1		2.36	3.07	3.16	3.2		3.5	4.2	4.3	4.3	
	M	F	M at F			M	F	M at F			M	F	M at F			M	F	M at F		
SE	0.51	0.39	0.78			0.70	0.55	1.09			0.08	0.06	0.12			0.11	0.08	0.17		
CD (p=0.05)	1.25	0.81	1.61			1.70	1.13	2.25			0.20	0.12	0.25			0.27	0.17	0.35		

4. CONCLUSION

The imbalanced use of inorganic fertilizers without organic fertilizers has led to the deterioration of soil health. Organic manure contains essential plant nutrients that can be reused for crop production. In the present study, use of organic manures, specifically poultry manure resulted in improve of soil physical and chemical properties viz., bulk density, water holding capacity, porosity and organic carbon. The highest organic carbon content was noticed with the application of poultry manure. The integrated nutrient combination of of poultry manure along with 125% recommended dose of inorganic fertilizer to fodder maize recorded higher soil available nutrients. Significantly higher green and dry fodder yield of fodder maize was recorded with application of poultry manure along with 75% recommended dose of fertilizer. Therefore, it is concluded that integrated nutrient management is crucial for sustainable crop production.

REFERENCES

1. Turan M and Yildirim E (Eds.). 2022. New Generation of Organic Fertilizers. Intech Open. doi: 10.5772/intechopen.95683.
2. Rekha SN and Prasad, RN. 2006. Pesticide residue in organic and conventional food-risk analysis. *Journal of Chemical and Health Safety*. 13: 12-19.
3. Singh TB. 2020. Role of Organic Fertilizers in Improving Soil Fertility. In: Naeem M, Ansari A, Gill S (eds) Contaminants in Agriculture. *Springer, Cham*.
4. Brewbaker J L. 2003. Corn production in the tropics: The Hawaii experience, *College of tropical agriculture and human Resources University of Hawaii at Manoa*.
5. Chaudhary DP, Jat SL, Kumar R, Kumar A and Kumar B. 2014. Fodder quality of maize its preservation In: Maize: *Nutrition Dynamics and Novel uses*. Pp: 153-160.
6. Piper CS. 1966. Soil and plant analysis. Hans pub., Bombay, pp: 368.
7. Jackson ML. 1973. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd, New Delhi.
8. Walkley A and Black CA. 1934. An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 40: 233-243.
9. Subbiah BV and Asija GL. 1956. A rapid procedure for estimation of available nitrogen in soils. *Current Science*, 25: 259-260.
10. Olsen SR, Cole CV, Watanabe FS and Dean AL. 1954. Estimation of available phosphorus in soils by extraction on with Sodium bicarbonate Circular no: 939, USDA.
11. Stanford S and English L. 1949. Use of flame photometer in rapid soil tests of K. *Canadian Journal of Agronomy*, 41: 446-447.
12. Panse VG and Sukhatme PV. 1985. Statistical methods for agricultural workers. Indian Council of Agricultural Research, New Delhi. pp: 100-174.
13. Agbede TM, Ojeniyi SO and Adeyemo AJ. 2008. Effect of poultry manure on soil physical and chemical properties, growth and grain yield of sorghum in southwest, Nigeria. *American-Eurasian Journal of Sustainable Agriculture*, 2(1): 72-77.
14. Agyenim SB, Zickermann J and Kornahrens M. 2006. Poultry manure effect on growth and yield of maize. *West African Journal of Applied Ecology*, 9: 1-11.
15. Jagadeesha N, Srinivasulu GB, Rathnakar M, Shet MR, Umesh, Gajanana Kustagi, Ravikumar B, Madhu L and Reddy VC. 2019. Effect of Organic Manures on Physical, Chemical and Biological Properties of Soil and Crop Yield in Fingermillet-Redgram Intercropping System *Int.J.Curr.Microbiol.App.Sci*. 8(5): 1378-1386.
16. Krishnakumar S, Saravanan A, Rajesh V and Mayil Samy P. 2007. Effect of organic farming on physical properties of rice grown soil. *J. Ecobiol*. 21(3): 263-267.
17. Rukmangada Reddy S, Reddy VC, Ramakrishna Parama VR. and Pampa Samanta. 2007. Effect of FYM, sewage sludge and urban compost on growth and yield of sweet sorghum (*Sorghum bicolor* L. Moench). *Journal of Soils and Crops*, 17(2): 211-216.

18. Dinesh Kumar. 2006. Integrated nutrient management for finger millet-soybean and soybean-finger millet cropping system. *Ph.D., Thesis*, UAS, Bangalore, Karnataka.
19. Hema RK, Bama S, Santhy P, Somasundaram E, Patil SG. Impact of different cropping and different nutrient management practices on soil carbon pools and soil carbon stock in verticustropept. *J Pharmacogn.andPhytochem.* 2019; 8(3):3424-3428.
20. Alagappan, S and Venkitaswamy R. 2019. Impacts of different sources of organic manures on soil physico-chemical properties, nutrient balance and yield of rice-green gram cropping sequence under organic farming. *International Journal of Agriculture & Environmental Science*, 6 (4). 65-85.
21. Singh RP, Yadav PK, Singh RK, Singh SN, Bisen, MK and J Singh. 2006. Effect of chemical fertilizer, FYM and biofertilizer on performance of rice and soil properties. *Crop Research*,32(3): 283-285.
22. Masthan Reddy BG, Patter PS and Kuchanur PH. 2005. Response of rice to poultry manure and graded levels of NPK under irrigated conditions. *Oryza*, 42(2): 109-111.
23. Rama Bharti, Gupta SK, Mainak Ghosh, Anshuman Kohli, Panda CK, Singh MK and Pathak SK. 2019. Effect of plant population and nitrogen levels on growth and yield attributes of fodder maize (*Zea mays L.*). *Multilogic in Science*, 9 (30): 215 - 218.
24. Mahmooda Buriro, Avinash Oad, Tahmina Nangraj, Allah WadhayoGandahi. 2014. Maize fodder yield and nitrogen uptake as influenced by farm yard manure and nitrogen rates. *European Academic Research*, 2(9):11624- 11637.
25. Rao SS and Shaktawat MS. 2002. Residual effect of organic manure, phosphorus and gypsum application in preceding groundnut (*Arachis hypogea*) on soil fertility and productivity of Indian mustard (*Brassica juncea*). *Indian Journal of Agronomy*, 47: 487-494.
26. Kanchikerimath M and Singh D. 2001. Soil organic matter and biological properties after 26 years of maize-wheat-cowpea cropping as affected by manure and fertilization in a Cambisol in semiarid region of India. *Agriculture, Ecosystems and Environment*, 86: 155–162.

UNDER PEER REVIEW