

**Antibiotics susceptibility profile of Non-Typhoidal *Salmonella* Species using Selected 3<sup>rd</sup> Generation Cephalosporins antibiotics against Humans and Poultry Faeces in Ido-Ekiti, Ekiti State.**

**ABSTRACT**

**INTRODUCTION:** Non-typhoidal *Salmonella* (NTS) isolates are among the most common bacterial pathogens associated with antimicrobial resistance (AMR) where animals are known to be the major reservoir of NTS despite improvements in hygiene and sanitation.

**AIM:** This study was conducted to investigate the prevalence of non-typhoidal *salmonella* species from humans and poultry faeces in Ido-Ekiti using selected 3<sup>rd</sup> generation cephalosporin antibiotics. **STUDY LOCATION AND DURATION:** The study was carried out at Federal Teaching Hospital Ido Ekiti for 3 months between July 2023 and September 2023.

**METHODOLOGY:** A total of 300 faecal samples were collected (100 from humans and 200 from poultry) by random sampling and analyzed by enriching on Selenite F broth and cultured on XLD agar using standard method. Faecal samples were identified using morphology, microscopic and biochemical characteristics. Antibiotic susceptibility test determined using disc diffusion method.

**RESULTS:** The prevalence of NTS in human and poultry sources were 9% and 11.5% respectively. Antibiotic susceptibility test revealed high resistance rates to 3<sup>rd</sup> generation Cephalosporins antibiotics among the poultry isolates, especially to Cefuroxime, Cefotaxime and ceftriaxone at 86.9%, 65.2% and 82.6% respectively while Ceftazidime at 88.9% only had a high resistance from human samples as compared to the poultry samples.

**CONCLUSION:** The study confirmed that poultries in Ido Ekiti are likely major sources of resistant NTS, and this poses a potential hazard to the public, especially those living close to the poultry and those rearing them.

**Keywords:** Non Typhoidal Salmonella (NTS), 3<sup>rd</sup> Generation Cephalosporins, Humans, Poultry birds.

**1. INTRODUCTION**

Nontyphoidal *Salmonella* (NTS) infections in Africa have been reported to have a mortality rate of 20-25% by 2020 [10]. Most non-typhoidal *Salmonella* (NTS) infections are caused by *Salmonella enterica* Typhimurium or *Salmonella enterica* Enteritidis. In 2019, there were 87,923 cases of NTS, the second most common bacterial gastrointestinal infection after campylobacteriosis [8]. Non Typhoidal *Salmonella* (NTS) is a symptomatic infection caused by bacteria of the genus *Salmonella* [20]. It is defined as an infectious or toxic disease caused by an agent that enters the body through food. Infection with nontyphoidal serotypes of *Salmonella* usually results in food poisoning. The elderly, the young, and others with weakened immune systems are more likely to develop NTS. It is usually spread by eating contaminated meat, eggs, water, or milk [19]. Other foods can spread disease if they come in contact with people. Several domestic animals such as cats, dogs and reptiles can carry and spread the infection. Diagnosis of NTS is made by stool examination or blood test [22]. The main source of this pathogen is poultry and cattle. Although most cases of salmonellosis are resolved on their own, antibiotic treatment may be necessary in severe cases, they may progress rapidly to systemic infection, particularly in infants, older adults, and individuals with weakened immune systems, requiring prompt clinical treatment [2]. The high prevalence of NTS infections in Nigeria is linked to various factors, including poor sanitation, inadequate access to clean water, and inadequate food handling and storage practices.

The problem of NTS infections is further complicated by the emergence of antimicrobial-resistant strains. Antibiotic resistant infections lead to longer, more severe illnesses and reduced treatment effectiveness [12]. NTS strains have been shown to develop resistance to antibiotics commonly used to treat infections. The spread of antimicrobial-resistant *Salmonella* is a major public health problem, as drug-resistant infections can cause high morbidity and mortality due to reduced disease duration and severity and reduced treatment efficacy [15,12]. Following Fajiladeet *al.* [7], antimicrobial drug resistance is prevalent among NTS species where isolates showed resistance to amoxicillin (100%), cotrimoxazole (100%), ampicillin (86%), chloramphenicol (100%) and ofloxacin (57%).

Cephalosporin C was discovered in the sea near a sewage outfall in Susiccu by the port of Cagliari in Sardinia by the Italian pharmacologist Giuseppe Brotzu in July 1945 [21]. Cephalosporin compounds were first isolated from a culture of *Acremonium strictum* from sewage in Sardinia in 1948 by the Italian scientist Giuseppe Brotzu [17]. Cephalosporins act similarly to beta-lactam antibiotics. They inhibit the synthesis of peptidoglycan in the cell wall of bacteria. Peptidoglycan is the bacterial exoskeleton that provides structural integrity and shape to cells and protects them from rupture [16]. These antibiotics are divided into several generations based on the period of drug development and their antimicrobial properties [16]. The difference lies in the molecular structure and has important therapeutic implications. The group of cephalosporins considered third generation is the most prescribed [18].

Third generation cephalosporins are drugs used to control and treat gram-negative and gram-positive bacteria. This drug belongs to the beta-lactam drug family, the most prescribed drug group. These cephalosporins are semi-synthetic analogues with different chemical substitutions in the C7 acylamide chain. This category includes ceftriaxone, cefdinir, cefixime, cefixime, cefpodoxime, ceftazidime, cefoprazone, ceftizoxime, and ceftibuten. It is a broad-spectrum antibacterial agent with activity against gram-negative and gram-positive bacteria and is more active against primary and secondary cephalosporin-resistant gram-negative bacteria. Interestingly, third generation cephalosporins are more resistant to beta-lactamase than first or second generation cephalosporins, especially *Salmonella*, *Klebsiella* and *Escherichia coli*. Several studies have reported 3rd generation cephalosporin resistance genes in non-typhoidal *Salmonella* isolated from humans and animals in Nigeria. A study conducted in Lagos, Nigeria by Iroha *et al.* [11] found that 38.4% of *Salmonella* typhoidal isolates were resistant to 3rd generation cephalosporins, some third generation cephalosporins include cefotaxime, ceftriaxone, and ceftazidime, as well as the oximino-monobactam aztreonam.

## **2. MATERIALS AND METHODS**

### **2.1 ETHICAL CONSIDERATION, QUESTIONNAIRE AND INFORMED CONSENT**

The ethical clearance (ERC/2022/06/06/794A) for this research was given by Federal Medical Centre (FMC) Ethical Committee after due processes had been followed. Administration of questionnaires was given to the subjects who consented to obtain their demographic data and other relevant information. The study was conducted between July 2022 and November 2022.

### **2.2 HUMAN SAMPLES**

A total of one hundred patients attending Federal Medical Centre, Ido Ekiti presenting with the study infection participated in this study and 200 poultry birds. One hundred faecal samples were collected and enriched in selenite F broth overnight and cultured on Xylose Lysine Deoxycholate (XLD) agar (Oxoid,uk) for 24hrs, NTS Colonies were then identified [4].

### **2.3 POULTRY SAMPLES**

A total of 200 faecal samples from poultry birds were collected with sterile swab sticks and transported within 1 hour of collection to the laboratory. The swab was pre-enriched in enriched in

selenite F broth overnight and cultured on Xylose Lysine Deoxycholate (XLD) agar (Oxoid,uk) for 24hrs ,NTS Colonies were then identified [4].

#### 2.4 IDENTIFICATION OF ISOLATES

The isolates were identified as non-typhoidal *Salmonella* with the following biochemical tests: Substrate Utilization test with Kligler Iron Agar, Urease, Citrate, Indole, Motility and Oxidase test using standard methods described by Cheesbrough [5].

#### 2.5 ANTIMICROBIAL SUSCEPTIBILITY TESTING

Antimicrobial susceptibility tests were performed using Mueller-Hinton agar by Disc diffusion method as described by CLSI [6]. The following antibiotics (Oxoid, Basingstoke, UK) were tested: Ofloxacin (OFX) (5µg), Pefloxacin (PEF) (5µg), Ciprofloxacin (CIP) (5µg), Levofloxacin (LEV) (5µg).

#### 2.6 STATISTICAL ANALYSIS

The data generated from this study was analysis using SPSS version 16.0 (SPSS Inc.Chicago IL).

### 3. RESULTS

The overall prevalence of non-typhoidal *Salmonella* isolated in human and poultry faecal samples was 32(10.7%) isolated from July 2022 to November 2022 (Table 1). Nine (9.0%) non-typhoidal *Salmonella* isolates from human faecal were gotten at Federal Medical Centre, Ido Ekiti (Table 1). Based on their age, 61-70 were most prevalent at 55.5% There were 3 males and 6 females, and based on their occupation, farmers had a high prevalence at 44.4% as compared to traders (Table 2). All the patients were showing presenting symptoms of NTS. However, the prevalence of NTS among the poultry birds selected at Ido Ekiti poultry locations showed that out of 200 faecal samples collected, 23(11.5%) non-typhoidal *Salmonella* were isolated (Table 1).

**Table 1: Overall prevalence of Non-typhoidal *Salmonella* (NTS) isolated from humans and poultry faecal.**

	Nontyphoidal <i>Salmonella</i> (n= 100)	No of Patients (n= 100)	No of Poultry (n=200)	Total (n=300)
Positive	9 (9.0%)	23 (11.5%)	32 (10.7%)	
Negative		91 (91.0%)	177 (88.5%)	268 (89.3%)
<b>Total</b>		<b>100 (100.0%)</b>	<b>200 (100.0%)</b>	<b>300 (100.0%)</b>

**Table 2: Prevalence of NTS in relation to age, sex and occupation**

	Positive(%)
<b>AGE</b>	
1-10	3 (33.3%)
11-20	1 ( 11.1%)
61-70	5 (55.5%)
<b>Total</b>	<b>9 (100%)</b>

<b>SEX</b>	
Male	3 (33.3%)
Female	6 (66.6%)
<b>Total</b>	<b>9 (100%)</b>

  

<b>OCCUPATION</b>	
Trader	2 (22.2%)
Farmer	4 (44.4%)
None	3 (33.3%)
<b>Total</b>	<b>9 (100%)</b>

The antimicrobial susceptibilities of the nontyphoidal *Salmonella* using four prescribed antibiotics were revealed in (Fig. 1). Of the 9 human NTS isolates, all 9 (100%) isolates showed resistance to Cefuroxime, 4 (44.4%) isolates showed resistance to Cefotaxime, 8 (88.9%) NTS were resistant to ceftazidime while 6 (66.6%) isolates showed resistance to Ceftriaxone. However, out of the 23 NTS isolated from poultry, 20 (86.9%) and 15 (65.2%) isolates were resistant to Cefuroxime and Cefotaxime respectively while 21 (91.3%) and 19 (82.6%) were resistant to ceftazidime and ceftriaxone respectively.

**Fig 1: Prevalence of antibiotic resistance of NTS isolates from human and poultry faecal samples**



#### 4. DISCUSSION

Non-typhoidal *Salmonella* (NTS) is a very important pathogenic organism that its prevalence shows a serious threat to man [3] and the isolation from poultry and clinical sources has not received much attention in Ekiti State, Nigeria. However, the aim of this study was to isolate NTS from poultry and human faecal samples in Ido Ekiti, and poultry is confirmed to remain one of the major routes of transmission of this organism. From the results obtained, Nine out of the one hundred gastroenteritis patients had NTS. Many studies have shown that Non-

typhoidal *Salmonella* (NTS) is among the very common pathogens causing bacterial bloodstream and gastroenteritis in adults and children in sub-Saharan Africa as reported by Akyala *et al.* [1]. Twenty-three out of two hundred poultry samples had NTS and the rate of isolation is higher compared to the isolation from humans. The prevalence of NTS from human faeces was 9.0% and poultry fecal sample had an 11.5% isolation rate, its prevalence among poultry faecal might be assumed as the main source of spread to humans. This agrees with Krawiec *et al.* [14] who reported a 64 (6.4%) prevalence rate of NTS among 100 samples collected from poultry and also agrees with Haeusler *et al.* [9] who reported that non-typhoidal *salmonellae* are isolated in 2–27% of stool culture-positive diarrhoeal illness. This is because Non Typhoidal *Salmonella* species are prevalent and they occur as pathogenic bacteria in the intestines of domestic, poultry, and wild animals.

In the demographic data of humans with NTS, based on their sex, there was no difference statistically between infections in males as compared to females suggesting that anybody can be infected either male or female, most especially when coming in contact with the bacteria. This agrees with an earlier work of Akyala *et al.* [1] who reported that sex is not a predisposing factor to NTS bacteremia in immunocompromised hosts. The age distribution showed a high prevalence in the age group 61-70 years at 55.5% which showed it was statistically significant to the study. This is the most immunocompromised age of the patients which could be as a result of weak immunity to various illness, and the prevalence of NTS tends to increase among these age groups and this could have predisposed them to high prevalence among the age group. Also, the study conducted by Kaushik and LeClercq [13] also showed a high prevalence of NTS among the age group 60 years and above. Based on their occupational status, the infection was more prevalent among farmers at 44.4%. This could be as a result of the farmers constant exposure to the poultry environment leading to an horizontal transmission from poultry to human and also the consumption of farmed food products that was grown with poultry faecal manure. This confirms that the spread of NTS is from poultry and livestock's which agrees with the study reported by WHO [23] that most of NTS are from poultry.

Antibiotic resistance of NTS isolates from sick patients and poultry faeces are shown in fig 1 suggests that antimicrobial resistance is widespread in both humans and poultry. High resistance rates to 3rd generation Cephalosporins antibiotics were seen among the poultry, especially to Cefuroxime, Cefotaxime and ceftriaxone while Ceftazidime only had an high resistance from human samples. This may be because poultry birds are usually given antibiotics which have created antibiotic resistance in humans. There have been reports indicating that the use of antimicrobials for growth promotion, prophylaxis, and treatment of food animals can increase the prevalence of resistance in human pathogens, particularly non-typhoidal *Salmonellosis*. Krawiec *et al.* [14] reported on the recent ban on the importation of poultry products into Nigeria that could have led to an increase in smuggling, which may be a likely reason for the emergence of resistance to 3rd generation cephalosporins.

## **CONCLUSION**

The results of the study helped to explain the antibiotic susceptibility prevalence of NTS in human and poultry faeces in Ado Ekiti. The study confirmed that poultries in Ido Ekiti are likely major sources of resistant NTS, and this poses a potential hazard to the public, especially those living close to the poultry and those rearing them. There is a need to study the molecular relatedness of the isolates from both humans and poultry to confirm the source of infection in sick patients.

## **AVAILABILITY OF DATA AND MATERIALS**

The authors declare that all the data supporting the findings are provided within the manuscript.

## REFERENCES

1. Akyala, S., Villanueva, M.A., Sison, F.B., Benigno, C.C., and Kreausukon, K. (2013). Distribution of bla CTX- m, bla SHV, bla TEM genes in extended- diaspason  $\beta$ -lactamase- producing *E. coli* isolates from broiler farms in the Philippines. *BMC Veterinary Research*. 15( 1), 1 – 8.
1. Antunes, P., Campos, J., and Peixe, L. (2016). *Salmonellosis* and its clinical contagious conditions. 41 Suppl 2( S2) S113- 9.
2. Anyanwu, B., Abdulfatai, K., Wartu, J.R., Mzungu, I., Muhammad, H.I.D. and Abdulsalam, A.O. (2010). Antibiotics vulnerability patterns and characterization of clinical *Salmonella* serotypes in Katsina State, Nigeria. *African Journal of Microbiology Res* 8915 – 921.
3. Benneth, T.J. and McCartney, J.E. (1997) Practical Medical Microbiology. In, 13th Edition, Vol. 12, Churchill Livingstone, London, 910.
4. Cheesbrough, M. (2006). District Laboratory Practice in Tropical Countries. Part 2, Alternate edition. *Cambridge University press* PP.182- 183.
5. Clinical Laboratory Standard's Institute (CLSI). (2020) Performance standards for antimicrobial susceptibility Testing. 26th ed. CLSI, supplement M100S. Wayne, PA *Clinical and Laboratory Standards Institute*.
6. Fajilade, O., Oluwafemi, A., Taiwo, L. (2022). Prevalence of multidrug- resistant *Salmonella spp.* in original food products vended in Ado- Ekiti, South Western Nigeria. *Microbes and contagious conditions*.3(2) 360- 365.
7. Food Drug Administration. (2019)" FDA issues peanut safety guidelines for foodmakers". Reuters. 2019-03-10.
8. Haeusler, S., Epple, H.J., Feurle, G.E., Kern, W.V. and Lynen, J.P. (2013) S2k- guideline gastrointestinal contagious conditions and Whipple's complaint. *Gastroenterology*, 53418 – 459.
9. Hald, T. (2021). Advances in microbial food safety Pathogen update *Salmonella*. *Elsevier Inc*. p.2.2. ISBN 9780128089606.
10. Iroha, I.U. (2017)." Extended- Spectrum  $\beta$ - Lactamase (ESBL)- producing non-typhoidal *Salmonella* (NTS) from humans and poultry in Lagos, Nigeria." *Journal of health, population, and nutrition*36.1 1- 10.
11. Jajere, S.M. (2019). A review of *Salmonella enterica* with particular focus on the pathogenicity and acidity factors, host particularity and antimicrobial resistance including multidrug resistance. *Veterinary World* 12, 504 – 521.
12. Kaushik, S., and Leclercq, R. (2014). Role of a qnr- suchlike gene in the natural resistance of *Enterococcus faecalis* to fluoroquinolones. *Antimicrobial. Agents of Chemotherapy*.513254- 3258.
13. Krawlec, C.A., Zarazaga, M., Sallem, R., Jouini, A., Slama, K. and Torres, C. (2015). Antibiotic resistance in *E coli* in animals husbandry. *Letter of Applied. Microbiology*.64( 5), 318 – 334).
14. Nair, D., Venkitanarayanan, K. and Kollanoor, J.A. (2018). Antibiotic resistant *Salmonella* in the food supply and the implicit part of antibiotic alternatives for control. *Foods* 7:167.
15. Pegler, I., Scott, V., Healy, B. and Brendan, H. (2017)." In cases antipathetic to penicillin, consider alternate and third generation cephalosporins for life hanging infections". *The Biomedical Journal*. 335(7627) 991.
16. Podolsky, T., Kaku, M., Onodera, S. and Yamachika, K. (1998). *Vibrio parahaemolyticus* chromosomal qnr homologue VPA0095 demonstration by metamorphosis with a shifted

- gene of its eventuality to reduce quinolone vulnerability in *Escherichia coli*. *Antimicrobial Agents of Chemotherapy*. 214 Avian Pathology 4- 2145.
17. Prince, A. (2022)" Cephalosporins and vancomycin". Columbia University. 15 October 2022.
  18. Santos, R.L., Shuping, Z., Renee, M., Tsolis, R.A., Kingsley, L. Gary, A. and Andreas, J.B. (2021)." Beast models of *Salmonella infections enteritis* versus typhoid fever". *Microbes and Infection*. 3(14 – 15) 1335 – 1344.
  19. Smith, J.L. and Bayles, D. (2017)." Postinfectious perverse bowel syndrome;a long- term consequence of bacterial gastroenteritis". *Journal of Food Protection*. 70( 7) 1762 – 9.
  20. Tilli, T. and Lois, R. (2000). Post Penicillin Antibiotics From acceptance to resistance? Welcome substantiations to Contemporary Medicine. History of ultramodern Biomedicine Research Group. ISBN 978-1-84129-012-6.
  21. WHO. (2021)." *Salmonella*(non-typhoidal)". World Health Organization. December 2021.
  22. World Health Organization. (2020). Non-typhoidal *Salmonella*.