

# HAEMATO-BIOCHEMICAL AND HORMONAL PROFILE ALTERATIONS IN POSTPARTUM TRUE ANESTRUS MEHSANA BUFFALOES TREATED WITH INDIGENOUS HERBAL HEAT INDUCER

## ABSTRACT

The present study was designed to explore the effects of a new Herbal heat inducer in Mehsana buffaloes with postpartum true anestrus and consecutive changes in Blood haematological, Biochemical and Hormonal profile. All the buffaloes were orally supplemented with Herbal heat inducer mixture on (Jantana powder-10gm)<sup>1<sup>st</sup></sup>, (Metrali powder-20gm)<sup>2<sup>nd</sup></sup> and (Hitali-powder-500mg)<sup>5<sup>th</sup></sup> day for augmentation of oestrus. Out of total treated animals 10 animals expressed the signs of estrus within 30 days. Post-treatment analysis of the overall blood packed cell volume percent in conceiving buffaloes had a significantly ( $p \leq 0.05$ ) elevated. Overall blood plasma total cholesterol concentration (mg/dl) was found to be a significantly ( $p \leq 0.001$ ) elevated in all treated and both conceiving and non-conceiving buffaloes, respectively. Overall blood plasma progesterone concentration (ng/ml) was found to be a significantly ( $p \leq 0.0001$ ) and ( $p \leq 0.001$ ) elevated in all treated and both conceived and non-conceived buffaloes, which indicate the positive favorable effect of all the drugs on postpartum true anestrus condition. However, Overall blood plasma tri-iodothyronine concentration (ng/ml) was found to be a significantly ( $p \leq 0.001$ ) and ( $p \leq 0.01$ ) elevated in all treated and all conceived buffaloes. The findings of present study revealed that strategic use of Indigenous herbal heat inducer in feed mixture may resolve the postpartum true anestrus problems in buffaloes.

**Keywords:** Mehsana Buffaloes, Postpartum true anestrus, Herbal heat inducer, Haematological profile, Blood Plasma profile

## INTRODUCTION

The Buffaloes in tropical and subtropical countries are used as a source of milk, meat and draught purpose. Postpartum True anestrus (PPTAE) is one of the main reproductive constraints in lactating buffaloes, and is much higher in conventionally managed herds than in organized farms. Reproductive efficiency of the animal is the primary determinant for sound and economic animal production system. A Large number of interacting factors viz. endocrine disturbances, poor nutrition, seasonal and systemic diseases might be attributed to the cause of primary infertility. Also the anestrus is multi-faceted problem, but inadequate nutrition, particularly dietary insufficiency of mineral, like calcium, phosphorus, copper, zinc and manganese, greatly contribute to anestrus (Kumaret *al.*, 2014), as these minerals play intermediate roles in the action of hormones and enzymes at cellular level, as reproductive cycle in animals are regulated by endocrine-neuroendocrine interactions between hypothalamic, gonadotropic, gonadal and other hormones (Terzano *et al.*, 2012). There are also certain biological and metabolic constituents that directly reflect the nutritional status and influence the reproductive performance of PPTAE either by acting as precursor of hormone synthesis or by stimulating response of target tissues. Disturbances in haematological, hormonal and biochemical milieu due to deficiency of trace minerals also cause anestrus where their estimation in serum samples might be a potential aid in

characterizing this problem in post parturient buffaloes. Several Indigenous Herbal inducer are being used since dates back to augment the reproductive performances as traditional medicine. There is a paucity of information regarding the blood haematological, biochemical and hormonal profile in estrus and anestrus phase. Based upon the above facts, the present study was designed to evaluate the influence of Herbal heat inducer mixture on reproductive performance of postpartum true anestrus Mehsana buffaloes.

## **MATERIALS AND METHODS**

The present investigation was carried out under field conditions of Banaskantha district north western part of the state of Gujarat and collaboration with the Banaskantha district cooperative Milk producers unions limited (Banas Dairy) to study the changes in haemato-biochemical and hormonal profile in postpartum true anestrus Mehsana buffaloes treated with Indigenous Herbal heat inducer.

The Mehsana buffaloes with good physical condition were gynaecologically examined. The buffaloes having smooth and inactive ovaries without palpable follicles or corpora lutea on either of ovaries and without pathological condition in the reproductive tract were randomly selected for the study. Repeated rectal examinations were made at 10-12 days interval until a definite conclusion could be drawn for cause (s) of anestrus. A total number of 10 buffaloes affected with postpartum true anestrus were selected for the study. All the buffaloes were orally supplemented with different Herbal heat inducer mixture Jantana powder-10g on 1<sup>st</sup> day, Metrali powder-20g on 2<sup>nd</sup> day and Hitali powder-500mg on 5<sup>th</sup> day of Fertikit, Mycon Pharma. All the buffaloes were observed for occurrence of estrus and were bred with fertile bull semen. Buffaloes which did not show the signs of estrus following the treatment were examined per rectally for pregnancy diagnosis on 60<sup>th</sup> day post-insemination to confirm the pregnancy and the buffaloes which did not conceive at first insemination were re-inseminated at subsequent estrus cycles.

The blood samples from different groups were collected on day 0 before treatment and on 7<sup>th</sup> day after treatment. A 10 ml of blood was collected by jugular vein puncture into a heparinized polystyrene tube, and the sample was maintained at 4°C and transported to laboratory within 1-2 hrs of collection. Blood haemoglobin (Hb) and packed cell volume (PCV) was estimated as per standard protocol (Benjamin, 1985). Plasma was separated by centrifugation at 3000 rpm for 15 minute and was stored in deep freeze (-20 °C) until analyzed. Plasma progesterone, tri-iodothyronine and thyroxine were estimated by using commercially available ELISA kits (Sigma Diagnostics Pvt. Limited, USA) and plasma total protein and total cholesterol were estimated by using standard procedures using commercial available diagnostic kits (LiquiCHEK™, AGAPPE Diagnostics Pvt. Limited, Kerala, India) in Biochemistry Analyzer RX-50V (Micro Lab, India). The data collected were suitably tabulated and analyzed by standard statistical Graphpad prism® Trial version 4 software. The data obtained pertaining to different groups were analyzed statistically using Two-way ANOVA. While the test of significance between and within the treatment groups were calculated by multiple comparison with Bonferroni test (Bonferroni, 1960) with 95% confidence interval.

## **RESULTS AND DISCUSSION**

### **Blood Haematological Profile**

Overall haemoglobin concentration (gm/dl) was found to be  $13.64 \pm 0.30$  (11.7-16.8) in treatment group buffaloes. Statistical analysis revealed a non-significant difference within various periods in the treatment group buffaloes with an increase at day 7 (Table 1). Statistical analysis also revealed a non-significant difference between and within the conceived and non-conceived buffaloes (Table 2). Haemoglobin levels (gm/dl) at various periods observed in the present study is in agreement with the Das *et al.*, (2016) in buffaloes. In the present study, blood haemoglobin profile revealed a non-significant difference within the group, agreed well with the Ray *et al.*, (2016) who have reported that adequate haemoglobin level is required to transport enough oxygen and nutrients to the vital organs including the ovarian tissues to be prepared for sending the signal to hypothalamus, which is in agreement with an increase in haemoglobin concentration after treatment in all groups.

**Table1. Blood Haemoglobin concentration (gm/dl) in postpartum true anestrous buffaloes group (Mean $\pm$ SEm).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Group	12.98 $\pm$ 0.36(11.7-14.9)	14.29 $\pm$ 0.39(12.8-16.8)	13.64 $\pm$ 0.30(11.7-16.8)

- Figures in parentheses indicate range.

**Table2. Blood Haemoglobin concentration (gm/dl) in all conceived and non-conceived true anestrous buffaloes (Mean $\pm$ SEm).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Conceived	12.39 $\pm$ 0.53(8.1- 16.2)	13.46 $\pm$ 0.48(9.2- 16.8)	12.93 $\pm$ 0.37(8.1-16.8)
Non-conceived	13.08 $\pm$ 0.73(9.2- 16.6)	14.26 $\pm$ 0.51(12.2- 16.8)	13.68 $\pm$ 0.46(9.2-16.8)

- Figures in parentheses indicate range.

Overall per cent packed cell volume was found to be  $34.27 \pm 0.83$  (28.9-42.9) in treatment group buffaloes. Statistical analysis revealed a non-significant difference within various periods in the treatment group (Table.3). The overall packed cell volume per cent observed at different periods in all the conceived and non-conceived buffaloes show non-significant difference are presented in table.4. Statistical analysis revealed a significant ( $P \leq 0.01$ ) difference within the conceived buffaloes with higher PCV per cent at day 7<sup>th</sup> ( $36.93 \pm 0.94$ ) as compared to that of day 0 ( $33.51 \pm 0.88$ ) (Table.4), similarly Das *et al.*, (2016) has reported in Mehsana buffaloes. An increase in PCV value was observed in all the groups parallel with an increase in haemoglobin concentration after treatment which is in agreement with Ray *et al.*, (2016) who have also reported in respective groups of postpartum anoestrus buffaloes.

**Table3. Blood Packed cell volume (PCV) per cent in postpartum true anestrous buffaloes group (Mean $\pm$ SEm).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Group	32.32 $\pm$ 0.76(28.9-36.8)	36.21 $\pm$ 1.22(23.9-42.9)	34.27 $\pm$ 0.83(28.9-42.9)

- Figures in parentheses indicate range.

**Table4. Blood Packed cell volume (PCV) per cent in all conceived and non-conceived true anestrous buffaloes (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Conceived	33.51±0.88 <sup>p</sup> (28.4- 41.1)	36.93±0.94 <sup>q</sup> (28.7- 42.9)	35.23±0.69(28.4-42.4)
Non-conceived	33.88±1.41(28.6- 40.9)	36.81±1.09(32.9- 42.6)	35.35±0.94(28.6-42.6)

- Figures in parentheses indicate range.
- Values bearing different superscripts within column differed significantly ( $P \leq 0.01$ ).

### Blood Plasma Biochemical Profile

Overall blood plasma total protein concentration (gm/dl) was found to be 8.81±0.34(5.99-11.22) in treatment group buffaloes. Statistical analysis revealed a non-significant difference within the treatment group (Table 5). The plasma total protein concentration (gm/dl) observed at various periods was non-significantly higher in conceiving as compared to non-conceiving buffaloes (Table 6). Which is an agreement with the study of Singh *et al.*, (2012) who have reported that the plasma total protein levels at different periods which gets the support of is increasing post-treatment in bovine using herbal estrus inducer therapy because its might due to herbal medicines that contain some amounts of proteins and concluded that the compound of herbal drugs influenced the biochemical profile of blood under normal range which ultimately affects the estrus response in bovine. However, the level was non-significantly higher at 7<sup>th</sup> day in all the conceiving buffaloes than non-conceived buffaloes under experiment.

**Table5. Blood plasma total protein concentration (gm/dl) in postpartum true anestrous buffaloes group (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Group	8.25±0.40(5.99-10.2)	9.36±0.49(6.2-11.22)	8.81±0.34(5.99-11.22)

- Figures in parentheses indicate range

**Table6. Blood plasma total protein concentration (gm/dl) in all conceived and non-conceived true anestrous buffaloes (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Conceived	8.49±0.29(5.99-10.93)	9.21±0.35(5.99-11.22)	8.85±0.23(5.99-11.22)
Non-conceived	7.69±0.38(6.44-9.98)	8.73±0.39(7.5-10.88)	8.44±0.27(6.78-10.88)

- Figures in parentheses indicate range.

Overall blood plasma total cholesterol concentration (mg/dl) was found to be 118.08±4.15 (77.6-145.2) in treatment group buffaloes. Statistical analysis revealed an increase in total cholesterol at day 7 post treatment with a significantly ( $P \leq 0.001$ ) higher value only in treatment group buffaloes (Table 7). However, significantly ( $P \leq 0.001$ ) higher cholesterol level was also observed at 7<sup>th</sup> day in all the conceiving and non-conceiving buffaloes under experiment (Table

8). Plasma total cholesterol concentration (mg/dl) level observed in the present study is in agreement with that of the 111.55±10.12 (mg/dl) level at expressed heat and 88.91±10.77 (mg/dl) level in dairy bovine which not expressing heat after treatment for postpartum anoestrus using herbal heat inducer (Ray *et al.*, 2016) and 160.76±0.52 (mg/dl) level at estrus and 130.98±0.31 (mg/dl) level in regular cyclic Murrah buffaloes (Balamurugan *et al.*, 2015). Significantly higher cholesterol level in treatment group animals might be an indication of positive action of the drug in reducing the service period since higher plasma cholesterol level facilitates early expression of oestrus in dairy bovine, as lipids are the precursors of gonadal steroid hormones (Westwood *et al.*, 2002).

**Table 7. Plasma total cholesterol concentration (mg/dl) in postpartum true anoestrous buffaloes group (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Group	107.44±5.82 <sup>P</sup> (77.6-130.1)	128.72±5.17 <sup>q</sup> (99.7-145.2)	118.08±4.15(77.6-145.2)

- Figures in parentheses indicate range.
- Values bearing different superscripts within column differed significantly (P≤0.001).

**Table 8. Plasma total cholesterol concentration (mg/dl) in all conceived and non-conceived true anoestrous buffaloes (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Conceived	101.45±4.27 <sup>P</sup> (72.8-132.7)	121.26± 4.47 <sup>q</sup> (74.5-148.9)	111.35±3.42(72.8-142.2)
Non-conceived	106.28±6.45 <sup>P</sup> (77.6-133.5)	127.99±4.72 <sup>q</sup> (97.7-142.2)	117.13±4.68(77.6-142.2)

- Figures in parentheses indicate range.
- Values bearing different superscripts within column differed significantly (P≤0.001).

### Blood Plasma Hormonal Profile

The overall blood plasma progesterone concentration (ng/ml) was found to be 1.37±0.11(0.80-1.99) in treatment group buffaloes. Statistical analysis revealed a significant (P≤0.0001) difference within in the treatment group buffaloes (Table 9). Statistical analysis also revealed a significant (P≤0.001) difference within the group of all conceived and non-conceived buffaloes (Table 10). However, the difference was non-significant between the groups in conceived as well as non-conceived buffaloes. A significant (P≤0.001) increase in P<sub>4</sub> at day 7 post treatment, in conceiving as well as non-conceiving buffaloes indicate the favorable effect of all the drugs on anoestrus condition. The treatment used in the present study have shown a significant increase in P<sub>4</sub> levels in the manner of normal cyclic animals and also increased conception rate in anoestrus buffaloes under field condition in particular with the use of Herbal treatment.

**Table 9. Plasma progesterone concentration (ng/ml) in postpartum true anoestrous buffaloes group (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Group	0.97±0.04 <sup>P</sup> (0.80-1.18)	1.74±0.13 <sup>q</sup> (0.93-1.99)	1.37±0.11(0.80-1.99)

- Figures in the parenthesis indicate range
- Values bearing different superscripts within column differ significantly (P≤0.0001).

**Table10. Plasma progesterone concentration (ng/ml)in all conceived and non-conceived true anestrus buffaloes (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Conceived	0.99±0.06 <sup>P</sup> (0.18- 1.50)	1.74± 0.14 <sup>q</sup> (0.93-2.98)	1.36±0.09(0.18-2.98)
Non-conceived	0.91±0.04 <sup>P</sup> (0.82-1.18)	1.72± 0.10 <sup>q</sup> (0.99-1.98)	1.32±0.11(0.82-1.98)

- Figures in parentheses indicate range.
- Values bearing different superscripts within column differ significantly (P≤0.001).

Overall blood plasma tri-iodothyronine (T<sub>3</sub>) concentration (ng/ml) was found to be 5.92±0.26 (4.21-9.19) in treatment group buffaloes. Statistical analysis revealed that the T<sub>3</sub> levels were found to be significantly (P≤0.001) increase on day 7<sup>th</sup>(6.58±0.38 ng/ml) after treatment as compared to (5.27±0.24 ng/ml) on day 0 (Table 11).

Statistical analysis revealed a non-significant difference for T<sub>3</sub> levels between the conceived and non-conceived buffaloes, but a significantly (P≤0.01) higher overall T<sub>3</sub> level was observed at day 7 in all the conceiving buffaloes (Table.12).The mean plasma tri-iodothyronine concentration (ng/ml) observed in the buffaloes under experiment is in agreement with that of the postpartum Egyptian buffaloes (Ashmawy *et al.*, 2015). In the present study, the mean plasma tri-iodothyronine concentration (ng/ml) profile revealed a significant (P≤0.001) rise in the herbal group after treatment which is in agreement with the study of Dutt *et al.*, (2011) in postpartum anoestrus buffaloes using herbal extract therapy.

**Table11. Plasma tri-iodothyronine concentration(ng/ml)in postpartum true anoestrusbuffaloes group (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Group	5.27±0.24 <sup>P</sup> (3.86-6.55)	6.58±0.38 <sup>q</sup> (5.29-9.19)	5.92±0.26(4.21-9.19)

- Figures in parentheses indicate range.
- Values bearing different superscripts within column differed significantly (P≤0.001).

**Table12. Plasma tri-iodothyronine concentration (ng/ml) in all conceived and non-conceived true anestrus buffaloes (Mean±SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
Conceived	5.45 ± 0.19 <sup>P</sup> (4.10- 7.11)	6.51 ± 0.19 <sup>q</sup> (5.36- 8.12)	6.08±0.15(3.86-8.12)
Non-	5.81±0.30(4.62- 7.52)	6.28±0.41(4.79- 9.19)	6.01±0.14(4.10-9.19)

<b>conceived</b>			
------------------	--	--	--

- Figures in parentheses indicate range.
- Values bearing different superscripts within column differed significantly ( $P \leq 0.01$ ).

Overall blood plasma thyroxine ( $T_4$ ) concentration (ng/ml) was found to be  $53.83 \pm 4.59$  (28.58-101.37) in treatment group buffaloes. Statistical analysis revealed a non-significant difference within treatment group buffaloes (Table.13). It was found to increase non-significantly at day 7 in treatment group.

Statistical analysis revealed a non-significant difference within the conceived and non-conceived buffaloes (Table.14). The mean plasma thyroxine concentration (ng/ml) observed in the buffaloes under experiment is in agreement with the study of Kumar *et al.*, (2010) and Raja Kumar *et al.*, (2010) in cows at normal cyclic condition and postpartum anoestrus. Whereas, lower plasma thyroxine concentration (ng/ml) in comparison to present findings have been reported in normal cyclic and postpartum true anoestrus acyclic Nili-Ravi buffaloes by Muhammad *et al.*, (2016). In the present study, the profile of plasma thyroxine concentration (ng/ml) revealed the non-significant difference between and within the all conceived and non-conceived buffaloes but it was found to increase non-significantly at day 7 post-treatment in all conceived and non-conceived buffaloes indicating the response to herbal heat inducers.

**Table 13. Plasma thyroxine concentration (ng/ml) in postpartum true anoestrus buffaloes group (Mean  $\pm$  SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
<b>Group</b>	50.10 $\pm$ 5.60 (28.58-80.97)	57.54 $\pm$ 7.37 (34.84-101.37)	53.83 $\pm$ 4.59 (28.58-101.37)

- Figures in parentheses indicate range.

**Table 14. Plasma thyroxine concentration (ng/ml) in all conceived and non-conceived true anoestrus buffaloes (Mean  $\pm$  SEM).**

Period	P-1 (0 day)	P-2 (7 <sup>th</sup> day)	Overall
<b>Conceived</b>	53.04 $\pm$ 3.49 (28.58-81.70)	53.87 $\pm$ 3.42 (28.78-95.22)	53.46 $\pm$ 2.42 (28.58-95.22)
<b>Non-conceived</b>	50.67 $\pm$ 3.55 (32.80-64.45)	68.19 $\pm$ 8.98 (28.65-101.37)	59.44 $\pm$ 5.14 (28.65-101.37)

- Figures in parentheses indicate range.

### Fertility Response

The per cent estrus induction was 80.00 with days at 15.5 (8-30 days) and intensity of estrus in terms of intense: moderate: mild were 25.00: 37.50: 37.50 in treatment group buffaloes, The conception rates were 70.00 per cent in treatment group buffaloes with lesser no of services per conception (1.57). The per cent buffaloes conceived with first two services were 85.72 in treatment group. Average days at fertile estrus post treatment was observed to be 29.57 (7/10) in conceived buffaloes of the treatment group. The major symptoms noticed were vaginal mucus discharge, edema, erythematic and wetness of vulva followed by excitement, occasionally bellowing and rarely off feed symptoms. Per cent conception rate following herbal mixture therapy in the present study agreed well with findings of Deshpande *et al.*, (2000); Hadiya *et al.*, (2015) and Sahatpure *et al.*, (2016) in anoestrus buffaloes.

The herbal treatment group buffaloes have shown a better overall breeding performance in terms of fertile days post treatment and a number of services required per conception. This effect of herbal drugs may be attributed to the collective beneficial effect of various ingredients as reported with their known effect in particular Vidang (Anthelmintic: useful against tapeworms, *Dama and Kirdak., 2002*), Palashbeej (recovers sexual dysfunction, *Huxley et al., 1992*), Sonamukhi (heat inducer, anthelmintic and strengthen immunity), Gokharu (to treat utero-genital tract infections and sexual dysfunction, *Samani et al.,2016*), *Gorakhmundi* (anthelmintic and immunomodulatory and anti-inflammatory), Shatawari (antiinflammatory, nourishing the ovum and cures infertility,increases libido, *Alok et al.,2013*), Ashwagandha (useful to cure female and male sexual disorders), Kalonji (anti-inflammatory, analgesic, cleanser of uterus, anthelmintic, *Liu et al. (2016)*),Balshepa (increases the fertility rate, *Santos et al., 2002*), Gajar beej (useful in oligomenorrhea, dysmenorrhea and irregular estrus cycle, *Afzal, et al., 2013*), Shivlingi (excellent anti-inflammatory, anti-infertility and sterility treatment infemale, provides nourishment to the female organs and support normal functioning, *Verma et al., 1993*) which seem to be implicated in better health status and reproductive performance of the buffaloes of Herbal treatment group.

## CONCLUSION

The haemoglobin concentration (gm/dl) and PCV per cent increased non-significantly at day 7 in treatment group. Whereas, per cent packed cell volume increase significantly at day 7 in conceived group of buffaloes. Total protein concentration levels was non-significantly higher in buffaloes of conceived as compared to non-conceived group. Overall blood plasma total cholesterol concentration (mg/dl) found to be significantly ( $P \leq 0.001$ ) increase at day 7 post treatment in treatment group buffaloes, However, significantly ( $P \leq 0.001$ ) higher cholesterol level was also observed at 7<sup>th</sup> day in all the conceiving and non-conceiving buffaloes under experiment. A significant rise in plasma progesterone concentration (ng/ml) at 7th-day post-treatment indicates a positive effect of all the herbal heat inducers on cyclicity. A significant increase in tri-iodothyronine and total cholesterol was noticed using herbal heat inducer at day 7 post treatment. The conceived buffaloes had significantly higher tri-iodothyronine levels at day 7 post treatment. Average days at fertile estrus occurred earlier in the herbal treatment group with a lesser number of services per conception. Conclusively, orally supplemented Herbal heat inducer can be used to induce the estrous in postpartum true anestrus buffaloes.

## REFERENCES

- Afzal, M.; Kazmi, I. and Kaur, R. (2013). Comparison of protective and curative potential of daucus carota root extract on renal ischemia reperfusion injury in rats. *Journal of Pharmaceutical Biology.* **51** (7): 856-62.
- Alok, S.; Jain, S. K.; Vermad A.; Kumar, M.; Mahor, A. and Sabharwal, M. (2013). Plant profile, phytochemistry and pharmacology of *Asparagus racemosus* (Shatawari): *Asian Pacific Journal of Tropical Disease.* **3** (3): 242–251.

- Ashmawy, N. A. (2015). Changes in peripheral plasma hormone concentrations and metabolites during the last trimester of pregnancy and around Parturition in the Egyptian buffalo and baladi cows. *International Journal of Advanced Research*. **3** (11): 1377-1390.
- Balamurugan, T. C.; Krupakaran, P. R.; Pandiyan, G. D. V.; Arunkumar, S. and Perumal, P. (2015). Comparative studies on serum biochemical profiling at various stages of reproduction in Murrah buffaloes. *Journal of Cell and Tissue Research*. **15** (1): 4823-4826.
- Benjamin, M. M. (1985). Outline of veterinary clinical pathology. 3rd eds kalyani publication, new delhi, india, pp 233-254.
- Bonferroni, C. E. 1960. Biostatistical analysis of variance (Two way-ANOVA) experiment. Bonferroni Test <http://www.investopedia.com/terms/b/bonferroni-test.asp#ixzz4jZXuzfJw>.
- Dama, L. B. and Kirdak, R. V. (2002). Effect of Vidhang seed extract against *Ascaridia galli* in naturally infected fowls (*Gallus domesticus*)". *Journal of Parasitic Disease*. **26**: 48-49.
- Das, H.; Lateef, A.; Panchasara, H. H. and Ali, A. M. (2016). Haemato-biochemical alterations during different stages of lactation in Mehshani buffaloes. *Buffalo bulletin*. **35** (3): 307-312.
- Deshpande, R. S.; Dhoble, R. L. and Sawale, H. (2000). Efficacy of indigenous drugs in treatment of postpartum anestrus in buffaloes. *Indian Journal of Animal Reproduction*. **21** (2): 115-116.
- Dutt, R.; Mehrotra, S.; Shanker, U. and Singh, G. (2011). Effect of murrayakoenigii and aeglemaramelos feeding on anestrus buffaloes. *Indian Journal of Animal Reproduction*. **32** (1): 1-4.
- Hadiya, K. K.; Ravikanth, K.; Adarsh, and Reothia, A. (2015). Efficacy of Exapar, Janova and Mintrus (EJM) in the treatment of various reproductive disorders in Jaffrabadi buffaloes. *International Journal of Advanced Research*. **3** (9): 1213- 1216.
- Kumar, R.; Gupta, M.; Rose, M. K.; Ghosh, M. and Singh, I. (2014). Hematological profile in postpartum anoestrous and normal cyclic Murrah buffaloes. *Journal of cell and Tissue Research*. **14** (2): 4241-4243.
- Kumar, S.; Saxena, A. and Ramsagar, (2010). Comparative studies on metabolic profile of anestrus and normal cyclic Murrah buffaloes. *Buffallo Bulletin*. **29** (1): 7-11.
- Liu, X.; Sahebkar, A. and Soranna, D. (2016). A systematic review and meta-analysis of randomized controlled trials investigating the effects of supplementation with *Nigella sativa* (black seed) on blood pressure. *Journal of Hypertension*. **34** (11): 2127-35.

- Muhammad, U. G.; Ahmad, I.; Ahmad, N.; Ijaz, N. and Mehfooz, A. (2016). Hematology, serum total cholesterol and thyroid hormone concentrations in cyclic and acyclic Nili-ravi buffaloes. *Pakistan Veterinary Journal*. **37** (1): 31-34.
- Raja Kuamr, P.; Lakshmana swamy, P.; Aswani Kumar, K. and Rao, K. S. (2010). Studies on the thyroid profile ( $T_3$ ,  $T_4$  and TSH) in estrous, cyclic and anestrous buffaloes. *Buffalo Bulletin*. **29** (2): 115-128.
- Ray, K.; Biswas, P.; Banerjee, U; Basu, S. and Sarkar, B. (2016). Supplementation of Herbal oestrus inducer and Mineral mixture combinations on haemato-biochemical profile of Crossbred cows. *Indian Journal of Animal Health*. **55** (2):141-148.
- Sahatpure, S.; Ravikantha, K.; Maini, S. and Pramod, A. (2016). Ovarian response after treatment with herbal estrus inducer and trace mineral supplement in postpartum anestrous in crossbreed cows. *International Journal of Advance Research*. **4** (7): 2069-2072.
- Samani, N. B.; Jokar, A; Soveid, M; Heydari, M. and Mosavat, S. H. (2016). Efficacy of the Hydroalcoholic Extract of *Tribulus terrestris* on the Serum Glucose and Lipid Profile of Women with Diabetes Mellitus: A Double-Blind Randomized Placebo-Controlled Clinical Trial". *J Evid Based Complementary Altern Med*.
- Santos, P. A. G.; Figueiredo, A. C.; Lourenço, P. M. L.; Barroso, J. G.; Pedro, L. G.; Oliveira, M. M.; Schripsema, J.; Deans, S. G. and Scheffer, J. J. C. (2002). *Hairy root cultures of Anethum graveolens (dill): establishment, growth, time-course study of their essential oil and its comparison with parent plant oils"* (PDF). *Biotechnology Letters*. **24** (12): 1031–1036.
- Singh, K.; Singh, N.; Chandy, A. and Manigauha, A. (2012). In vivo antioxidant and hepatoprotective activity of methanolic extracts of *Daucus carota* seeds in experimental animals. *Asian Pacific Journal of Tropical Biomedicine*. **2** (5): 385-8.
- Terzano, G. M. (2012). Overview on reproductive endocrine aspects in buffalo. *Journal of buffalo science*, **1**(4):126-138.
- Verma, D. M.; Balakrishnan, N. P. and Dixit, R. D. (1993). Flora of Madhya Pradesh (Vol. I). Pbl. Botanical survey of India, Kolkata, India: 490.
- Westwood, C. T.; Lean, I. J. and Garvin, J. K. (2002). Factors influencing fertility of Holstein dairy cows: a multivariate analysis. *Journal of Dairy Science*. **85**: 3225-3237.