

“GROWTH, YIELD AND FRUIT QUALITY OF CAPSICUM HYBRIDS (*Capsicum annuum*) AS AFFECTED BY INTEGRATION OF INORGANIC FERTILIZERS AND ORGANIC MANURES (FYM) IN PROTECTED CONDITION”

ABSTRACT

A trial was conducted at Horticulture Research Farm, Department of Horticulture, during the year 2023-24. Five Capsicum varieties Names Nemalite, Cebrail, Bungl, Volante and Shehzadi. were evaluated at SHUATS, Prayagraj in randomized block design with three replications during Jan-April 2023-24 to evaluate the best performing variety in terms of growth, yield and quality. The variety Cebrail performed best in terms of plant height (68.97 cm), number of branches (5.65), days to first flower initiation (42.89), chlorophyll content (SPAD Value) (45.71 cm), number of fruits per plant (8.53), fruit length (9.2 cm), individual fruit weight (160.44 g), fruit yield per plant (1.09 kg), maximum fruit yield per plot(6.56 kg). The highest benefit cost ratio was found at 3.15 in the same variety.

Keywords: Capsicum; Performance; Varieties

1. Introduction

Capsicum (*Capsicum annuum L.*) moreover called Simla Mirch in India is one of the most important vegetables grown in open situations further to beneath managed situations. Its financial significance as a excessive-fee vegetable crop both in home and remote places markets. due to more consumer alternatives and use in diverse culinary products, satisfactory production of capsicum is the need of the day.

1.1. Why Integrated Nutrient Management?

It is the combined application of chemical fertilizers and organic manures for crop manufacturing. it's miles ecologically, socially and economically feasible. It will increase crop yield and first-rate, It minimise the quantity of synthetic and natural fertilisers you use, It reduce ammonia emissions and enhance air first-class and human health, It lessen damage to sensitive habitats due to extra nutrients carried in the air or via water, It lessen greenhouse fuel emissions and assist lessen the outcomes of climate alternate.

Importance of Integrated Nutrient Management

- Nutrient management helps to reduce the contamination of waterways by plant nutrients.
- Improve soil fertility.
- Enhance plant productivity.
- Reduce the cost of chemical fertilizers.
- Providing balanced nutrition to crops.
- Promotes carbon sequestration and prevents the deterioration of soil, water, ecology, and also leaching of nutrients from the soil.
- Help to check the emerging deficiency of nutrients other than NPK.
- It brings economy and efficiency to fertilizer use and favourably affects the physical, chemical and biological environment of soil.

2.MATERIALS AND METHODS

This experiment was carried out during in 2023 at Horticulture Research Farm, SHUATS, Prayagraj, Uttar Pradesh, India which is located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. This area is situated on the right side of the river Yamuna by the side of Prayagraj Rewa Road about 5 km away from Prayagraj, city. The experiment was done in the polyhouse. The highest O₂ and CO₂ concentration was 20.2% and 480 ppm respectively. In the polyhouse the maximum temperature was 34.7 °C. The maximum light intensity inside and outside the polyhouse was 63600 lux and 110600 lux.

The temperature reached up to 48°C in summer and in winter it goes down to as low as 2-3 °C. The experiment was laid out in a randomized block design with 9 treatments and three replications. The crop was grown in a naturally ventilated polyhouse. 30 days old seedlings were used for transplanting. November was the first week that transplants were performed at 60 cm × 45 cm spacing on the raised bed. The plants were trained along a plastic thread tied to GI wire stretched over beds. The necessary recommended cultural practices like fertilizer application, irrigation and weeding.

Treatment Combinations

TREATMENT	COMBINATION
T0(Nemalite)	100% NPK+15t FYM
T1(Nemalite)	100% NPK + 15t FYM + AZOTOBACTER (10 g)
T2(Cebrail)	100% NPK+15t FYM
T3(Cebrail)	100% NPK + 15t FYM + AZOTOBACTER (10 g)
T4(Bungi)	100% NPK+15t FYM
T5(Bungi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)
T6(Volante)	100% NPK+15t FYM
T7(Volante)	100% NPK + 15t FYM + AZOTOBACTER (10 g)
T8(Sehezadi)	100% NPK+15t FYM
T9(Sehezadi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)

3. STATISTICAL ANALYSIS

The data recorded during the course of the investigation were subjected to statistical analysis as per the method of analysis of variance Fisher (1950). This investigation was done in the RBD statistical method. The significance and non-significance of the treatment effect were judged with the help of 'f' value (variance ratio) and compared with the table value at a 5% level of significance. If the calculated value exceeds then the value, the effect of considered to be significant. The significant difference between the means was tested against the critical difference at a 5% level of significance.

4. Results and Discussion

Observations were recorded for growth parameters which are Plant height (cm), Number of primary branches, Days taken for flower initiation and chlorophyll content. In yield parameters are number of fruits per plant, fruit length (cm), fruit diameter (cm), average fruit weight (g), fruit yield (kg plant⁻¹), individual fruit Weight (gm), fruit yield (200sq. m). In quality parameters are total soluble solids (°Brix), vitamin-C (mg/100g). In economics parameters are cost of cultivation (Rs/200sq. m), Gross Return (Rs/200sq. m), Net Return (Rs/200sq. m) and Benefit Cost Ratio (Rs/200sq. m).

4.1 Growth Parameter

4.1.1 Plant height (cm)

The Maximum plant height at 90 DAP was recorded in the Variety Capsicum Cebrail (T2) (68.97 cm) given in (Table 1). This might be due to the genetic constitution of the varieties. The differential response of vegetative growth of the different may be due to differences in genetic constituents of the varieties.

4.1.2 Number of primary branches

The Maximum Number of branches was recorded in the Variety Capsicum Cebrail (T2) (5.65) given in (Table 1). It is due to the organic treatment of cebrail (T2) was good consumption for those capsicum plants. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition. The differential response of vegetative growth

of the different varieties may be due to differences in genetic constituents of the varieties and microclimate condition.

4.1.3 Days taken for flower initiation

The Early days to first flower initiation were recorded in the Variety Capsicum Cebraill (T2) (42.89) given in (Table 1). This may be due occurrence of early flowering is basically a genetic character of each variety. However, favourable temperature regime in protected conditions for a longer period showed a great impact on the genetic constitution of plant to express its full genetic potential. Better environmental conditions and available nutrients seems to have brought quick changes in plant growth and development.

Table 1 Height of Plant (cm), number of primary branches, days taken for flower initiation and chlorophyll as affected by inorganic fertilizers and Organic manures in Capsicum varieties in

Treatment	Treatment Combination	Plant height (cm)	Number Of Branches	Days to First Flower Initiation
T0(Nemalite)	100% NPK+15t FYM	67.76	5.03	43.56
T1(Nemalite)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	65.86	5.06	43.44
T2(Cebraill)	100% NPK+15t FYM	68.97	5.65	42.89
T3(Cebraill)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	67.91	5.61	43.33
T4(Bungi)	100% NPK+15t FYM	61.34	4.01	48.56
T5(Bungi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	61.30	4.01	48.67
T6(Volante)	100% NPK+15t FYM	62.78	4.03	46.67
T7(Volante)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	62.67	4.04	47.00
T8(Sehezadi)	100% NPK+15t FYM	63.00	4.35	49.22
T9(Sehezadi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	63.18	4.36	49.00
	F test	S	S	S
	S.E (d) (±)	0.89	0.03	1.21
	CD 0.05	1.87	0.07	2.55
	C.V	1.69	0.84	3.21
protected				conditio

4.2 Yield parameters

4.2.1 Number of Fruits Per Plant

The Maximum Number of fruits per plant was recorded in the Variety Capsicum Cebraill (T2) (8.53) given in (Table 2). This might be due to the favourable climatic conditions and sufficient accumulation of photosynthesis in the polyhouse condition.

4.2.2 Fruit length (cm)

The Maximum Fruit length (cm) was recorded in the Variety Capsicum Cebraill (T2) (9.2 cm) given in (Table 2). Increased fruit size in different hybrids, might be due to enhanced photosynthesis accumulation of carbohydrates and favourable effect on vegetative growth which increased the fruit variety besides increasing fruit size.

4.2.3Fruit diameter (cm)

The Maximum Fruit diameter was recorded in the Variety Capsicum Volante (T6) (8.04 cm) given in (Table 2). It might be increased fruit size attributed in different hybrids might be due to enhanced

photosynthesis, accumulation of carbohydrates and favourable effect on vegetative growth which increased the fruit variety besides increasing the fruit size

4.2.4 Individual fruit weight (g)

The Maximum individual fruit weight was recorded in the Variety Capsicum Cebrail (T2) (160.44 g) given in (Table 2). Due to increased fruit weight may be attributed to the favourable microclimate that prevailed in the polyhouse compared to other structures.

4.2.5 Fruit yield per plant (kg)

The Maximum fruit yield per plant was recorded in the Variety Capsicum Cebrail (T2) (1.09 kg) given in (Table 2). It might be due to the organic condition of Cebrail (T2) was higher number of flowers per plant, fruits per plant, more pollination, lesser flower drop, maximum percent fruit set, maximum mean fruit weight and fruit volume.

4.2.6 Fruit yield (kg/m²)

The Maximum fruit yield per plot was recorded in the Variety Capsicum Cebrail (T2) (6.56 kg) given in (Table 2). The higher fruit yield under this condition may be attributed to the favourable climatic conditions that prevailed under polyhouse and also due to its protective ability against major biotic stress, which reduces the effect of the excess rainfall, water logging, and provides a controlled environment to the crop.

Table 2 Number of Fruits Per Plant, Fruit Length (cm), Fruit Diameter (cm), Individual Fruit Weight (gm), Fruit Yield Per Plant (kg) and Fruit yield per plot (kg/m²) as affected by inorganic fertilizers and Organic manures in Capsicum varieties in protected condition

Treatment	Treatment Combination	Number of Fruits Per Plant	Fruit Length (cm)	Fruit Diameter (cm)	Individual Fruit Weight (gm)	Fruit Yield Per Plant in(kg)	Fruit yield per plot (kg/m ²)
T0(Nemalite)	100% NPK+15t FYM	7.34	8.7	7.03	141.56	1.03	6.18
T1(Nemalite)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	7.31	8.7	7.04	144.44	0.99	5.96
T2(Cebrail)	100% NPK+15t FYM	8.53	9.2	6.98	160.44	1.09	6.56
T3(Cebrail)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	8.43	8.9	7.00	158.56	0.99	5.96
T4(Bungi)	100% NPK+15t FYM	7.22	6.3	5.97	102.11	0.84	5.06
T5(Bungi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	7.90	6.2	6.26	99.00	1.07	6.44
T6(Volante)	100% NPK+15t FYM	7.42	8.8	8.04	139.00	0.96	5.74
T7(Volante)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	7.44	8.9	8.00	132.22	0.90	5.42
T8(Sehezadi)	100% NPK+15t FYM	7.93	5.6	5.38	85.67	0.53	3.20
T9(Sehezadi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	7.33	5.5	5.34	88.11	0.60	3.60
F test		S	S	S	S	S	S
S.E (d) (±)		0.29	0.14	0.35	3.34	0.01	0.09
CD 0.05		0.61	0.30	0.74	7.02	0.03	0.18
C.V		4.65	2.28	6.41	3.27	1.85	1.96

4.3 Quality parameters

4.3.1 TSS Content (°Brix)

The Maximum Tss (°Brix) was recorded in the Variety *Capsicum* Volante (T7) (9.34) is given in (Table 3). TSS is an important quality attribute of capsicum fruit. An increase in this parameter improves the flavour and increases the palatability. Since capsicum is used for salad making, fruits with high TSS are highly preferred.

4.3.2 Vitamin C (mg/100g)

The Maximum Vitamin C content (mg/100g) was recorded in the Variety *Capsicum* Sehezadi (T8) (154.66) is given in (Table 3). Generally, the higher ascorbic acid content would increase the nutritive value of capsicum, which would help better retention of colour and flavour. Capsicum varieties and hybrids possessing high ascorbic acid content are in great demand in export markets which may be due to differences in genetic constituents of the varieties and microclimate condition.

4.3.3 Chlorophyll (mg/cm⁻²)

The Maximum Chlorophyll content (mg/cm⁻²) was recorded in the Variety *Capsicum* Cebrail (T2) given in (Table 3). This means that less photosynthesis would occur in the leaves of the plant, so less glucose is made as a result. Therefore there is less energy released for growth as glucose is needed for respiration.

Table 3 TSS (°Brix) & Vitamin C (mg/100g) content as affected by inorganic fertilizers and Organic

Treatment	Treatment Combination	Tss(°Brix)	Vitamin C (mg/100g)	Chlorophyll (mg m ⁻²)		
T0(Nemalite)	100% NPK+15t FYM	7.56	152.00	42.32		
T1(Nemalite)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	9.26	153.22	43.03		
T2(Cebrail)	100% NPK+15t FYM	8.70	152.03	45.71		
T3(Cebrail)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	8.74	152.15	43.31		
T4(Bungi)	100% NPK+15t FYM	9.00	152.56	45.20		
T5(Bungi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	9.08	154.10	43.86		
T6(Volante)	100% NPK+15t FYM	7.59	154.48	42.40		
T7(Volante)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	9.34	152.56	44.39		
T8(Sehezadi)	100% NPK+15t FYM	8.83	154.66	43.41		
T9(Sehezadi)	100% NPK + 15t FYM + AZOTOBACTER (10 g)	8.68	154.05	43.07		
	F test	S	S	S		
	S.E (d) (±)	0.05	1.19	1.48		
	CD 0.05	0.12	2.50	3.11		
	C.V	0.77	0.95	4.16		
manures	in	Capsicum	varieties	in	protected	condition

Table 4 Cost of agronomical practices of cultivation as affected by inorganic fertilizers and Organic manures in Capsicum varieties in protected condition

S. No	Particular	Requirement	Rate/unit Rs.	Cost
(A)	Land preparation			
I.	Soil Pulverization	4 labours	350 Rs/labours	1,400
II.	Layout of field	2 labours	350 Rs/labours	700
III.	Mulching (30 Micron) Silver black			6,000
(B)	Planting	1 labour	350 Rs/labour	350
(C)	Manures and fertilizer			
I.	FYM	300kg	10 Rs/kg.	3,000
II.	Urea	60 Kg	10 Rs/Kg	600
III.	DAP	50 Kg	30 Rs/Kg	1,500
IV.	MOP	40 Kg	50 Rs/Kg	2,000
V.	Labour	1 labours X 3 time	350 Rs/labour	1,050
(D)	Irrigation	1 Labour X 3 time	350	1,050
(E)	Weed Management	1 labour X 3 time	350 Rs/labour	1,050
(F)	Harvesting	1 labours X 3 time	350 Rs/labour	1,050
	Total cost of cultivation			19,750

Table 5 Cost economics of growing different varieties of capsicum as affected by inorganic fertilizers and Organic manures in Capsicum varieties in protected condition

Treatments	Cost of cultivation (Rs/200sq. mt)	Total yield 200sq. mt	Selling Rate (Rs/q)	Gross return (Rs/200sq. mt)	Net return Rs./ 200sq. mt)	Benefit cost ratio
T0(Nemalite)	19,750	15.45	5000	77,250	57,500	2.91
T1(Nemalite)	19,750	14.90	5000	74,500	54,750	2.77
T2(Cebrail)	19,750	16.40	5000	82,000	62,250	3.15
T3(Cebrail)	19,750	14.90	5000	74,500	54,750	2.77
T4(Bungi)	19,750	12.65	5000	63,200	43,500	2.20
T5(Bungi)	19,750	16.10	5000	80,500	60,750	3.07
T6(Volante)	19,750	14.35	5000	71,750	52,000	2.63
T7(Volante)	19,750	13.55	5000	67,750	48,000	2.43
T8(Sehezadi)	19,750	8.00	5000	40,000	20,250	1.02
T9(Sehezadi)	19,750	9.00	5000	45,000	25,250	1.27

5. Conclusion

From the present investigation it is concluded that variety Cebrail (T2) performed best in terms of Growth parameters viz., plant height (68.97 cm), Number of branches (5.65), Days to first flower initiation (42.89), Chlorophyll content (45.71). In terms of Yield Parameter number of fruits per plant (8.53), Fruit length (9.2 cm), individual fruit (160.44 g), fruit yield per plant (1.09 kg), fruit yield per plot (6.56 kg). Also, in terms of Economics, Variety Cebrail (T2) recorded the highest Benefit cost ratio (3.15).

REFERENCE

- Jeevansab. (2000).** Effect of nutrient sources on growth, yield and quality of capsicum grown under different environments. M.Sc. (Agri.) Thesis, Univ. Agric. Sci. Dharwad, Karnataka (India).
- Jose, J. L., Jesus, L. E., Huez, L., Alejandro, M. and Garcia., 2013.** Post harvest quality and shelf life of green pepper (*Capsicum annuum* L.) grown under open field and greenhouse conditions. *Egyptian J. Hort.*, 19 (5): 326-337.
- Joshi, V.C.Yadav V.K, Nautiyal, M K and Girish Tiwari. 2010.** Assessment of different capsicum (*Capsicum annuum* L.) genotypes under protected conditions in mid hills of Himalaya. National Symposium on Abiotic and Biotic Stress Management in Vegetable Crops.
- Kamaruddin, R. (2007).** Design and development of naturally ventilated tropical crop protection structures and hydroponics systems. *Acta-Horticulturae.* (742): 139-153.
- Kohnic, A., Ostojic, I. and Karic, N. (2006).** Vegetable pests in greenhouses in territory of Herzegovina. *Radovi Poljoprivrednog Fakulteta Univerziteta u Sarajevu* Works of the Faculty of Agriculture University of Sarajevo. 51(57(2)): 139-140.
- Sreedhara, D. S., Kerutagi, M. G., Basavaraja, H., Kunnal, L. B. and Dodamani, M. T., 2013,** Economics of capsicum production under protected conditions in northern Karnataka. *Karnataka J. Agric. Sci.*, 26 (2): 217-219.
- Sweta Rani. 2003.** Variability and path coefficient studies in bell pepper (*Capsicum annuum* L.) M.Sc thesis submitted to Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan.(H.P).
- Takte, R. L, Ambad, S. N, Kadam, U. S, and Dhawale, B. C. 2003.** Greenhouse cladding material, shade nets, ventilation etc. Proceedings of All India on seminar on Potential and Prospects for Protective Cultivation, organized by the Institute of Engineers, Ahmednagar, December, 12-13, 2003, pp. 117-119.
- Takte, R. L., Ambad, S. N., Kadam, U. S. and Dhawale, B. C. (2003).** Greenhouse cladding material, shade nets, ventilation etc. Proceedings of All India on seminar on Potential and Prospects for Protective Cultivation, organized by the Institute of Engineers. pp. 117-119.
- Valerio, E., Cecilio, A. and Mexia, A. (2007).** Interactions between aphid species and beneficial organisms in sweet pepper protected crop. *Boletín de Sanidad-Vegetal,-Plagas.* 33(2): 14
- Verma, T.S, Yadav, H.L, Singh, R.V, Sharma, S.C. and Thakur, P.C 2003.** Assessment of hybrids in capsicum for marketable suitability. *Scientific Horticulture.* 8: 101-106. 5.
- Vethamoni, P. I. and Natarajan, S. 2008.** Cultivation of sweet pepper cultivars (*Capsicum annuum* var. *grossum* L.) under shade net in tropical plains of Tamil Nadu. *Asian Journal of Horticulture.* 3(2):372-376
- Vimlesh, K., Naresh, K., Pathania and Nimit, K., 2015,** Evaluation of bell pepper (*Capsicum annuum* L. var. *grossum* Sendt.) genotypes for quality traits in modified naturally ventilated polyhouse. *Ann. Hort.*, 8 (1): 69-72.
- Yellavva and Patil, A. A. 2009.** Performance of coloured capsicum hybrids under different protected structures. *Karnataka Journal of Agricultural Sciences.* 22(5): 1058-1061.
- Zende, U. M., 2008,** Investigation on production techniques in capsicum under protected cultivation. *M. Sc. (Agri.) Thesis*, Univ. Agric. Sci., Dharwad, Karnataka (India).
- Abdullah, A., Mahmoud, W., Hesham, A. R. and Abdullah, I., 2013,** Effects of pruning systems on growth, fruit yield and quality traits of three greenhouse grown bell pepper (*Capsicum annuum* L.) cultivars. *Australian J. Crop Sci.*, 7 (9) : 1309-1316.
- Ahmed, M. A., Mahmoud and Ahmed, A.S., 2015,** Production and evaluation of high yielding sweet pepper hybrids under greenhouse conditions. *J. Agric. Environ. Sci.*, 15 (4): 573-580.
- Akhilesh, O, Mishra, S, Singh, R.V. and Harihar Ram. 2001.** Character association in capsicum (*Capsicum annuum* L.) genotypes. *Progressive Horticulture.* 33:138-143.
- Aruna, P. and Sudagar, I.P. 2010.** Evaluation of capsicum varieties under poly house conditions. *Asian Journal of Horticulture.* 4 (2):336-337.
- Arya, P. S., 1999,** Vegetable Breeding and Seed Production. Kalyani Publishers, Ranjinder Nagar, Ludhiana, India. pp. 318.
- Awani Kumar Singh, Balraj Singh and Ramawant Gupta.2011.** Performance of sweet pepper (*Capsicum annuum*) varieties and economics under protected and

open field conditions in Uttarakhand. *Indian Journal of Agricultural Sciences*. 81(10):973-5.

Backer, J. C., 1989, The effect of air humidity on flowering, fruit set and fruit growth of glass house sweet pepper (*Capsicum annuum* L.). *Scient. Hort.*, 40: 1-8.

Banaras, M., Bosland, P. W. and Lownds, N. K., 2005, Effects of harvest time and growth conditions on storage and post-storage quality of fresh peppers (*Capsicum annuum* L.). *Pakistan J. Biotech.*, 37 (2): 337-344.

Basavaraja, N, Nandi, V. R. and Praveen Jhologikar, 2003. Protected cultivation of capsicum and bhendi. Proceedings of All India Seminar on Potential and Prospects for Protective Cultivation, Organized by the Institute of Engineers, Ahmednagar, December, 12-13, pp. 197-199.

Belko, I. and Valsikova, M., 2004, Evaluation of sweet pepper (*Capsicum annuum* L.) assortment. *Capsicum Eggplant Newsletter*, 23: 21-24.

Bergefurd, B. R., Lewis, W., Harker, T., Miller, L., Welch, A. and Weaks, E., 2011, Bell pepper cultivar performance trial grown in southern Ohio. www.southcenters.osu.edu.

Bhatt, R. M. and Rao, N. K. S., 1993, Response of bell pepper to photosynthesis, growth, flower and fruit setting to night temperature. *Photosynthetica*, 28: 127-132.

Biwalkar, N., Singh, K. G., Jain, A. K., Sharda, R., Jindal, S. K., Singh, K and Chawla, N. 2015. Response of coloured sweet pepper (*Capsicum annuum* L. var. grossum) to fertigation and irrigation levels under naturally ventilated greenhouse. *Agricultural Research Journal*. 52(1): 19-25.

Chate, B.R, Mangave, K.K. and Jogdand, S.M. 2012. Performance of different cultivars of sweet pepper (*Capsicum annuum* L.) under shade house condition. *Ecology, Environment and Conservation paper*. 18 (3):573-578.

Chaudhary, G-N. and Mirza, M. (2012). Greenhouse pepper production economics. *Peppers: botany, production and uses*. 255-269.

Choudhary, N., Pathania, N. K., Singh, S. P., Pardeep, K. and Yudhvir, S., 2011, Evaluation of red and yellow capsicum hybrids for quality attributes in naturally ventilated polyhouse in mid hills of western Himalayas. *Veg. Sci.*, 38: 218-220.

Clemencia, L. and Sumagaysay, 2015, Plastic tunnel production of bell pepper grown during wet and dry season. *Int. J. Edu. Res.*, 3 (7): 45.

Dargie, T., Bizuayehu, T., Haddis, Y. and Andnet, B., 2013, Effects of harvesting stage and storage duration on postharvest quality and shelf life of bell pepper (*Capsicum annuum* L.) varieties under passive refrigeration system. *Global Sci. Res. J.*, 1 (1): 027-033.

Vimala et al. (2007), Effect of organic and inorganic fertilizers on growth, yield and nutrient content of bird chilli (*Capsicum frutescense*) (*J. Trop. Agric. and Fd. Sc.* 35(1)(2007): 29– 40)

Laware et al., (2010), Effect of Novel Organic Liquid Fertilizer on Growth and Yield in Chilli (*Capsicum annuum* L.) (*ASIAN J. EXP. BIOL. SCI. SPL*.2010 :15-19)

Malik et al., (2011), Growth, yield and fruit quality of sweet pepper hybrid SH-SP-5 (*Capsicum annuum* L.) as affected by integration of inorganic fertilizers and organic manures (FYM) (*Journal of Agricultural Technology* 2011 Vol. 7(4):1037-1048)

Dorji et Al., (2011), Effect of different rates and combinations of farm yard manure and inorganic fertilizers on chilli (*Capsicum annuum*) yield (*National Soil Services Centre, Ministry of Agriculture*)

López et al., (2011), Response of chilli pepper (*Capsicum annuum* L.) to salt stress and organic and inorganic nitrogen sources: growth and yield (tropical and subtropical agroecosystems, 14 (2011): 137 – 147)

Reddy et al., (2017), Study of different organic manure combination on growth and yield of chilli (*Capsicum annuum* L.) (*Plant Archives* Vol. 17 No. 1, 2017 pp. 472-474)

Altaf et al., (2019), Effect of NPK, organic manure and their combination on growth, yield and nutrient uptake of chilli (*Capsicum Annum* L.) (*Horticulture Int J*. 2019;3(5):217–222.)

Elad, Y, Y, Messika, Y, Brand, M, David, D.R. and Szejnberg, A. 2007. Effect of colored shade nets on pepper powdery mildew (*Leveillulataurica*). *Phytoparasitica*. 35:285-299.

Hajarica, T. K. and Phookan, D. B., 2005, Performance of tomato cultivars for polyhouse cultivation during spring summer in Assam. *Indian J. Hort.*, 62 (3): 268-271.

Hutton, M.G. and Handley, D.T. 2007. Bell pepper cultivar performance under short, variable growing seasons. *Hort Technology*. 17 (1):136 141.9.

Ilic, Z, Milenkovic, L, Bodroza-Solarov, M, Marinkovic, D. and Sunic, L. 2012. Tomato fruits quality as affected by light intensity using color shade nets. 47th Croatia and 7th International Symposium on Agriculture. Opatija. Croatia. *Sym. Proc.* 414-418.

Rahul Rathore and Vijay Bahadur, 2024 Assessment of Capsicum (*Capsicum annuum*) Varietal Performance in Natural Ventilation Polyhouse Settings International Journal of Environment and Climate Change Volume 14, Issue 2, Page 128-134, 2024; Article no. IJECC.109247 ISSN: 2581-8627