

## Original Research Article

### Economic cost analysis Feasibility of including Probiotics in Supplementation the feed of caged Caged broilers Broilers

#### Abstract

As part of the Sam Higginbottom University of Agriculture, Technology, and Sciences in Prayagraj, this experiment was carried out at the Small Animal Lab that which is part of the research unit. In this study, we investigated the “Economic cost analysis of including probiotics in the feed of caged broilers”. We used a total of 96 day-old chicks and were randomly assigned. Each of the four groups of 24 chicks (T<sub>0</sub>, T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub>) contained eight replicates each, which meant that each group contained twenty-four chicks. A total of 96 day-old chicks were randomly divided into these groups. The groups were as follows: T<sub>0</sub> control (untreated) group was fed the basal diet (BD), T<sub>1</sub> was given fed *Bacillus coagulans* at a concentration of 40 parts per million ppm in addition to the basal diet BD, T<sub>2</sub> was given fed *Bacillus B. subtilis* at a concentration of 50 parts per million mixed ppm in with the basal diet BD, and T<sub>3</sub> was fed *B. licheniformis* and *B. coagulans* given the basal diet at a concentration of 20 parts per million ppm in BD together with *Bacillus licheniformis* and *Bacillus coagulans*. After a 35-day (five week) trial period, an analysis of the economic effects of adding the probiotics supplementation in feed was calculated. The major difference was noted due to results revealed a decrease in FCR as birds achieved more weight, leading to increased body weight and resulting in a higher price per kg. It was also noted that adding probiotics to the feed doesn't increase the variable cost, and treatment groups had a lower variable cost than the control group. From the above study, it was found that the maximum profit was obtained in T<sub>3</sub>, i.e., (Rs. 5771.32), followed by T<sub>0</sub> (Rs. 5297.3), and T<sub>2</sub> (Rs. 5074.92), whereas the T<sub>1</sub> group had the least revenue. Therefore, it can be concluded that mixing probiotics in poultry feed increases profit without much affecting the production cost.

**Keywords:** Cost, Economics, FCR, Probiotics, Profit and Revenue.

#### 1. INTRODUCTION

Among the agricultural and veterinary industries, poultry is experiencing rapid expansion. Poultry production spends a significant amount on feed, which accounts for 70% of the total. Poultry growers

Formatted: Font: Not Bold

are seeing a decline in their profit margins due to the ever-increasing cost of compounded feed and other poultry feed additives. Consequently, the most crucial requirement for optimal genetics for economically viable chicken production is balanced and effective feeding. ~~A number of~~ ~~Some~~ feed additives, including growth promoters, synthetic hormones, and antibiotics, have been widely utilized to increase ~~the poultry output of chickens~~. Nevertheless, **antibiotic growth promoters (AGPs)** used in poultry have the potential to impact human health through residues in chicken products and the development of antibiotic-resistant microorganisms. Using **AGPs** in chicken feed is illegal in many European nations. As a result, a lot of studies have focused on finding alternatives that can meet the needs of consumers and international markets without compromising production or being hazardous to animals or humans. One of these approaches that stands out is ~~adding probiotics to animal feed~~. Feed additives containing probiotics reduce the risk of gastrointestinal diseases by increasing the ~~human~~ immune response, secreting antimicrobial compounds, and ~~preventing the exerting a~~ competitive exclusion ~~of of~~ harmful bacteria [1]. These products have been shown to have no residue in animal products and have been found to enhance animal performance and health. This has been supported by various studies ~~conducted by~~ [2-6]. The reason behind this improvement is that these products enhance the digestibility of the ~~animals' animal's~~ diet, ~~as demonstrated by~~ [7]. This leads to better utilization of nutrients and, consequently, higher productivity, as indicated ~~by studies conducted by~~ [8-10]. Probiotics can be administered to chickens either by mixing them with feed or ~~by by~~ adding them to drinking water [11]. The growth of pathogenic ~~bacteria,~~ including *Escherichia coli* and *Salmonella sp.*, can be inhibited by lactic acid bacteria probiotics, which have been shown to have a favorable effect [12-16]. In addition to producing lactic acid and bacteriocins, lactic acid bacteria ~~have the ability to~~ can colonize the intestines and survive [17]. The probiotic and antimicrobial properties of *Lactobacillus casei* make it a promising candidate for use as a functional probiotic in animal feed [18]. You can treat your birds/animals with probiotics in both feed and water, but it's important to note that probiotics in water can make them gain weight and reduce the feed conversion value. ~~You can find out if~~ ~~Investigating whether it's~~ more profitable to give broiler probiotics in water ~~should be carried out~~ by doing an economic study. ~~Thus we aimed to explore~~ ~~Finding out~~ how much money broiler farms make by switching from ~~antibiotic growth promoters (AGPs)~~ to probiotics like *Bacillus-B.Subtilis*, *Bacillus-B.Coagulans*, and *Bacillus slicheniformis* ~~in order~~ to increase feed conversion and weight was the main motivation for this research.

Formatted: Highlight

Formatted: Highlight

## 2. MATERIALS AND METHODS

As part of the Sam Higginbottom University of Agriculture, Technology, and Sciences in Prayagraj, this experiment was carried out at the Small Animal Lab that which is part of the research unit. In this study, we investigated the economic cost analysis of including probiotics in the feed of caged broilers. We used a total of 96 day-old chicks and were randomly assigned four groups of 24 chicks (T<sub>0</sub>, T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub>) and eight replicates each, into these groups. The groups were as follows: T<sub>0</sub> control group was fed the basal diet (BD), T<sub>1</sub> was fed *Bacillus coagulans* at 40 ppm in BD, T<sub>2</sub> was fed *B. subtilis* at 50 ppm in BD, and T<sub>3</sub> was fed *B. licheniformis* and *B. coagulans* at 20 ppm in BD. Each of the four groups (T<sub>0</sub>, T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub>) contained eight replicates, which meant that each group contained twenty-four chicks. A total of 96 day-old chicks were randomly divided into these groups. T<sub>0</sub> control (untreated) group was fed the basal diet, T<sub>1</sub> was given *Bacillus coagulans* at a concentration of 40 parts per million in addition to the basal diet, T<sub>2</sub> was given *Bacillus subtilis* at a concentration of 50 parts per million mixed in with the basal diet, and T<sub>3</sub> was given the basal diet at a concentration of 20 parts per million together with *Bacillus licheniformis* and *Bacillus coagulans*. After a 35-day (five week) trial period, an analysis of the economic effects of adding probiotics in feed was calculated.

Formatted: Font: Not Bold

### 3. RESULTS AND DISCUSSION

Table 1: Variable cost of treatments

Description	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>
Transport (Rs)	12.5	12.5	12.5	12.5
Feed additive (Rs)	0	0.18	1.58	0.30
DOC broiler (Rs)	768	768	768	768
Feed (Rs)	1533.6	1440	1382.76	1313.88
<b>Total (Rs)</b>	<b>2314.1</b>	<b>2220.68</b>	<b>2164.08</b>	<b>2094.68</b>
T <sub>0</sub> control group was fed the basal diet (BD), T <sub>1</sub> was fed <i>Bacillus coagulans</i> at 40 ppm in BD, T <sub>2</sub> was fed <i>B. subtilis</i> at 50 ppm in BD, and T <sub>3</sub> was fed <i>B. licheniformis</i> and <i>B. coagulans</i> at 20 ppm in BD.				

Formatted Table

Formatted: Font: 10 pt

Formatted: Left

The provided table outlines the breakdown of costs associated with different treatments (Treatment<sub>0</sub>, Treatment<sub>1</sub>, Treatment<sub>2</sub>, and Treatment<sub>3</sub>), along with a Treatment<sub>0</sub> (control group), across various categories. Here's a detailed explanation of each category:

**Transport (Rs):**

• ~~Transport~~ represents the cost of transportation for each treatment group as well as the control group.

• The cost of transportation ~~is was~~ consistent across all groups ~~at Rs 12.5~~.

#### Feed additive (Rs):

• ~~Feed additive~~ displays the cost of feed additives (probiotics) for each treatment group and the control group ~~as shown in Table 1~~.

➤ ~~The cost of feed additives varies across treatments:~~

• ~~Treatment<sub>0</sub> (Control): No cost (Rs 0)~~

• ~~Treatment<sub>1</sub>: Rs 0.18~~

• ~~Treatment<sub>2</sub>: Rs 1.58~~

• ~~Treatment<sub>3</sub>: Rs 0.30~~

#### DOC broiler (Rs):

• ~~DOC~~ indicates the cost associated with Day Old Chicks (DOC) for each treatment group and the control group.

• The cost of DOC broiler ~~remains remained~~ constant across all groups ~~(Table 1) at Rs 768~~.

#### Feed (Rs):

~~Feed~~ represents the cost of feed for each treatment group and the control group.

The cost of feed varies across treatments ~~as shown in Table 1~~.

• ~~Treatment<sub>0</sub>: Rs 1533.6~~

• ~~Treatment<sub>1</sub>: Rs 1440~~

• ~~Treatment<sub>2</sub>: Rs 1382.76~~

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Normal, No bullets or numbering

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Normal, No bullets or numbering

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Normal, No bullets or numbering

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Normal, No bullets or numbering

- ~~Treatment<sub>3</sub>: Rs 1313.88.~~

Formatted: Font: (Default) Times New Roman, 12 pt

**Total (Rs):** ~~is the~~

~~Total Rupees~~ ~~sumsup of~~ the total variable costs for each treatment group and the control group.

It ~~includes~~ ~~included~~ the costs of transportation, feed additives, DOC broiler, and feed.

The total cost varies across treatments ~~as shown in Table 1.~~

- ~~Treatment<sub>0</sub>: Rs 2314.1~~
- ~~Treatment<sub>1</sub>: Rs 2220.68~~
- ~~Treatment<sub>2</sub>: Rs 2164.08~~
- ~~Treatment<sub>3</sub>: Rs 2094.68~~

Each row in the table provides a detailed breakdown of the variable costs associated with transportation; feed additives, DOC broiler, feed, and the total cost for each treatment group and the control group. This breakdown allows for a comprehensive comparison of costs across different treatments and the control group.

**Table 2: Total of revenue and profit for 96 broiler chicken treatment**

Description	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>
Final of body weight production (kg)	40.06	34.40	38.10	41.40
Final of body weight sell/kg (Rs)	190	190	190	190
Total of Revenue (Rs)	7611.4	6536	7239	7866
Total of cost (Rs)	2314.1	2220.68	2164.08	2094.68
Profit (Rs)	5297.3	4315.32	5074.92	5771.32

~~T<sub>0</sub> control group was fed the basal diet (BD). T<sub>1</sub> was fed *Bacillus coagulans* at 40 ppm in BD. T<sub>2</sub> was fed *B. subtilis* at 50 ppm in BD, and T<sub>3</sub> was fed *B. licheniformis* and *B. coagulans* at 20 ppm in BD.~~

Formatted Table

Formatted: Left

### Final body weight production (kg):

~~Final body weight production~~ shows the average final body weight production in kilograms for each treatment group at the end of the treatment period. It indicates the weight gain achieved by the broiler chickens (Table 2).

**Final body weight sell/kg (Rs):**

The selling price per kilogram of broiler chickens remains constant ~~at Rs 190~~ across all treatment groups (Table 2). This indicates the price at which the broiler chickens are sold.

**Total Revenue (Rs):**

~~Total Revenue~~ represents the total revenue generated from selling broiler chickens for each treatment group. It is calculated by multiplying the final body weight production by the selling price per kilogram.

**Total Cost (Rs):**

~~Total Cost~~ displays the total cost incurred for treating 96 broiler chickens in each treatment group. The costs include transportation, feed additives, DOC broiler, and feed, ~~as shown in the previous tables.~~

**Profit (Rs):**

~~Profit~~ shows the profit generated from the sale of broiler chickens for each treatment group. It is calculated by subtracting the total cost from the total revenue, indicating the financial gain from the broiler chicken treatment.

Each row in the table provides detailed information about the final body weight production, revenue, cost, and profit for each treatment group, facilitating comparison and analysis of the effectiveness and profitability of different treatments.

**Discussion:** Probiotics of the type T2, T3, T4, and T5 administered by feed or water can enhance body weight while decreasing feed conversion[19]. Adding 0.25 g probiotic/liter through drinking water yields the highest production performance results and the most profitable economic analysis. Therefore, for optimal agricultural performance and profit, farmers can administer 0.025 g probiotic/liter through water, and study by [20] during the study period, 60% of broiler farms were found to be in operation. When it came to economic losses, broiler flocks were the worst hit, ~~with the exception of except for~~ two farms (B1 and B6), caused by factors such as reduced body weight gain, ~~and~~ increased feed conversion ratio (FCR), mortality, chemotherapy, and chemoprophylaxis. There was no subclinical form of coccidiosis documented in farm B6, loss due to chemotherapy was higher

compared to poor FCR, whereas in farm B6 no subclinical form of coccidiosis was recorded. Chemotherapy was higher than low FCR in farm B2, and mortality was the main cause of loss in farm B1 due to concomitant infection of inflammatory bowel disease.

#### 4. SUMMARY AND CONCLUSION

~~From the above study it~~ was found that the maximum profit was obtained in T<sub>3</sub> ~~i.e.,~~ (Rs. 5771.32), ~~followed by~~ T<sub>0</sub> (Rs. 5297.3), and T<sub>2</sub> (Rs. 5074.92) whereas the T<sub>1</sub> group had the least revenue. The major difference was noted due to a decreased in FCR as birds achieved more weight leading to increased Body-body weight and resulting in more price per kg. ~~It was also noted that~~ a Adding Probiotics-probiotics in to the feed doesn't increased the variable cost and treatment groups had variable cost less than the Control group.

~~Therefore, from the above study it can be concluded that m~~ Mixing probiotics in poultry feed increases the profit/revenue without much affecting the production cost. Hence, probiotics can be included in poultry feed however large-large-scale studies in the future are recommended.

#### REFERENCES

1. Corcionivoschi, N., Drinceanu, D., Stack, D., Ștef, L., Julean, C. and Bourke, B. The effect of probiotics on animal health. Review. Scientific Papers Animal Science and Biotechnologies. 2010; 43(1): 35-35.
2. Fuller, R. Probiotic in man and animals. J. Appl. Bacteriol. 1989; 66: 131-139.
3. Jin, L.Z., Ho, Y.W., Abdullah, N. and Jalaludin, S. Probiotics in poultry: modes of action. World's Poultry Science Journal. 1997; 53(4):351-368.
4. Zulkifli, I., Abdullah, N., Azrin, N.M. and Ho, Y.W. Growth performance and immune response of two commercial broiler strains fed diets containing Lactobacillus cultures and oxytetracycline under heat stress conditions. British poultry science. 2000; 41(5):593-597.

**Comment [es1]:** Although the references are sufficient as 30% (6 out of 20) of the listed references were published in the past five years. Preferred to update and increase the percentage to at least 35-40%. The topic contains a lot of studies including the impact of adding probiotics forms on immunity and performance

5. Ferreira A J P, Pizarro L D C R and Leme I L. Probióticos e prebióticos. Spinosa, H.S.; Gorniak, S.L.; Bernardi, M.M.(Eds.) Farmacologia aplicada à medicina veterinária. 3. ed. Rio de Janeiro: Guanabara Koogan. 2002;574–578.
6. Patterson JA and Burkholder KM. Application of prebiotics and probiotics in poultry production. *Poultry Science*. 2003;82:627–31.
7. Apata D F. Growth performance, nutrient digestibility and immune response of broiler chicks fed diets supplemented with a culture of *Lactobacillus bulgaricus*. *Journal of Science Food Agriculture*. 2008;88:1253–58.
8. Kabir SML, Rahman MM and Rahman MB. The dynamics of probiotics on growth performance and immune response in broilers. *International Journal of Poultry Science*. 2004;3: 361–64.
9. Mountzouris KC, Tsirtisikos P and Kalamara E. Evaluation of the efficacy of a probiotic containing *Lactobacillus*, *Bifidobacterium*, *Enterococcus* and *Pediococcus* strains in promotion of broiler performance and modulation of cecal microflora composition and metabolic activities. *Poultry Science*. 2007; 86:309–17.
10. Mountzouris K C, Tsirtisikos P and Palamidi I. Effects of probiotic inclusion levels in broiler nutrition on growth performance, nutrient digestibility, plasma immunoglobulin, and cecal microflora composition. *Poultry Science*. 2010; 89: 58–67.
11. Utomo A M. Effect of probiotic giving with *Lactobacillus* sp. and *Saccharomyces cerevisiae* against broiler weight gain. (Thesis), Airlangga University. Surabaya. 2012.
12. Huda K, Lokapirnasari W P and Soeharsono S. Addition of probiotic *Lactobacillus acidophilus* and *Bifidobacterium* sp. to the analysis of laying hens infected by *Escherichia coli*. *Mimbar Agribusiness*. 2019; 5:176–82.
13. Najwan R, Lokapirnasari W P and Huda K. Effect of the addition of probiotic *Lactobacillus acidophilus* and *Bifidobacterium* to the production of laying hens infected with *Escherichia coli*. *Briliant*. 2019;4:125–31.
14. Wardhani HCP, Lokapirnasari W P and Soepranionondo K. Use of combination probiotic *Lactococcus lactis* and *Lactobacillus acidophilus* as a substitute of antibiotics in chicken layer infected by *Escherichia coli* on business analysis. *Mimbar Agribusiness*. 2019;5:183–92.
15. Rahman N F N, Lokapirnasari W P, Soepranionondo K, Hidanah S and Sarmanu S. The effect of using *Lactobacillus casei* and *Bifidobacterium* sp. probiotics on feed

conversion ratio and feed consumption in laying hens infected with *Escherichia coli*. *Briliant*. 2019;4:167–73.

16. Nouri, M., Rahbarizadeh, F., Ahmadvand, D., Mousakhani, F., Sadeghzadeh, E., Lavasani, S. and Khodami, V.V. Inhibitory effects of *Lactobacillus salivarius* and *Lactobacillus crispatus* isolated from chicken gastrointestinal tract on *Salmonella enteritidis* and *Escherichia coli* growth. 2010.
17. Salminen S, Wright A V and Ouwehand A. 2004. Lactic acid Bacteria: microbiological and functional aspects, revised and expanded. 3rd ed. Marcel Dekker, Inc. New York. 2004; 177.
18. Amaravathi SC, Mallam M, Manthani GP and Komireddy KR. Effect of dietary supplementation of probiotics and enzymes on the haematology of rabbits reared under two housing systems *Veterinary World*. 2012;5:748–53.
19. Andriani, A.D., Lokapirnasari, W.P., Karimah, B., Hidanah, S. and Al-Arif, M.A., 2020. Potency of probiotic on broiler growth performance and economics analysis. *Indian Journal of Animal Sciences*. 2020; 90(8): 1140-1145.
20. Pant, S., Bhatt, P. and Singh, A.K. Economic impact assessment and disease prevalence of coccidiosis in broilers around tarai region. 2019; 548-655.