

Anti-hyperlipidaemic effect of *Moringaoleifera* ethanol extract on high fat diet fed rat

Abstract

Background: The art or practice of herbal remedies refers to the use of herbs and herbal treatments for the purpose of maintaining health and preventing, treating, or curing sickness. In some areas, herbal treatments can also be referred to as herbal medicine.

Aim: The present study was designed to investigate the antihyperlipidemic and antiatherogenic potentiality of ethanolic extract of *Moringaoleifera* extract in high fat diet-induced hyperlipidemic rats.

Materials and method: *Moringaoleifera* ethanolic extract was prepared using Soxhlet apparatus. Male rats were made hyperlipidemic by giving high fat diet. *M. oleifera* was administered in a dose of 250, 500 and 750 mg/kg.b.w./day for 30 days in high fat diet-induced hyperlipidemic rats. The SGPT, SGOT, total cholesterol, triglyceride, LDL, HDL, Urea and creatinine levels were measured after the treatment.

Results: For both the SGPT and the SGOT, it was seen that groups 5 and 6 exhibited statistically significant ($p < 0.05$) outcomes in the case of the SGPT. When conducting the renal function test, it was observed that the levels of creatinine and urea were statistically significant ($p < 0.05$) in the cases of groups 4, 5, and 6. In the case of high-density lipoprotein (HDL) and low-density lipoprotein (LDL), groups 4, 5, and 6 showed statistically significant findings ($p < 0.05$) in HDL levels, while groups 5 and 6 showed statistically significant LDL levels. The triglyceride levels in the group were found to be statistically significant ($p < 0.05$), while the findings obtained from groups 5 and 6 were also found to be statistically significant.

Conclusion: The study revealed that *M. oleifera* has anti hyperlipidemic activity.

Keywords: Herbal medicine, *Moringaoleifera*, HDL, LDL, Phytochemicals.

Introduction

The liver, the biggest glandular organ, regulates the bulk of an individual's physiological processes. Throughout the day, the liver receives the whole blood supply of a person on many occasions. It plays a crucial role in human metabolism [1]. The abundance of lipoproteins and triglycerides (TG), which are subject to the effects of hereditary and environmental variables, serves as an indicator of the lipid metabolism process. Dyslipidemia may result from either intrinsic factors, exogenous factors, or a combination of genetic predisposition and external effects [2]. Statins are the most recognised and widely accepted therapy for hypercholesterolemia. Their efficacy is well acknowledged, and when used at the maximum allowable dose in combination with ezetimibe, they often facilitate the achievement of LDL-C goals [3]. But these drugs associated with damage of liver enzymes, cancer, diabetes, renal alternation and stroke etc[4]. So medicinal plant researcher are searching for new treatment with lower cost and fewer side effects. Some chemical components derived from medicinal plants may have therapeutic uses, say specialists in the subject. It follows that researchers are always on the lookout for new herbal cures and other plant-based therapies to treat a wide range of illnesses [5]. While phytotherapy is based on scientific study, herbalism is more concerned with the practical applications of medicinal plants. Plants have played a significant role in human medicine for thousands of years due to the wide diversity of chemicals they contain, many of which have medicinal characteristics [6]. The vast variety of chemical components found in medicinal plants allows them to exert a broad spectrum of pharmacological and therapeutic effects. Tanning agents, glycosides, alkaloids, saponins, polysaccharides, essential oils, terpenoids, resins, and plant lipids are all examples of such components [7-9]. Modifying plants genetically allows for the precise regulation of chemical concentrations, allowing for the desired therapeutic effect. Reverse genetics has many potential applications, one of which is to enhance the production of secondary metabolites like alkaloids [10]. Recent scientific progress around the world has led to more research into the healing properties of plants [11]. This is because plants are safe, have strong pharmacological activity, and are more cost-effective than man-made drugs.

Moringa oleifera has nutritional and therapeutic features as a result of its tremendous medical potential; however, this is only the case if the economic worth of the plant's nutritional value, medicinal applications, and animal feed is significant. This tropical deciduous tree, which is endemic to the southern Himalayas in northern India, is a perennial and belongs to the Moringaceae family. Antioxidant, anti-inflammatory, neuroprotective, hypoglycemic, and blood

lipid-lowering are just a few of the nutritional and medicinal benefits of *Moringa oleifera* extracts. *Moringaoleifera's* rich phytochemical content—including flavonoids, glucosinolates, isothiocyanates, and phenolic acids—is strongly associated with its positive effects [12].

The purpose of our present study is to evaluate the anti-hyperlipidemic effects of *Moringaoleifera*.

MATERIALS AND METHODS

Plant Collection and Extract Preparation

Moringa oleifera were collected from local market of Dhaka. The material was authenticated by National herbarium, Bangladesh. Firstly *Moringaoleifera* was cleaned properly with water and it was then air-dried. Finally dried leaves were crushed in powder. The powder was soaked for 10 days in 70% ethanol. Vigorous shaking was also performed occasionally. Next, the solution was filtered. The collected filtrate was dried in a rotary evaporator at a low temperature and pressure. Finally, the crude residue was subjected to the required pharmacological testing.

Drugs and Chemicals

Atorvastatin drug was obtained from inceptapharmaceuticals as a gift sample. Ethanol was sourced from Merck in Germany.

Experimental Animal Procurement, Nursing, and Grouping

A total of 90 male rats weighing between 120 and 150 grams were obtained from Jahangirnagar University in Savar, Dhaka. Each of them was housed in a climate-controlled environment (temperature $25\pm 3^{\circ}\text{C}$, relative humidity $55\pm 5\%$, and a 12-h light/dark cycle) at the University of Dhaka's Institute of Nutrition & Food Science (INFS). They were given a conventional food and were permitted to drink clean water. All of the animals were maintained in this habitat for at least one week prior to the research for adaptation. 90 rats were randomly distributed into 9 groups where each group contains 10 rats.

High Fat Diet: The Levin and Dunn-Meynell composition served as the basis for adjusting the high-fat diet. The high-fat diet consists of 50% lipids, 40% carbohydrates, and 10% proteins. The dietary composition is shown in Table 1.

Table 1: Composition of high fat diet

Food Ingredients	Composition
Lipid (50%)	Milk powder (10%) Ghee (30%) Mutton fat (40%) Coconut oil (10%) Butter (10%)
Carbohydrate (40%)	Boiled rice (40%) Smashed potato (40%) Boiled corn (20%)
Protein (10%)	Dry powdered prone (40%) Dry boiled mutton (20%) Cheese (20%) Egg (20%)

After mixing the ingredients thoroughly, the high fat diet was given to the rats to induce obesity for 10 weeks. [13]

Experimental design

Rats were individually weighed and then divided into nine independent groups for research on anti-hyperlipidemic action. The distribution of rodents among the groups was based on their body weight, with each group consisting of five rats. Group 1 received normal saline and group 2 received the standard drug atorvastatin at 10mg/kg body weight. The atorvastatin control group in Table 2 shows rats that were given atorvastatin with a high-fat diet since using simply atorvastatin would result in the animals dying. The rest of the groups received different dose of plant extract at 250,500 and 750mg/kg dose. N/A indicates that rats in this group did not receive any therapeutic treatment.

Table 2: Antihyperlipidemic activity analysis

Group number	Group Status	Treatment specimen & Dose	Group Abbreviation
1	Negative Control	Physiological Saline	N
2	Positive Control	High Fat Diet	HF
3	High Fat Diet + Atrovastatin	High Fat Diet + Atrovastatin	HFD + ATV
4	High Fat Diet + <i>M. oleifera</i>	High Fat Diet+ MO ₂₅₀	HFD + MO ₂₅₀
5	High Fat Diet + <i>M. oleifera</i>	High Fat Diet + MO ₅₀₀	HFD + MO ₅₀₀
6	High Fat Diet + <i>M. oleifera</i>	High Fat Diet + MO ₇₅₀	HFD + MO ₇₅₀
7	<i>M. oleifera</i>	MO ₂₅₀	MO ₂₅₀
8	<i>M. oleifera</i>	MO ₅₀₀	MO ₅₀₀
9	<i>M. oleifera</i>	MO ₇₅₀	MO ₇₅₀

Evaluation of anti-hyperlipidemic Activity

For this experiment, 90 rats were randomly picked and equally divided into nine groups

Table 3: Application of treatment efficacy

Group Number	Group Specification	Treatment species	Dose treatment species (mg/kg)	Abbreviation of Groups
1	Negative control	Physiological saline	10 ml/kg	N
2	High Fat	N/A	N/A	HF
3	HF+At ₁₀	Atrovastatin 10mg/kg	10	At ₁₀
4	HF+MO ₂₅₀	<i>Moringa oleifera</i>	250	MO ₂₅₀
5	HF+MO ₅₀₀	<i>Moringa oleifera</i>	500	MO ₅₀₀
6	HF+MO ₇₅₀	<i>Moringa oleifera</i>	750	MO ₇₅₀

7	MO ₂₅₀	<i>Moringa oleifera</i>	250	MO ₂₅₀
8	MO ₅₀₀	<i>Moringa oleifera</i>	500	MO ₅₀₀
9	MO ₇₅₀	<i>Moringa oleifera</i>	750	MO ₇₅₀

Biological Sample Collection.

Blood was drawn from the animal as soon as its heart was punctured and transferred to a micro centrifuge tube after the killing. The samples were centrifuged at 5,000 rpm for 5 minutes to create the supernatant fluid. Biochemical testing subsequently required the transfer of this fluid to an additional micro centrifuge tube. We carefully took the kidney and liver from the animal after sacrifice and cleaned them in ice-cold saline to assess their function.

Estimation of Biochemical Parameters

For the evaluation of lipid profile total cholesterol, triglyceride, HDL and LDL level were examined. For kidney function test the level of Urea and Creatinine in blood were measured and for liver function tests SGOT and SGPT levels were evaluated. All the tests were performed by using Humaluzer 3000.

Statistical analysis:

All of our findings (raw data) in terms of numerical parameters were recorded and analyzed on a spreadsheet using the MS Excel application. The gathered data were subjected to descriptive statistics, with the findings reported as mean SD. To evaluate statistical significance, we used the SPSS 16 software's "One-way Anova test" to interpret inter-group heterogeneity in terms of several biological factors. The occurrences are considered statistically significant since the 'p' value was less than 0.05 ($p < 0.05$).

Results and discussion:

The global use of herbal medicines for addressing diverse health conditions is seeing significant growth. There is a significant increase in the acceptability and public interest in natural medicines in both emerging and established nations. These herbal remedies are now not only

accessible in pharmacy shops but also in grocery stores. For both the SGPT and the SGOT, it was seen that groups 5 and 6 exhibited statistically significant ($p < 0.05$) outcomes in the case of the SGPT. However, in the case of the SGOT, there were no statistically significant findings. There were two other investigations that came to the same conclusions [14, 15]. When conducting the renal function test, it was observed that the levels of creatinine levels were statistically significant ($p < 0.05$) in the cases of groups 4, 5, and 6 (Table 4). Also in case of urea the results were statistically significant in groups 4, 5 and 6. Two separate investigations [16, 17] came to the same conclusions about the subject which provide as the same results of our experiment. When examining the levels of high-density lipoprotein (HDL) and low-density lipoprotein (LDL), it was seen that groups 4, 5, and 6 exhibited statistically significant discoveries ($p < 0.05$) in HDL levels. On the other hand, groups 5 and 6 further showed statistically significant levels of LDL. It was determined that the triglyceride levels in Group 4 were statistically significant ($p < 0.05$). Additionally, the data obtained from groups 5 and 6 were also found to be statistically significant in the case of total cholesterol. Two further studies produced the same findings of our investigation [18, 19].

M. oleifera has many potential pathways via which it reduces serum lipid levels. Some people think that stopping 3-hydroxy-3-methylglutaryl coenzyme-A reductase (HMG-CoA reductase) from working could lower the production of cholesterol. HMG-CoA reductase is the main enzyme that makes cholesterol. MEPA may also enhance the activity of lecithin-cholesterol acyl transferase (LCAT). This enzyme has a crucial function in integrating unbound cholesterol into HDL. This will enhance the process of reverse cholesterol transport and effectively block the absorption of LDL by endothelial cells via competition [20, 21].

Table 4: Lipid profile of *Moringa oleifera*:

Groups	SGPT	SGO T	Creati- nine	Urea	TC	HDL	LDL	TG
C	36.28±4.2	36.52	0.5±0.0	29.49±	94.82±3.19	66.25±3.62	35.22±3.	54.28±3.29
5		±3.12	5	2.32			26	
HF	95.32±6.8	88.73	2.24±0.	87.72±	177.84±9.5	38.54±3.84	69.32±5.	104.24±6.26
0		±8.18	06	6.91	3		69	
HF+At ₁₀	74.59±2.0	74.33	1.1±0.0	59.51±	119.26±7.5	57.36±4.58	45.40±4.	70.64±5.94

	9	±6.28	8	6.39	6		28	
HF+MO ₂₅₀	93.28± 6.51	88.18 ±3.59	1.97±0. 08*	82.39± 4.81*	172.42±6.2 1	42.60±2.81 *	37.54±3. 57	97.29±6.82*
HF+MO ₅₀₀	90.27±6.5 8*	87.29 ±8.13	1.70±0. 07*	78.30± 3.59*	166.17±7.5 3*	47.84±5.33 *	41.66±4. 58*	93.10±5.28
HF+MO ₇₅₀	86.81±5.2 9*	85.52 ±3.61	1.29±0. 05*	74.53± 5.60*	162.23±6.1 8*	54.54±4.28 *	45.08±3. 28*	90.25±6.73
MO ₂₅₀	34.67±2.6 2	39.42 ±2.85	0.60±0. 08	28.08± 3.06	90.18±4.51	63.28±4.20	37.30±2. 21	56.29±4.10
MO ₅₀₀	36.37±4.1 2	34.73 ±2.80	0.70±0. 06	32.53± 2.84	92.30±2.81	66.53±3.29	35.18±3. 26	53.29±3.10
MO ₇₅₀	34.12±3.2 9	35.79 ±3.18	0.9±0.0 8	30.80± 2.35	94.95±3.22	65.19±4.24	38.42±4. 20	55.53± 4.23

Note: The results were expressed in Mean±SEM (standard mean error) *p< 0.05, **p< 0.01, and ***p< 0.001 were considered as statistically significant. The statistical analysis followed by one-way analysis of variance (Dunnett's test) compared to the control.

Conclusion

Within the scope of this investigation, the anti-hyperlipidemic property of *Moringaoleifera* ethanolic extract were investigated. Based on the findings of this research, it seems that an ethanol extract derived from the plant *M. oleifera* may be able to provide protection against excessive cholesterol, damage to the liver, and impaired kidney function. As a result, more research is necessary in order to determine the active components in the entire extract that have the ability to reduce hyperlipidemia. After the active chemicals have been discovered, it is possible to conduct a comprehensive investigation.

Ethical Approval

All experimental methods followed the recommendations of the Institutional Animals Ethics Committee (IEAC) with the ethical approval no (301/LUB.Pharm).

References:

1. Bhawna S, Kumar SU. Hepatoprotective activity of some indigenous plants. *Int J Pharm Tech Res.* 2009 Oct;4:1330-4.
2. Mosca S, Araújo G, Costa V, Correia J, Bandeira A, Martins E, Mansilha H, Tavares M, Coelho MP. Dyslipidemia diagnosis and treatment: risk stratification in children and adolescents. *Journal of nutrition and metabolism.* 2022 Feb 21;2022.
3. Colin, S.; Chinetti-Gbaguidi, G.; Kuivenhoven, J.A.; Staels, B. Emerging Small Molecule Drugs. *Handb. Exp. Pharmacol.* **2015**, 224, 617–630.
4. Ruscica M, Ferri N, Banach M, Sirtori CR, Corsini A. Side effects of statins: from pathophysiology and epidemiology to diagnostic and therapeutic implications. *Cardiovascular Research.* 2022 Dec 1;118(17):3288-304.
5. Baroi JA, Hossian MR, Chowdhury MM, Dolon NN, Maliha F, Rupak MA, Lima NN, Ullah MR, Tahsin R. An assessment of anti-hyperlipidemic potentialities of ethanolic extract of hemidesmusindicus in high fat induced rat model. *Asian Journal of Food Research and Nutrition.* 2023 Jul 9;2(4):323-30.
6. Mim IJ, Peza FY, Chowdhury MM, Khan TR, Mandal SK, Maliha F, Alam M, Rahman T, Tashin R. An evaluation of anti-diabetic activity of ethanolic extract of asparagus racemosus in alloxan induced rat model. *International Journal of Advances in Nephrology Research.* 2023 Aug 2;6(1):60-8.
7. Lima NN, Dolon NN, Maliha F, Ullah MR, Humayra F, Chowdhury MM, Rupak MA, Baroi JA, Shohan FS, Tashin R. An Evaluation of Analgesic and Anti-Inflammatory Activity of Ficusracemosa in Rat Model.
8. Yang L, Stöckigt J. Trends for diverse production strategies of plant medicinal alkaloids. *Natural product reports* 2010;27(10):1469-1479.
9. Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *Journal of Pharmacognosy and Phytochemistry.* 2013;1(6):168-18210. Rupak MA, Chowdhury MM, Shurov

10. Rupak MA, Chowdhury MM, Shurovi FS, Ferdous J, Tahsin MR, Sarif S, Hasan MM, Chowdhury JA, Kabir S, Chowdhury AA, Aktar F. An Evaluation of Analgesic and Anti-Inflammatory Activity of Ethanolic Extract of *Cynodon Dactylon* on Stressed Rodent Model. *Biomedical Journal of Scientific & Technical Research*. 2022;42(3):33550-7
11. Pracheta SS, Sharma V, Paliwal R, Sharma S, Yadav S, Singh L et al (2011) Chemoprotective activity of hydro-ethanolic extract of *Euphorbia nerrifolia* Linn. Leaves against DENA-induced liver carcinogenesis in mice. *Biol Med*. 3(2):36–44.
12. El-Hadary AE, Ramadan MF (2019) Antioxidant traits and protective impact of *Moringa oleifera* leaf extract against diclofenac sodium-induced liver toxicity in rats. *J Food Biochem*. 43(2):e12704
13. Abdul, N. A., & Rahmat, A. (2015). jaafar Hz. Protective Effects of Tamarillo.
14. Puspitasari D, Lestari I, Rahayuningsih CK, Christyaningsih J. The Effectiveness of *Moringa Oleifera* Leaf Extract on Hepatotoxic Case Reviewing from Cadmium, SGOT and SGPT Blood Levels in White Rats (*Rattus Norvegicus*) Induced with Cadmium (Cd). *Jurnal Teknokes*. 2022 Sep 16;15(3):137-46.
15. Islam R, Alam MJ. Evaluation of liver protective activity of *Moringa oleifera* bark extract in paracetamol induced hepatotoxicity in rats. *BioRxiv*. 2019 Jan 7:513002.
16. Saleh SS, Sarhat ER. Effects of ethanolic *Moringa oleifera* extract on melatonin, liver and kidney function tests in alloxan-induced diabetic rats. *Indian Journal of Forensic Medicine & Toxicology*. 2019 Oct 1;13(4):1015-9.
17. Nafiu AO, Akomolafe RO, Alabi QK, Idowu CO, Odujoko OO. Effect of fatty acids from ethanol extract of *Moringa oleifera* seeds on kidney function impairment and oxidative stress induced by gentamicin in rats. *Biomedicine & Pharmacotherapy*. 2019 Sep 1;117:109154.
18. Madkhali HA, Alharthy KM, Asiri MA, Ganaie MA, Ansari MN, Rehman NU, Hamad AM. *Moringa oleifera* Lam. (family Moringaceae) leaf extract attenuates high-fat diet-induced

dyslipidemia and vascular endothelium dysfunction in Wistar albino rats. *Tropical Journal of Pharmaceutical Research*. 2019;18(12):2597-604.

19. Helmy SA, Morsy NF, Elaby SM, Ghaly MA. Hypolipidemic effect of *Moringa oleifera* Lam leaf powder and its extract in diet-induced hypercholesterolemic rats. *Journal of medicinal food*. 2017 Aug 1;20(8):755-62.
20. Geetha, G., Kalavalarasariel, G.P., Sankar, V. (2011). Antidiabetic effect of *Achyranthes rubrofusca* leaf extracts on alloxan induced diabetic rats. *Pak. J. Pharm. Sci.*, 24(2), 193-199.
21. Kolawole OT, Kolawole SO, Ayankunle AA, Olaniran IO. Methanol leaf extract of *Persea americana* protects rats against cholesterol-induced hyperlipidemia. *British Journal of Medicine and Medical Research*. 2012 Mar 18;2(2):235-42.