

Genetic Variability and Selection Parameters for Yield and its Contributing Traits in Bread Wheat (*Triticum aestivum* L. em. Thell)

ABSTRACT

Ten bread wheat genotype and their all possible cross combinations excluding their reciprocals (10 parents+45 F₁s+45 F₂s) were evaluated in a randomized block design with three replications during Rabi2022-2023 at Student's Instructional Farm, C. S. Azad University of Agriculture and Technology-208002, Kanpur, U.P to get the extent of variability and related parameters to select effectively for grain yield and quality character. It is the big challenge for researchers to get high grain yield with high protein. The data was recorded on fourteen characters viz., days to heading (50%), days to maturity, plant height (cm), number of productive tillers per plant, chlorophyll content, spike length (cm), number of spikelets per spike, number of grains per spike, weight of grain per spike (g), 1000 grain weight (g), biological yield (g), harvest index (%), protein content (%) and grain yield per plant (g). The variability among parents was highest for days to maturity it ranged from 117.33 to 137.67 with the mean value 126.60. The variation in F₁ progenies was highest for plant height ranged from 85.97 to 120.53 with the mean value of 96.71. The variation in F₂ progenies was highest for plant height ranged from 83.70 to 120.20 with the mean value of 94.99. The result showed higher estimated values for phenotypic coefficient of variability(PCV) of all the fourteen characters as compared to their corresponding genotypic coefficient of variability (GCV). Highest value of PCV and GCV were noted for grain yield per plant in both generation F₁s and F₂s except GCV were observed for none of the character F₁s. High heritability estimates were recorded for all characters in both generation F₁s and F₂s. High genetic gain was observed for grain yield per plant followed by biological yield per plant in both generation F₁s and F₂s.

Keywords: Variability, heritability, genetic advance mean and range.

INTRODUCTION

Wheat (*Triticum aestivum* L.) is one of the most important cereal crop grown in different environments due to its versatile nature over the world. At global level, it's cultivated over 219.15 million ha and production of 808.44 million tons with an average productivity of 36.88 quintals per hectare. In India, it is grown in area of 30.45 million hectares with a production of 107.74 million tons and productivity of 35.37 quintals per hectare (FAO, 2022). Wheat is grown under diverse agro-climatic conditions leading to wide fluctuation in productivity from region to region. Wheat is used for both human and animal nutrition and plays an important role in the nutrition of rapidly growing populations both in our country and the world. A number of biometrical techniques are extensively used for genetic evaluation of plants. Out of them, diallel, partial diallel and line X tester analyses have been frequently used by breeders for genetic study because they provide more genetic information about the material studied. Wheat has been playing an important role in the economy of several countries. The present study was based on fourteen quantitative characters measured on ten genotypes with one check. Information on the variability was measured by estimating the genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV), heritability and genetic advance for individual quantitative character and through equilibrium

distance over the characters. In addition, the present study will also helpful in identification of high yielding genotypes with better quality. In crop improvement only the genetic components of variation are important because this is the only component which transmits to the next generation. Heritability is an important parameter in the selection process as it determines the response to selection. Heritability is used by plant breeders to enhance the precision of single field trial and /or of series of trials (Allard, 1962). Study of the estimates of heritability with genetic advance is more meaningful and reliable than considering the parameters individually.

MATERIAL AND METHODS

Ten genetically divers genotypes were crossed in all possible combinations excluding reciprocal during *Rabi* 2022-23 at Student's Instructional Farm, C. S. Azad University of Agriculture & Technology, Kanpur-208002 U.P. Basic material consisting of ten morphological diverse genotypeviz., HD 2733, DBW 187, DBW 222, HD 3086, HUW 666, K 1317, KRL 19, HI 1563, DBW 14 and K 9423. The salient features of these parental genotypes are given in Table-1.

Table 1 :Salient features of parental genotypes

Sr. No.	Name of Parents	Pedigree	Salient feature	Origin place
1.	HD 2733	ATTLA/3/TUI/CARC//CHEN/CHTO/4/ATTLA	Resistance to brown rust and tolerant to leaf blight.	IARI, N. Delhi
2.	DBW 187	NAC/TH.AC//3*PVN/3/MIRLO /BUC/4/2*PASTOR/5/KACHU/6/KACHU	Good biscuit spread factor (8.6cm), High Fe content (43.1 ppm), resistance to yellow and brown rust	IIWBR, Karnal
3.	DBW 222	KACHU/SAUAL/8/ATTLA*2/PBW65/6/PVN//CAR422/ANA/5/BOW/CROW//BUC/PVN/3/YR/4/TRAP#1/7/ATTLA/2*PASTOR	Resistant for stripe and leaf rust; Highly resistant to Karnal bunt (9.1%) and loose smut (4.9%).Chapatti quality score (7.5/10).	IIWBR, Karnal
4.	HD 3086	DBW 14/ HD 2733/HUW 468	Resistance to yellow and brown rust	IARI, N. Delhi
5.	HUW 666	-	Drought tolerance	BHU, Varanasi
6.	K1317	K 0307/K9162	Resistant to brown rust and leaf blight. Good Chapatti quality	CSAUA&T, Kanpur
7.	KRL 19	PBW 255/KRL 1-4	Salt affected Wheat Varieties Also suitable for water logging area and tolerant to yellow and brown rust	CSSRI, Karnal
8.	HI 1563	MACS 9496*2/MC10	Resistance to all the pathotypes of the three rusts at seedling stage	IARI, RS, Indore
9.	DBW 14	RAJ 3765/PBW 343	Tolerant to leaf blight	IIWBR, Karnal
10.	K 9423	HP1633/KAL/UP262	Tolerance to heat at grain development stage	CSAUA&T, Kanpur

The genotype parents were sown in a randomized complete block design (RCBD) with three replication where each replication, there were 145 rows in three tiers. Parents and F₁s were sown of single row while F₂s were sown in two row plots. The length of each line was 3 meter with row to row spacing of 22.5 cm and plant to plant spacing of 10 cm. All the recommended agronomical practices were adopted to raise the crop. For fourteen characters *viz.*, days to heading (50%), days to maturity, plant height (cm), number of productive tillers per plant, chlorophyll content, spike length (cm), number of spikelets per spike, number of grains per spike, weight of grain per spike(g), 1000 grain weight (g), biological yield (g), harvest index (%), protein content (%) and grain yield per plant (g) observation were recorded from the five randomly selected plants from each genotype HD 3086 bread wheat variety of this region were used as 'check varieties' in this experiment.

Estimation of coefficient of variability: The computation of genotypic coefficient of variability (GCV) and phenotypic coefficient of variability (PCV) was done by the formula given by **Burton and de Vane (1953)**.

$$GCV = \frac{\sqrt{V_g}}{\bar{X}} \times 100$$

Where,

V_g = Genotypic variance, and

\bar{X} = Germplasm mean

$$PCV = \frac{\sqrt{V_p}}{\bar{X}} \times 100$$

Where,

V_p = phenotypic variance, and

\bar{X} = Germplasm mean

Heritability (in narrow sense): in F₁ generation was calculated by the formula proposed by **Crumpacker and Allard (1962)**, which is as follows:

$$h^2 = \frac{\left(\frac{1}{4}\right) \hat{D}}{\left(\frac{1}{4}\right) \hat{D} + \left(\frac{1}{4}\right) \hat{H}_1 - \left(\frac{1}{4}\right) \hat{F} + \hat{E}}$$

Heritability: in F₂ generation was calculated according to the methodology proposed by **Verhalen and Murray (1969)**.

$$h^2 = \frac{\left(\frac{1}{4}\right) \hat{D}}{\left(\frac{1}{4}\right) \hat{D} + \left(\frac{1}{16}\right) \hat{H}_1 - \left(\frac{1}{8}\right) \hat{F} + \hat{E}}$$

Heritability (in per cent) = Heritability coefficient (\hat{h}^2) \times 100

Where,

\hat{h}^2 = estimate of heritability coefficient and \hat{D} , \hat{H}_1 , \hat{F} and \hat{E} are the

Genetic advance: The calculation of the estimates of genetic advance was done according to the formula given by **Robinson et al. (1949)**.

$$GA = (k) \left(\hat{h}^2\right) (\delta_{ph})$$

and genetic advance over mean of the character

$$GA (\%) = \frac{GA}{\bar{X}} \times 100$$

Where,

GA	=	estimate of genetic advance
k	=	selection differential at 5% selection intensity, i.e., 2.06
σ_{ph}	=	phenotypic standard deviation
\hat{h}^2	=	estimate of heritability coefficient (broad sense)
\bar{x}	=	mean of the character concerned

RESULT AND DISCUSSION

A. Analysis of variance (ANOVA)

The analyses of variance (ANOVA) for parents + F₁s and parents + F₂s for all the fourteen characters are presented in Table-2 and Table-3.

In case of Table-2, highly significant differences were observed among the parents for all the 14 characters. Similarly, highly significant differences were found among F₁s for all the 14 traits. It was indicated significant variability in the base material as well as the material generated subsequently, involved all possible combinations except reciprocal cross. Highly significant differences were noted among parents vs. F₁s for all characters except for days to 50% heading, chlorophyll content and weight of grain per spike which reflected significant amount of heterotic response in these attributes.

In case of Table-3, highly significant differences were observed among the parents and F₂s for all the attributes. Parents vs. F₂s also exhibited highly significant differences for all the traits except for days to maturity, chlorophyll content and weight of grain per spike.

The magnitude of variability in F₁s was higher for days to heading (50%), days to maturity, plant height, number of productive tillers per plant, spike length, weight of grains per spike, biological yield per plant and protein content in comparison to F₂s and *vice-versa* for other characters.

B. Mean and Range

The mean values and range of parents, their F₁ and F₂ progenies for all the fourteen traits are presented in Table-4.

The variability among parents was highest for days to maturity it ranged from 117.33 to 137.67 with the mean value 126.60. Mean value for plant height reported 90.94 with the range from 80.60 to 100.83. The average days to heading (50%) was 81.77 with the minimum 74.33 and maximum values 93.67. Biological yield per plant ranged from 35.53 to 51.27 with the mean value of 41.57. Harvest index ranged from 30.61 to 45.64 with the mean value 38.21 and the average grain yield per plant 15.99 ranged from 12.33 to 23.40 while number of grains/spike ranged from 30.47 to 38.1 with the mean value of 34.28. 1000 grain weight ranged from 38.98 to 44.31 with the average value of 41.72. Chlorophyll content range from 38.47 to 43.39 with the mean value of 40.56 and the average spike length 11.12 ranged from 9.27 to 13.47 while number of productive tillers/plant ranged from 6.33 to 10.00 with the mean value of 7.50. Mean number of spikelets/spike found 17.05 with the range of 15.47 to 18.87 while the

average protein content found 10.34 with the minimum and maximum values 9.22 and 11.54 respectively. Weight of grains/spike ranged from 1.53 to 1.87 with the mean value of 1.67.

The variation in F₁ progenies was highest for plant height ranged from 85.97 to 120.53 with the mean value of 96.71 followed by biological yield per plant ranged from 20.67 to 52.07 with the mean value of 38.77. Days to maturity ranged from 118.00 to 138.67 with the mean value of 125.88 and the average days to heading (50%) 81.20 ranged from 72.00 to 89.67 while grain yield per plant ranged from 8.53 to 22.87 with the mean value of 15.59. chlorophyll content ranged from 34.23 to 48.07 with the mean value of 40.27. number of grains per spike ranged value 28.53 to 41.60 with the mean value of 36.95 and the average harvest index 40.24 ranged from 36.78 to 44.98 while 1000 grain weight ranged from 39.12 to 46.12 with the mean value of 42.88. Mean number of spikelets per spike found 18.43 with the ranged of 14.80 to 21.20 while the average number of productive tillers 8.63 with the minimum and maximum values 6.00 and 11.67 respectively. Spike length ranged from 8.30 to 13.47 with the mean value of 10.63 and the average protein content 10.14 ranged from 8.81 to 11.26 while and weight of grain per spike ranged from 1.30 to 2.13 with the mean value of 1.66.

The variation in F₂ progenies was highest for plant height ranged from 83.70 to 120.20 with the mean value of 94.99 followed by biological yield/plant ranged from 26.77 to 57.00 with the mean value of 36.70. day to maturity ranged from 117.33 to 141.33 with the mean value of 126.64 and the average days to heading (50%) 80.73 ranged from 73.67 to 90.33 while chlorophyll content ranged from 34.98 to 48.71 with the mean value of 40.36. grain yield/plant ranged from 10.87 to 24.1 with the mean value of 14.84. mean number of grains/spike found 36.35 with the range of 29.13 to 41.93 while the average harvest index 40.44 with the minimum and maximum values 34.96 and 45.78 respectively. 1000 grain weight ranged from 39.65 to 46.61 with the mean value of 43.02. number of productive tiller per plant ranged from 6.00 to 12.67 with the mean value of 8.79 and the average number of spikelets per spike 18.23 ranged from 14.87 to 21.13 while spike length ranged from 8.07 to 13.4 with the mean value of 10.52., protein content ranged from 8.89 to 11.26 with the mean value of 10.02 and weight of grain per spike ranged from 1.33 to 2.27 with the mean value of 1.66.

C. Coefficients of variation

In Table-5 the estimates variability parameters i.e., genotypic coefficient of variation and phenotypic coefficient of variation were computed for all the traits having significant difference between the genotypes. GCV and PCV are categories viz., high (above 20%), moderate (above 10 and below 20%) and low (below 10 %).

In F₁ generation, at level of phenotypic coefficient of variation high values of phenotypic coefficient of variation (PCV) were observed for grain yield per plant and exhibited moderate for number of productive tiller per plant, biological yield per plant, spike length and weight of grain per spike. whereas recorded low for number of spikelets per spike, number of grains per spike, chlorophyll content, plant height, harvest index, protein content, days to heading (50 %), days to maturity and 1000 grain weight. At level of genotypic coefficient of variation, high values of genotypic coefficient of variation (GCV) were observed for none of the characters and exhibited moderate for grain yield per plant, biological yield per plant, number of productive tiller per plant, spike length and weight of grain per spike whereas recorded low for number of spikelets per spike, number of grains per spike, plant height, chlorophyll content, harvest index, protein content, days to heading (50 %), days to maturity and 1000 grain weight

In F₂ generation, at level of phenotypic coefficient of variation high values of phenotypic coefficient of variation (PCV) were observed for grain yield per plant and exhibited moderate for biological yield per plant, number of productive tiller per plant, spike length weight of grain per spike, number of grains per spike and number of spikelets per spike whereas recorded low for chlorophyll content, harvest index, plant height, protein content, days to heading (50 %), 1000 grain weight and days to maturity. At level of genotypic coefficient of variation, high values of genotypic coefficient of variation (GCV) were observed for grain yield per plant and exhibited moderate for biological yield per plant, number of productive tiller per plant, spike length and weight of grain per spike whereas recorded low for number of spikelets per spike, number of grains per spike, chlorophyll content, plant height, harvest index, protein content, days to heading (50 %), 1000 grain weight and days to maturity.

D. heritability and genetic advance

Estimates of heritability (narrow sense) and genetic advance for 14 characters, both in F₁ and F₂ generations, are presented in Table-5. There are three categories for the expression of heritability viz., high (above 30%), moderate (above 10 and below 30%) and low (below 10%) and genetic advance viz., high (above 20%), moderate (above 10 and below 20%) and low (below 10%).

(a) Heritability

In F₁ generation, high heritability estimates were recorded for all characters namely days to heading (50%), days to maturity, plant height (cm), number of productive tillers per plant, chlorophyll content (%), spike length (cm), number of spikelets per spike, number of grains per spike, weight of grain per spike(g), 1000 grain weight (g), biological yield (g), harvest index (%), protein content (%) and grain yield per plant (g).

In F₂ generation, high heritability was observed for all characters namely days to heading (50 %), days to maturity, plant height (cm), number of productive tillers/plant, chlorophyll content (%), spike length (cm), number of spikelets per spike, number of grains per spike, weight of grain per spike(g), 1000 grain weight (g), biological yield (g), harvest index (%), protein content (%) and grain yield per plant (g).

(b) Genetic advance:-

Estimates of genetic advance in per cent over mean ranged from 7.74 (1000 grain weight) to 37.88 (grain yield per plant) in F₁ generation. High genetic gain was observed for grain yield per plant followed by biological yield per plant, number of productive tiller per plant, spike length and weight of grain per spike. Moderate genetic advance was recorded for number of spikelets per spike followed by number of grains per spike, plant height, chlorophyll content, harvest index and protein content and low for days to heading followed by days to maturity and 1000 grain weight.

Estimates of genetic advance in per cent over mean ranged from 9.14 (1000 grain weight) to 40.20 (grain yield per plant) in F₂ generation. High genetic gain was observed for grain yield per plant followed by biological yield per plant, number of productive tiller per plant, and spike length. Moderate genetic advance was recorded for number of spikelets per spike followed by number of grains per spike, weight of grain per spike, plant height, chlorophyll content, harvest index, protein content and days to heading. Low for days to maturity followed by 1000 grain weight.

CONCLUSIONS

The present study generally indicated there was plenty of genetic variability among the genotypes. Thus, there is vast opportunity in the improvement of protein content and grain yield in bread wheat genotype. Therefore the result and information generated from current study need to be used by interested plant breeders. Since the experiment was carried out at one location and in one season, it is advisable to continue over several years and locations with this study for the improvement of bread wheat. Studies revealed that the days to maturity with high heritability and low genetics advance over mean followed by biological yield per plant and grain yield per plant high heritability and low genetics advance over mean in both generation F₁s and F₂s. plant height, chlorophyll content, number of spikelets per spike, number of grain per spike, harvest index and protein content were the major yield attributing characters coupled with moderate genetic advance over mean in both generation F₁s and F₂s, so emphasis should be given for these characters at the time of sample selection.

REFERENCES:

1. **Adhikari BN, Joshi BP, Shrestha J, Bhatta NR.** Genetic variability, heritability, genetic advance and trait association study for yield and yield components in advanced breeding lines of wheat. *Nepalese Journal of Agricultural Sciences.*2018;17:229-238.
2. **Ajmal SU, Zahi N, Mujahid MY.** Estimation of genetic parameters and character association in wheat. *J Agric Biol Science.*2009;1(1):15-18.
3. **Ali Y, Atta BM, Akhter J, Monneveux P, Lateef AZ.** Genetic variability, association and diversity studies in Wheat (*Triticum aestivum* L.) germplasm. *Pak J Bot* 2008;40(5):2087-2097.
4. **Allard RW.** Principals of plant breeding. John Willy and Sons Inc., USA 1962
5. **Arya VK, Singh J, Kumar L, Kumar R, Kumar P, Chand, P.**Genetic variability and diversity analysis for yield and its components in wheat (*Triticum aestivum* L.).*Ind J Agric Research.*2017;51(2):128-134.
6. **Arya VK, Singh J, Kumar L, Nagar SS, Ahalawat NK, Chand P.** Genetic variability and heritability studies in relation to grain yield and its component traits in wheat (*Triticum aestivum* L.). *Int J Agricult Stat Sci.* 2018;14: 215-222.
7. **Aycicek M, Yildirim T.**Heritability of yield and yield components in bread wheat (*Triticum aestivum* L.) genotypes. *Bangl J Bot.* 2006;35(1):17-22.
8. **Balkan A.** Genetic variability, heritability and genetic advance for yield and quality traits in M_{2,4} generations of bread wheat (*Triticum aestivum* L.) genotypes. *Turk J of Fiel Cro* 2018;23(2):173-179.
9. **Baranwal DK, Mishra VK, Vishwakarma MK, Yadav PS, Arun B.** Studies on genetic variability, correlation and path analysis for yield and yield contributing traits in wheat (*T. aestivum* L.em.Thell.).*Plant Archives.*2012;12 (1):99-104.
10. **Dedaniya AP, Pansuriya AG, Vekaria DM, Memon JT, Vekariya TA.** Combining ability analysis for yield and its components in bread wheat (*Triticum aestivum* L.). *Electronic Journal of Plant Breeding.* 2019;10(3):1005-1010.

11. **Devesh P, Moitra KP, Shukla RS, Shukla SS, Pandey S, Arya G.** Analysis of variability, heritability and genetic advance of yield, its components and quality traits in wheat. *IJAEB*. 2018; 855-859.
12. **Dhonde SH, Kute NS, Kanawade DG, Sarode ND.** Variability and characters association in wheat (*Triticum aestivum*). *Agricultural science digest*. 2000;20(2):99-101.
13. **Dutamo D, Alamerew S, Eticha F, Assefa E.** Genetic variability in bread wheat (*Triticumaestivum L.*) germplasm for yield and yield component Traits. *Journal of Biology, Agriculture andHealthcare*. 2015;5(17):140–147.
14. **Farshadfar E, Estehghari MR.** Estimation of genetic architecture for agro morphological characters in common wheat. *International Journal of Biosciences*. 2014;5(6):140-147.
15. **Firouzian A.** Heritability and genetic advance of grain yield and its related traits in wheat. *Pak. J. Bio. Science*. 2003;6(24):2020-2023.
16. **Gaur SC.** Genetic improvement through variability, heritability and genetic advance for grain yield and its contributing traits in wheat (*Triticum aestivum L.*). *Int J Pure Appl Biol*. 2019;7(1): 368-373.
17. **Ghallab KH, Sharran AAN, Al-SayedShalby NAN.** Genetic parameters for yield and yield components traits of some wheat genotypes grown in newly reclaimed soils. *International Journal of Agronomy and Agricultural Research*. 2016;9(4):1-8.
18. **Gupta SK, Verma SR.** Variability, Heritability, and genetic advance under normal and rainfed conditions in durum wheat (*Triticum durum desj.*). *Ind J Agri Res*. 2000;34(2): 122-125.
19. **Imran M, Kerkhi SA, Ali K, Singh S, Ahmad A, Singh S.** Variability heritability and genetic advance for yield and some quality traits in spring wheat (*Triticum aestivum L.*). *Plant Archives*. 2018;18, 51-56.
20. **Iqbal A, Khalil IH, ShahSMA, Kakar MS.** Estimation of heritability, genetic advance and correlation for morphological traits in spring wheat. *Sarhad Journal of Agriculture*, 2017;33(4):674-679.
21. **Jamil A, Khan S, Sayal OU, Waqas M, Ullah Q, Ali S.** Genetic variability, broad sense heritability and genetic advance studies in bread wheat (*Triticum aestivum L.*) germplasm. *Pur and Appli Biol*. 2017;6(2):538-543.
22. **Kashif M, Khaliq I.** Heritability, Correlation and Path Coefficient Analysis for Some Metric Traits in Wheat. *International Journal of Agriculture & Biology*. 2004;6(1):138–142.
23. **Khan AS, Salim I, Ali Z.** Heritability of various morphological traits in wheat. *International journal of agriculture & biology*. 2003;5(2):138-140.
24. **Korkut KZ, Başer I, Bilgin O.** Genotypic and phenotypic variability, heritability and phenotypic correlation for yield and yield components in bread wheat varieties. *Acta agronomicahungarica*. 2001;49(3):237-242.
25. **Kumar B, Singh CM, Jaiswal KK.** Genetic variability, association and diversity studies in bread wheat (*Triticum aestivum L.*). *An international quarterly journal of life sciences*. 2013;8(1):143-147.
26. **Kumar N, Markar S, Kumar V.** Studies on heritability and genetic advance estimates in timely sown bread wheat (*Triticum aestivum L.*). *BioscieDiscov*. 2014;5(1):64-69.
27. **Kumar S, Yadav RK.** Study of economic heterosis and inbreeding depression in bread wheat (*Triticum aestivum L.*) under late sown condition. *Depression*. 2016;9(9):1082-1086.

28. **Kumar S, Choudhary AM, Singh V, Chauhan MP, Yadav RDS** Studies of variability, heritability and genetic advance in some quantitative characters in bread wheat (*Triticum aestivum* L.). Journal of pharmacognosy and phytochemistry.2019;8(4):402-404.
29. **Majumder DAN, Shamsuddin AKM, Kabir MA, Hassan L.** Genetic variability, correlated response and path analysis of yield and yield contributing traits of spring wheat. J of the Bangl Agri Univer. 2008;6(2):227-234.
30. **Memon S, Iqbal N, Zaman F, Memon S, Abro MA, Jatoi GH et al.** Heritability estimates in F₂ segregating population in hexaploid wheat (*Triticum aestivum* L.). Pak J Biotechnol.2007;15(3):803-810.
31. **Novoselovic D, Baric M, Drezner G, Gunjaca J, Lalic A.** Quantitative inheritance of some wheat plant traits. Genetics and Molecular Biology. 2004;27(1):92-98.
32. **Prasad J, Dasora A, Chauhan D, Rizzardi DA, Bangarwa SK, Nesara K.** Genetic Variability, Heritability and Genetic Advance in Bread Wheat (*Triticum aestivum* L.) Genotypes. Genetics and Molecular Research.2021;20(2): 194-199.
33. **Reena R, Punia MS, Singh V.** Estimation of genetic variability parameters for various quantitative traits and rust resistance in bread wheat (*Triticum aestivum* L.). Int. J Curr Microbiol App Science.2018;7(7):1955-1966.
34. **Saleem B, Khan AS, Shahzad MT, Ijaz F.** Estimation of heritability and genetic advance for various metric traits in seven F₂ populations of bread wheat (*Triticum aestivum* L.). Journal of Agricultural Sciences.2016;61(1):1-9.
35. **Saxena M, Qureshi M, Ansari BA, Sial MA.** Genetic heritability for grain yield and its related characters in spring wheat (*Triticum aestivum* L.). Pak J Bot.2007;39(5):1503-1509.
36. **Shrimali J, Shekhawat AS, Kumari S.** Genetic variation and heritability studies for yield and yield components in barley genotypes under normal and limited moisture conditions. J of Pharmacognosy and Phytochemistry. 2017;6(4):233-235.
37. **Varsha, Verma P, Saini P, Singh V, Yashveer S.** Genetic variability of wheat (*Triticum aestivum* L.) genotypes for agro-morphological traits and their correlation and path analysis. Journal of Pharmacognosy and Phytochemistry.2019;8(4):2290-2294.
38. **Yadav N, Dashora A, Choudhary U, Dave M, Rathore J.** Assessment of genetic variability, heritability and genetic advance in bread wheat (*Triticum aestivum* L.) Genotypes. The Pharma Innovation Journal. 2021;10(9):528-530.
39. **Yaqoob, M.** Estimation of genetic variability, heritability and genetic advance for yield and yield related traits in wheat under rainfed conditions. J Agric Res. 2016;54(1):1-14.

Table-2 Analysis of variance for fourteen characters of 10 Parents and their 45 F₁s in bread wheat

Source of variation	d.f.	Days to heading (50%)	Days to maturity	Plant height (cm)	Chlorophyll content	No. of productive tillers/plant	Spike length (cm)	No. of spikelets/spike
Replication	2	3.28	7.17	5.05	2.41	0.03	0.34	0.52
Treatment	54	61.51**	109.64**	149.93**	25.90**	6.99**	5.27**	9.27**
Parents	9	111.93**	143.91**	96.82**	8.16**	4.46**	6.69**	5.81**
F ₁	44	52.42**	104.84**	145.68**	30.06**	6.95**	4.96**	9.14**
Parents vs F ₁	1	7.89	12.68**	815.79**	2.07	31.53**	5.93**	46.24**
Error	108	4.64	2.38	3.54	1.37	0.58	0.12	0.27
Source of variation	d.f.	No. of grains/spike	Weight of grain/ spike(g)	1000 grain weight (g)	Biological yield/plant (g)	Harvest index (%)	Protein content (%)	Grain yield /plant (g)
Replication	2	1.98	0.008	0.19	2.30	0.72	0.028	0.17
Treatment	54	31.79**	0.109**	10.05**	145.23**	21.96**	1.178**	27.88**
Parents	9	26.94**	0.051**	11.94**	84.27**	58.18**	1.700**	36.62**
F ₁	44	29.54**	0.124**	9.14**	156.61**	12.77**	1.076**	26.64**
Parents vs F ₁	1	174.44**	0.002	33.08**	193.55**	100.59**	0.979**	4.02*
Error	108	1.23	0.009	0.68	3.31	2.05	0.069	0.80

*, ** significant at 5% and 1% level, respectively

Table-3 Analysis of variance for fourteen characters of 10 Parents and their 45 F₂s in bread wheat.

Source of variation	d.f.	Days to heading (50%)	Days to maturity	Plant height (cm)	Chlorophyll content	No. of productive tillers/plant	Spike length (cm)	No. of spikelets/spike
Replication	2	0.77	1.82	4.97	0.97	0.48	0.08	0.35
Treatment	54	60.81**	104.45**	132.41**	28.56**	6.14**	4.99**	9.21**
Parents	9	111.93**	143.91**	96.82**	8.16**	4.46**	6.69**	5.81**
F ₂	44	51.13**	98.75**	133.56**	33.35**	5.69**	4.56**	9.34**
Parents vs F ₂	1	26.59**	0.06	402.61**	1.05	41.01**	8.74**	34.10**
Error	108	2.74	2.11	3.54	1.99	0.48	0.12	0.31
Source of variation	d.f.	No. of grains/spike)	Weight of grain/ spike(g)	1000 grain weight (g)	Biological yield/plant (g)	Harvest index (%)	Protein content (%)	Grain yield /plant (g)
Replication	2	1.88	0.051	2.12	0.34	0.73	0.045	0.06
Treatment	54	36.60**	0.093**	12.37**	143.78**	22.65**	1.092**	28.67**
Parents	9	26.94**	0.051**	11.94**	84.27**	58.18**	1.700**	36.62**
F ₂	44	37.01**	0.104**	11.79**	145.99**	13.13**	0.933**	26.96**
Parents vs F ₂	1	104.99**	0.003	41.67**	582.59**	121.64**	2.630**	32.34**
Error	108	1.36	0.009	0.42	3.01	2.74	0.090	0.74

*, ** significant at 5% and 1% level, respectively

Table-4 Mean and range of parents and their F₁s and F₂s for fourteen characters in bread wheat.

S. No.	Characters	Parent			F ₁			F ₂		
		Mean	Range		Mean	Range		Mean	Range	
1	Days to heading (50%)	81.77	74.33	93.67	81.20	72.00	89.67	80.73	73.67	90.33
2	Days to maturity	126.60	117.33	137.67	125.88	118.00	138.67	126.64	117.33	141.33
3	Plant height (cm)	90.94	80.60	100.83	96.71	85.97	120.53	94.99	83.70	120.20
4	Chlorophyll content	40.56	38.47	43.39	40.27	34.23	48.07	40.36	34.98	48.71
5	No. of productive tillers/plant	7.50	6.33	10.00	8.63	6.00	11.67	8.79	6.00	12.67
6	Spike length (cm)	11.12	9.27	13.47	10.63	8.30	13.47	10.52	8.07	13.40
7	No. of spikelets/spike	17.05	15.47	18.87	18.43	14.80	21.20	18.23	14.87	21.13
8	No. of grains/spike	34.28	30.47	38.10	36.95	28.53	41.60	36.35	29.13	41.93
9	Weight of grain/ spike(g)	1.67	1.53	1.87	1.66	1.30	2.13	1.66	1.33	2.27
10	1000 grain weight (g)	41.72	38.98	44.31	42.88	39.12	46.12	43.02	39.65	46.61
11	Biological yield/plant (g)	41.57	35.53	51.27	38.77	20.67	52.07	36.70	26.77	57.00
12	Harvest index (%)	38.21	30.61	45.64	40.24	36.78	44.98	40.44	34.96	45.78
13	Protein content (%)	10.34	9.22	11.54	10.14	8.81	11.26	10.02	8.89	11.26
14	Grain yield /plant (g)	15.99	12.33	23.40	15.59	8.53	22.87	14.84	10.87	24.10

Table-5 Grand mean, heritability (narrow sense), genetic advance and coefficient of variance for fourteen characters in bread wheat.

Characters		Mean	Min	max	Heritability (%)	GA	GA% mean	GCV (%)	PCV (%)
Days to heading (50%)	F ₁	81.30	72.00	93.67	80.35	8.04	9.89	5.36	5.97
	F ₂	80.92	73.67	93.67	87.60	8.48	10.48	5.44	5.81
Days to Maturity	F ₁	126.01	117.33	138.67	93.76	11.93	9.47	4.75	4.90
	F ₂	126.64	117.33	141.33	94.16	11.67	9.22	4.61	4.75
Plant height (cm)	F ₁	95.66	80.60	120.53	93.22	13.89	14.52	7.30	7.56
	F ₂	94.25	80.60	120.20	92.37	12.98	13.77	6.95	7.24
Chlorophyll content	F ₁	40.32	34.23	48.07	85.62	5.45	13.52	7.09	7.66
	F ₂	40.39	34.98	48.71	81.68	5.54	13.72	7.37	8.15
Number of productive tillers plant	F ₁	8.43	6.00	11.67	78.72	2.67	31.71	17.35	19.56
	F ₂	8.56	6.00	12.67	79.53	2.52	29.46	16.04	17.99
Spike length	F ₁	10.71	8.30	13.47	93.22	2.60	24.30	12.22	12.65
	F ₂	10.63	8.07	13.47	93.13	2.53	23.84	11.99	12.43
Number of spikelets per spike	F ₁	18.18	14.80	21.20	91.81	3.42	18.81	9.53	9.95
	F ₂	18.02	14.87	21.13	90.58	3.38	18.74	9.56	10.05
No. of grains per spike	F ₁	36.46	28.53	41.60	89.19	6.21	17.03	8.75	9.27
	F ₂	35.97	29.13	41.93	89.61	6.68	18.58	9.53	10.06
Weight of grain/ spike(g)	F ₁	1.66	1.30	2.13	79.68	0.34	20.29	11.03	12.36
	F ₂	1.66	1.33	2.27	74.89	0.30	17.97	10.08	11.65
1000 seeds grain weight (g)	F ₁	42.67	38.98	46.12	82.16	3.30	7.74	4.14	4.57
	F ₂	42.78	38.98	46.61	90.43	3.91	9.14	4.66	4.91
Biological yield per plant (g)	F ₁	39.28	20.67	52.07	93.45	13.70	34.87	17.51	18.11
	F ₂	37.59	26.77	57.00	93.96	13.68	36.39	18.22	18.80
Harvest index (%)	F ₁	39.87	30.61	45.64	76.37	4.64	11.63	6.46	7.39
	F ₂	40.03	30.61	45.78	70.77	4.46	11.15	6.44	7.65
Protein content (%)	F ₁	10.18	8.81	11.54	84.32	1.15	11.30	5.97	6.50
	F ₂	10.08	8.89	11.54	78.76	1.06	10.49	5.74	6.46
Grain yield per plant (g)	F ₁	15.66	8.53	23.40	91.84	5.93	37.88	19.19	20.02
	F ₂	15.05	10.87	24.10	92.65	6.05	40.20	20.27	21.06