

# Impact of Land Use and Land Cover Changes on Ecosystem Services Value in Mwanza City, Tanzania

## Abstract

Ecosystem services are vital services that support life and are the basis for human socio-economic progress. However, changes in land use and land cover (LULC) brought about by urban expansion degrade them. Thus, analysing the impact of land use and land cover (LULC) change on ecosystem service values (ESVs) is crucial for understanding and informing resource policy decisions. This study aims to analyse the impact of land use and land cover changes on ecosystem service values in Mwanza City, Tanzania. To achieve that, the benefits transfer approach was employed to analyse the changes in ESV in response to LULC. We estimated and analysed changes in ESV using satellite image datasets from 1999, 2009, and 2019. The LULC classes that were identified are vegetated land, agricultural land, waterbodies, built-up area, and bareland. The results exhibit that Mwanza City experienced significant LULC changes. While vegetated land, agricultural land, and bareland decreased by 49%, 15%, and 36%, respectively, the built-up area and water bodies increased by 568% and 48%, respectively, during the two decades. The total ESV decreased from 31.35 million US dollars to 26.3 million US dollars between 1999 and 2009 and to 23.96 million US dollars between 2009 and 2019. The waterbodies increased due to the increased volume of water in streams that expanded the floodplains, which resulted from surface runoff attributed to increased paved surfaces as more land was converted into a built-up environment upstream. The built-up area and bareland contributed nothing to ESV. However, the built-up area was the driving force behind the reduction of ESV in other LULC classes, as it was encroaching on them. The study concludes that the decrease in ESV reflects the degradation of ecosystem services due to the change in LULC. Hence, it is recommended that sustainable management of ecosystems be adhered to for the proper functioning of the earth's life-support system.

**Keywords:** Benefits transfer method, Ecosystem services, land use/cover, Tanzania, Urbanisation.

## 1. INTRODUCTION

Healthy ecosystems are vital for producing ecosystem services crucial for human survival [1]. Ecosystems generate oxygen, purify the water and air, and stabilise the climate. Organisms decompose waste, create soil for food production, and recycle nutrients for agriculture. Animals pollinate and fertilise plants, while humans use and trade these species for various purposes, including food, shelter, medicine, and aesthetics [2].

Costanza et al. [3] and Millennium Ecosystem Assessment (MEA) [4], define ecosystem services as the direct and indirect benefits of the ecosystem to people. Costanza et al. [3] grouped ecosystem services into seventeen major categories (Table 1).

**Table 1: Ecosystem Services and Functions Adopted from Costanza et al. (1997)**

	<b>Ecosystem services</b>	<b>Ecosystem functions</b>
1	Gas regulation	Regulation of atmospheric chemical composition.
2	Disturbance regulation	Capacitance, damping and integrity of ecosystem response to environmental fluctuations.
3	Climate regulation	Regulation of global temperature, precipitation, and other biologically mediated climatic processes at global or local levels
4	Water regulation	Regulation of hydrological flows.
5	Erosion control and sediment retention	Retention of soil within an ecosystem
6	Water supply	Storage and retention of water
7	Soil formation	Soil formation processes.
8	Waste treatment	Recovery of mobile nutrients and removal or breakdown of excess or xenic nutrients and compounds
9	Nutrient cycling	Storage, internal cycling, processing and acquisition of nutrients.
10	Pollination	Movement of floral gametes
11	Food production	That portion of gross primary production extractable as food.
12	Biological control	Trophic-dynamic regulations of populations.
13	Refugia	Habitat for resident and transient populations
14	Raw materials	That portion of gross primary production extractable as raw materials.
15	Genetic resources	Sources of unique biological materials and products

16	Cultural	Providing opportunities for non-commercial uses
17	Recreation	Providing opportunities for recreational activities.

**Source:** Adopted from Costanza et al. (1997)

However, more than ever, humans have changed ecosystems drastically and quickly [5], mostly to accommodate urbanisation and the world's expanding resource needs [6]. Resources such as wood, fresh water, air and water purification, controlling local and regional climates, and mitigating natural dangers are either being depleted or used in an unsustainable manner [5]. Converting landscapes (such as paving) and harvesting biological populations (such as logging), are a threat to the ongoing supply of ecosystem services [7, 4].

The rapid growth of urban areas, which are home to 56% of the world's human population as of 2022 [8], is exerting enormous pressure on ecosystem services within and around cities [7]. Urban land change has disproportionately huge consequences given the area it occupies worldwide [6]. It is projected that by 2030, the built-up area will have added to the world 1.2 million km<sup>2</sup> of urban land [9,10], concentrating pressures on the ecosystems that support those urban regions. However, much of the urban land growth is expected to be in Africa [11].

More natural ecosystems on the outskirts of cities are being converted to industrial, residential, or agricultural uses [12]. Leading to the loss of agricultural lands, habitat fragmentation, biodiversity loss, and inefficient use of natural resources [10]. The process hampers the ecosystems and the services they provide for human well-being in cities [13, 6]. This is because the changing LULC triggers the demand and supply flow of ecosystem services as well as the economic benefits of commercial, industrial, and residential land uses for urban dwellers [14].

Urbanisation affects the structure, composition, function, and efficiency of ecosystems, thereby affecting the value of ecosystem services [14, 15, 16, 17]. Its effects might be bigger than those caused by natural events, especially if ecosystem surfaces are changed so much that natural processes of succession and recovery are harmed, which makes the system less resilient [12,18,7]. When landscapes are changed for farming, ecosystem services are often swapped, with one service (like increasing crop yields) getting better at the expense of others (like losing the ability to support soil formation) [14,7; 4].

Thus, addressing the impacts of LULC changes on ecosystems in and around cities is crucial [19, 14]. However, this cannot be done if policymakers and decision-makers are not well informed on the effects of land use and land cover changes on ecosystem services in monetary terms. Because of this, measuring ESVs and their variations has attracted a lot of attention since Costanza et al. [3] published their list of biome ecological service value coefficients (LULC categories) and estimations of global ESVs [20].

Since then, several ecosystem services assessment and valuation methods have been developed, including benefits transfer, replacement cost, contingent valuation, avoided cost, travel cost, and market prices approaches [21, 14]. However, the benefits transfer method has been widely used because it is easy to apply as it "uses a given valuation study to assess a new location of similar characteristics" [21].

The benefits transfer method refers to "the transfer of original ecosystem service value estimates from an existing 'study site' or multiple study sites to an unstudied 'policy site' with similar characteristics that is being evaluated" [22]. Costanza and colleagues estimated the monetary value of 17 ecosystem services from 16 different global landscapes using the benefits transfer method. Later, they conducted case studies of more than 300 places from different parts of the world, which updated the estimations more [21].

The resultant estimates provided the foundation for using them to study other regions with similar characteristics elsewhere. Eventually, researchers have utilised this method to evaluate the impact of land use and land cover changes on ecosystem services, considering land use and land cover changes as a substitute for the supply of ecosystem services [21, 23].

Similarly, ecosystem services in Tanzania, like in any other country in the world, are vital for supporting life systems, including the socio-economic development of the people. For instance, more than 70% of the population in the country is employed in agriculture. The sector contributes approximately 25% of foreign exchange, 30% of the GDP, 65% of industrial raw materials, and 100%

of food needs [24]. But also, ecosystems serve as the main source of energy in the country, whereby 90% of households are using either firewood (69%) or charcoal (21%) for cooking [25].

With an average population growth rate of 3.2% per year, mainly attributed to natural population growth, Tanzania is among the top 8 contributors to global population growth in the world [8] and the fifth in Africa [26]. By 2050, its population is expected to exceed 140 million people [27, 26]. Such growth has some implications for the demand and supply flow of ecosystem services. This is because ecosystem services have to save additional populations in various ways. For example, the country was losing 469,000 ha of forest per year as of 2020, attributed to firewood and charcoal production, agricultural expansion, overgrazing, the development of human settlements, uncontrolled fires, the development of infrastructure and industry, timber extraction, and refugees [28].

However, of particular concern has been the rate at which towns and cities are growing in the country. For instance, in the past three decades, the urban lands of eleven towns and cities in the country grew by 480 km<sup>2</sup> [29]. Such expansion has already resulted to urban sprawl in some cities due to the lack of a clear urban policy in the country [30].

Mwanza City is among the cities in the country that are experiencing fast urban expansion. Mwanza, the second-largest city in the country, was home to 1,245,000 people in 2022 [31]. The population of the city is expected to reach 2.4 million by 2035 [32], assuming the present urban growth rate of 3% per year is maintained [33]. Yet the city is facing the development challenge of unguided urban expansion [34].

For instance, between 1999 and 2009 and between 2009 and 2019, Mwanza City grew by 107% (2179.9 hectares) and 118% (4957.53 hectares), respectively [35]. This expansion changed most of the land use and land cover for urban use [35]. Most land cover and land use turned into physical development for the construction of infrastructure, institutional buildings, commercial, industrial, and residential buildings (34).

However, since the majority of Mwanza City residents remain heavily dependent on local ecosystem services, as is the case in the country, there is a need for managing ecosystems in order to enhance services. Thus, to improve understanding and inform resource policy decisions in the management of ecosystems, it is important to quantify and analyse changes in ecosystem service values (ESVs). Therefore, the goals of this study were to (i) quantify LULC changes between the years 1999, 2009 and 2019, and (ii) analyse the economic value of ecosystem service changes (ESVs) associated with LULC dynamics in Mwanza City. The aim of the paper is to enhance understanding and inform Mwanza City authorities on the impact of land use and land cover change on ecosystem service values for sustainable ecosystem service management.


## 2. MATERIALS AND METHODS

### 2.1 Description of the study site

Mwanza City is located in northwest Tanzania, on Lake Victoria's southern shores. With a total area of 256.45 km<sup>2</sup>, of which 184.90 km<sup>2</sup> (72%) is dryland and 71.55 km<sup>2</sup> (28%) is covered by water, approximately 173 kilometres of the 184.90-kilometre dry land area are urbanised; the remaining parts are made up of farmed plains, valleys, grassy, undulating rocky hill areas, and forests [34]. It is situated between latitudes 2°15'S and 2°45'S and between longitudes 32°45'E and 33°05'E (Figure 1).

Mwanza City is divided into two administrative districts: the Nyamagana District (Mwanza City Council) and the Ilemela District (Ilemela Municipal Council). On October 1, 2012, the Mwanza City Council was split into the new Mwanza City Council (Nyamagana) and the Ilemela Municipal Council (Ilemela). According to the United Republic of Tanzania (URT) population and housing census of 2022, the Nyamagana District (MCC) had a population of 594,834 and an average household's size of 3.9, while the Ilemela District (IMC) had a population of 509,687 and an average household size of 4.0 [27].

Mwanza City has been experiencing unguided urban expansion [34], which has resulted in land use and land cover changes [35]. According to [34], most land cover and land use turned into physical development for the construction of infrastructure, institutional buildings, commercial, industrial, and residential buildings. Thus, increasing awareness about the loss of ecosystem benefits through quantification in monetary terms is potentially important as it informs environmental managers and planners.



**Figure 1: Location of the Study Site**

### 2.2 Pre-Processing and Classification of Images

The land use and cover classes were classified and analysed using Landsat 5 TM and 8 OLI satellite images. The satellite images were obtained through (<https://earthexplorer.usgs.gov/>) (Table 1). Image data pre-processing and processing were done using QGIS 3.34.1 Prizren. The Semi-Automatic Classification (SCP) plugin tool was used to do this. First, the images were georeferenced and geometrically corrected using the UTM zone 36<sup>0</sup>S coordinate system and the Datum ARC 1960. Next, the Maximum Likelihood Classification under supervised classification was used to classify LULC categories. Five LULC categories were created, namely vegetated land, cultivated land, waterbodies, built-up area and bare land (Table 2).

**Table 2: Landsat Images**


Landsat image	Sensor	Path/row	Acquisition	Resolution	Source
Landsat 8	Operational Land Imager	171/62	4 <sup>th</sup> June 2019	30m	United States Geological Survey
Landsat 5	Thematic Mapper	171/62	22 <sup>nd</sup> June 2009	30m	United States Geological Survey
Landsat 5	Thematic Mapper	171/62	8 <sup>th</sup> June 1999	30m	United States Geological Survey

### 2.3 Land Use and Land Cover Change Detection

The amount of change in LULC in hectares was calculated. Equally, the percentage change at a ten-year interval throughout the study period was as well computed (Table 4). The following formula (Eq. 1) was used to determine the percent rate of change:

$$\Delta\% = 100 \times \left( \frac{\text{final} - \text{initial}}{\text{initial}} \right) \dots \dots \dots \text{Equation 1}$$





**Figure 2: The 1999, 2009 & 2019 Classified LULC Maps**

In the period between 1999 and 2019, the findings show that the total built-up area increased by 568% (Table 4). Whereby, 4,368.3 hectares (217%) of built-up area increased between 1999 and 2009, and 7,074.27 hectares (111%) increased between 2009 and 2019 (Table 4). This increase was at the cost of other LULC types. Also, between 1999 and 2019, waterbodies increased by 48% (236.78 hectares). Conversely, between 1999 and 2019, there was a decrease of 5,214.57 hectares (49%), 4,570.51 hectares (15%), and 1,894.27 hectares (36%), in vegetated land, agricultural land, and bareland, respectively (Table 4).

**Table 4: LULC Change**

LULC classes	1999			1999-2009			2009			2009-2019			2019			2019-1999		
	Area (ha)	Area (%)	Rate of change (%)	Area (ha)	Area (%)	Rate of change (%)	Area (ha)	Area (%)	Rate of change (%)	Area (ha)	Area (%)	Rate of change (%)	Area (ha)	Area (%)	Rate of change (%)			
Vegetated land	10555.56	21	-30	7404.6	15	-28	5340.99	10	-49									
Agricultural land	30212.28	62	-2.3	29516.85	60	-13	25641.77	53	-15									
Waterbodies	499.05	1	19	593.46	1.2	24	735.83	1.5	48									
Built-up area	2014.83	5	217	6383.13	13.8	111	13457.4	28.5	568									
Bare land	5287.5	11	-12	4671.18	10	-27	3393.23	7	-36									
Total	48569.22	100	-	48569.22	100	-	48569.22	100	-									

**Source:** Computed from Satellite Images

The findings imply that within the period of 20 years, Mwanza City experienced the loss of vegetated land, agricultural land, and bareland to built-up areas. The slight increase in waterbodies was due to the increased volume of water in streams resulting from surface runoff attributed to increased paved surfaces as more land was converted into a built-up environment [42]. The increase in the built-up area corresponds to a population increase in the city. The population of Mwanza City increased from 241,923 in 2002[34 ] to 706,353in 2012, with an annual growth rate of 3%[43]. As of 2022, the total population of Mwanza City was estimated to 1,245,000 [31]. This growth heightened the demand for land to construct infrastructure, institutional buildings, commercial, industrial, and residential buildings [34].

On the other hand, within the same time period, the bareland decreased significantly. It changed to other LULC types, specifically through the removal of rocks to get building sites and quarrying rocks and sand for construction activities [44]. Mwanza City is geographically a rocky landscape, hence being dubbed "the rocky city." Thus, any physical landscape changes are easily noticeable.

### 3.2 The Impact of LULC Changes on Ecosystem Services Values

The study established that, in 1999, vegetated land (21.19 million US\$), agricultural land (2.78 million US\$), and waterbodies (7.38 million US\$) had the greatest contribution to ecosystem services as they had both higher ecosystem service values per hectare and total areas compared to the following years (Table 5). With the exception of the waterbodies which were increasing with an increase in time. In total, they contributed 31.35 million US dollars to ecosystem services (Table 5).

**Table 5: Estimated ESV for 1999**

LULC class	1999 ESV (million us \$ )			
	Area (ha)	ES Coefficient (US \$ ha-1/year)	Estimated ESV	
Vegetated land	10555.56	2007	21,185,008.92	21.19
Agricultural land	30212.28	92	2,779,529.76	2.78
Waterbodies	499.05	14785	7,378,454.25	7.38
Built-up area	2014.83	0	0	0
Bare land	5287.5	0	0	0
<b>Total</b>				31.35

However, the changes in vegetated land and agricultural land that took place between 1999 and 2009 significantly reduced their contribution to ecosystem services. Their ecosystem service value was 17.58 million US dollars in total (Table 6). This was because much of those LULC types changed to built-up areas, which was not contributing anything to ecosystem services. On the contrary, the waterbodies contributed more to the ecosystem services as the ecosystem service value grew to 8.77 million US dollars (Table 6) from 7.38 million US dollars in 1999 (Table 5). This was attributed to the increase in the total area. The increase in area may be due to the increased volume of water in streams that expanded the floodplains, which resulted from surface runoff attributed to increased paved surfaces as more land was converted into a built-up environment upstream[42].

**Table 6: Estimated ESV for 2009**

LULC class	2009 ESV (million US \$ )			
	Area (ha)	ES Coefficient (US \$ ha-1/year)	Estimated ESV	
Vegetated land	7404.6	2007	14,861,032.2	14.86
Agricultural land	29516.85	92	2,715,550.2	2.72
Waterbodies	593.46	14785	8,774,306.1	8.77
Built-up area	6383.13	0	0	0
Bare land	4671.18	0	0	0
<b>Total</b>				26.3

Similarly, the study indicates that in 2019, vegetated land and agricultural land had the lowest contribution to ecosystem service values. While the water bodies had the highest contribution to ecosystem service values. Table 7 shows that vegetated land and agricultural land had ecosystem service values of 10.72 million US dollars and 2.36 million US dollars, respectively. Compared to the past two decades, this amount was equivalent to losing ecosystem service values of 10.47 million US

dollars (49%) and 0.42 million US dollars (15%) of vegetated land and agricultural land, respectively (Table 8).

**Table 7: Estimated ESV for 2019**

LULC class	2019 ESV (million us \$ )			
	Area (ha)	ES Coefficient (US \$ ha-1/year)	Estimated ESV	
Vegetated land	5340.99	2007	10,719,366.93	10.72
Agricultural land	25641.77	92	2,359,042.84	2.36
Waterbodies	735.83	14785	10,879,246.55	10.88
Built-up area	13457.4	0	0	0
Bare land	3393.23	0	0	0
<b>Total</b>				23.96

Conversely, the contribution of waterbodies to ecosystem service value continued to rise as the acquisition of land from other LULC types continued. Table 8 indicates that between 1999 and 2019, waterbodies increased by 47% (3.5 million US dollars). Also, the total decrease in ecosystem service value over the two decades was 14.61 million US dollars (24%) (Table 8). This was attributed to a decrease in the area of vegetated land and agricultural land.

**Table 8: Changes in ESVs from 1999 to 2009, 2009 to 2019, and 1999 to 2019 Time Periods**

LULC categories	Ecosystem service values changes (million US \$ ) between study periods					
	1999-2009		2009-2019		1999-2019	
	Million US\$	Change %	Million US\$	Change %	Million US\$	Change %
Vegetated land	-6.33	-30	-4.14	-28	-10.47	-49
Agricultural land	-0.06	-2	-0.36	-13	-0.42	-15
Waterbodies	1.39	19	2.11	24	3.5	47
Built-up area	0	0	0	0	0	0
Bare land	0	0	0	0	0	0
Total	-10.55		-4.06		-14.61	
% Overall change		-16		-9		-24

#### 4. CONCLUSION

This study analysed the impact of land use and land cover changes on ecosystem service values in Mwanza City, Tanzania. The results have shown that LULC changed significantly over the two decades. However, while significant losses are noticed with the vegetated land (49%), there are gains in areal size with the built-up area (568%) during the study period. Subsequently, these LULC dynamics influenced the change in value of the ESV. While the vegetated land lost an ESV of 10.47 million US dollars, the waterbodies gained a total ESV of 3.5 million US dollars during the same time period. This study concludes that the LULC dynamics affected the value of ecosystem services over the two decades. It is recommended that findings on ESV changes and LULC dynamics provide important evidence for policymakers, suggesting decision-making and management strategies. This can be done by Mwanza City authorities through the design of sustainable ecosystem service management strategies with a great emphasis on sustainable land use management.

#### AUTHORS' CONTRIBUTIONS

Both authors read and approved the final manuscript.

#### REFERENCES

1. Adla K, Dejan K, Neira D, Dragana S. Chapter 9 - Degradation of ecosystems and loss of ecosystem services. 2022, 281-327. <https://doi.org/10.1016/B978-0-12-822794-7.00008-3>.
2. Sekercioglu CH. Ecosystem functions and services. In: Sodhi NS, Ehrlich PR, (edn). Conservation biology for all. Oxford Academic. 2010. doi: 10.1093/acprof:oso/9780199554232.003.0004.
3. Costanza R, d'Arge R, de Groot R, Farberk RS, Grasso M, Hannon B, Limburg K, Naeem S, O'Neill RV, Paruelo J, Raskin RG, Suttonk P, van den MB. The value of the world's ecosystem services and natural capital. Nature. 1997, 387.

4. Millenium Ecosystem Assessment (MEA). Ecosystems and humans'well-being: current state and trends. Milleniumecosystem assessment. Washington: Island Press; 2005.
5. Guo Z, Zhang L, Li Y. Increased dependence of humans on ecosystems services and biodiversity. Plos One. 2010, 5(10). doi:10.1371/journal.pone.0013113.
6. BaiX, McPhearsonT, CleughH, NagendraH, TongX, ZhuT,ZhuY. Linking urbanisation and the environment: conceptual and empirical advances. Annual Review of Environment and Resources. 2017; 42:215–40. <https://doi.org/10.1146/annurev-environ102016-061128>
7. Ervin D, Brown D, Chang H, Dujon V, Granek E, Shandas V, Yeakley A. Growing cities depend on ecosystem services. Solutions. 2012; 74-86. [www.thesolutionsjournal.org](http://www.thesolutionsjournal.org)
8. UN-Habitat. World cities report 2022: envisaging the future of cities. United Nations Human Settlements Programme (UN-Habitat). 2022. <https://unhabitat.org/>
9. Marcotullio PJ, Sorensen A. Editorial: Future urban worlds: theories, models, scenarios, and observations of urban spatial expansion. Front. Built Environ. 2023; 9. doi: 10.3389/fbuil.2023.1194813
10. Mahtta R, Fragkias M, Güneralp B, Mahendra A, Reba AM, Wentz EA, Seto KC. Urban land expansion: the role of population and economic growth for 300+ cities. NPJ Urban Sustain. 2022; 2, (5). <https://doi.org/10.1038/s42949-022-00048-y>
11. Angel S. Urban expansion: theory, evidence and practice. Buildings and Cities. 2023; 4(1): 124–138. <https://doi.org/10.5334/bc.348>
12. Theodorou P. The effects of urbanisation on ecological interactions. Current Opinion in Insect Science. 2023; 52. <https://doi.org/10.1016/j.cois.2022.100922>
- 13.Lapointe M, Gurney GG, Cumming GS. Urbanisation affects how people perceive and benefit from ecosystem service bundles in coastal communities of the Global South. Ecosystems and People. 2021; 17(1): 57–68. <https://doi.org/10.1080/26395916.2021.1890226>
14. Degefu MA, Argaw M, Feyisa GL, Degefa S. Dynamics of urban landscape nexus spatial dependence of ecosystem services in rapid agglomerate cities of Ethiopia. Science of the Total Environment. 2021; 798. <https://doi.org/10.1016/j.scitotenv.2021.149192>
15. KingEG., NelsonDR, McGreevyJR. Advancing the integration of ecosystem services and livelihood adaptation. Environ. Res. Lett. 2019; 14. <https://doi.org/10.1088/1748-9326/ab5519>
16. WiselySA, AlexanderK, MahlabaT, CassidyL. Linking ecosystem services to livelihoods in southern Africa. Ecosystem Services. 2018; 30: 339–341. <https://doi.org/10.1016/j.ecoser.2018.03.008>
17. Silori CS.Ecosystem services and sustainable development: opportunities and challenges. In: Ecosystem services and its mainstreaming in development planning process: SundriyalM, Dhaundiya VK, editors. Uttarakhand Science Education and Research Centre (USERC), &Bishen Singh Mahendra Pal Singh, Dehradun, India. 2015; 22-32.
18. BurtonPJ, JentschA, Walker LR. (2020). The Ecology of Disturbance Interactions. BioScience. 2020; 70: 854–870. doi:10.1093/biosci/biaa088
- 19.RotondoF, PerchinunnoP, L'AbbateS, MongelliL.Ecological transition and sustainable development: integrated statistical indicators to support public policies. Sci Rep 2022; 12. <https://doi.org/10.1038/s41598-022-23085-0>
20. Kindu M, SchneiderT, TeketayD, Knoke T. Changes of ecosystem service values in response to land use/land cover dynamics in Munessa–Shashemene landscape of the Ethiopian highlands. Science of the Total Environment 2016; 547, 137–147 <http://dx.doi.org/10.1016/j.scitotenv.2015.12.127>
- 21.NajmuddinO, Li Z, KhanR, ZhuangW. Valuation of Land-Use and Land-Cover-Based Ecosystem Services in Afghanistan-An assessment of the past and future. Land. 2022;11, 1906. <https://doi.org/10.3390/land11111906>
22. Richardson L, Loomis J, Kroeger T, Casey F. The role of benefit transfer in ecosystem service valuation. Ecological Economics. 2014;<http://dx.doi.org/10.1016/j.ecolecon.2014.02.018>
23. YiH, Güneralp B, Filippi AM, Kreuter UP, Güneralpl. Impacts of land change on ecosystem services in the San Antonio River Basin, Texas, from 1984 to 2010. Ecol. Econ.2017; 125–135. doi: 10.1016/j.ecolecon.2016.11.019
24. Mpogole H, Dimoso P,Mayaya H. Agriculture for Rural Development in Tanzania June 2020. In book: The game changer: President Magufuli'sfirst term in office. TEMA Publishers Co. Limited. 2020.
25. Doggart N, Ruhinduka R,Meshack CK, IshengomaRC, Morgan-Brown T, Abdallah JM, Spracklen DV, Sallu SM. The influence of energy policy on charcoal consumption in urban households in Tanzania. Energy for Sustainable Development. 2020; 57, 200–213. <https://doi.org/10.1016/j.esd.2020.06.002>

26. World Bank. Tanzania economic update. Clean water, bright future: The transformative impact of investing in WASH. The World Bank group | East and Southern Africa region macroeconomics, trade and investment global practice. Washington DC 20433. 2022;18. [www.worldbank.org](http://www.worldbank.org)
27. The United Republic of Tanzania (URT). The 2022 population and housing census: Administrative units population distribution report; Tanzania. Ministry of finance and planning, Tanzania National Bureau of Statistics and President's Office - finance and planning, Office of the chief government statistician, Zanzibar. 2022. <https://sensa.nbs.go.tz/>
28. Nzunda EF, Yusuph AS. Forest degradation in Tanzania: A systematic literature review. *Intech open*. 2022. doi: 10.5772/intechopen.107157
29. Sumari, NS, Ujoh F, Swai CS, Zheng M. Urban growth dynamics and expansion forms in 11 Tanzanian cities from 1990 to 2020. *International Journal of Digital Earth*. 2023; 16 (1): 1985–2001. <https://doi.org/10.1080/17538947.2023.2218114>
30. Tanzania Urbanisation Laboratory (TULab). Harnessing Urbanisation for development: roadmap for Tanzania's urban development policy. Paper for the coalition for urban transitions, London and Washington DC. 2019. <http://newclimateeconomy.net/content/cities-working-papers>
31. UN DESA (United Nations Department of Economic and Social Affairs). World population prospects 2022: Summary of results. UN DESA/POP/2021/TR/NO. 3. United Nations Department of Economic and Social Affairs. 2022.
32. Worrall L, Colenbrander S, Palmer I, Makene F, Mushi D, Mwijage J, Martine M, Godfrey N. Better urban growth in Tanzania: Preliminary exploration of the opportunities and challenges. Coalition for urban transitions. 2017. <http://newclimateeconomy.net/content/cities-working-paper>
33. UN-Habitat. Voluntary local review City of Mwanza. A review of the implementation of the sustainable development goals. Mwanza City Council. 2023. <https://unhabitat.org/2023/06>
34. Mwanza City Council (MCC). Mwanza City council strategic plan 2016/2017-2020/2021. President's office regional administration and local government. 2017.
35. Kaganga LS. Data on land use land cover changes of urban and peri-urban areas: The case of Mwanza City in Tanzania. *Current Urban Studies*. 2023; 11, 604-618. <https://doi.org/10.4236/cus.2023.114031>
36. Sharma S, Hussain S, Singh AA. Impact of land use and land cover on urban ecosystem service value in Chandigarh, India: a GIS-based analysis. *Journal of Urban Ecology*. 2023; 9 (1). <https://doi.org/10.1093/jue/juac030>
38. Kreuter UP, Harris HG, Matlock MD, Lacey RE. Change in ecosystem service values in the San Antonio area, Texas. *Ecological Economics*. 2001; 39: 333–346
39. de Groot R, Brander L, van der Ploeg S, Costanza R, Bernard F, Braat L, Christie M, Crossman N, Ghermandi A, Hein L, Hussain S, Kumar P, McVittie A, Portela R, Rodriguez LC, ten Brink P, van Beukering P. Global estimates of the value of ecosystems and their services in monetary units. *Ecosyst. Serv*. 2012; 1: 50–61.
40. Costanza R, Chichakly K, Dale V, Farber S, Finnigan D, Grigg K, Heckbert S, Kubiszewski I, Lee H, Liu S, Magnuszewski P, Maynard S, McDonald N, Mills R, Ogilvy S, Pert PL, Renz J, Wainger L, Young M, Ziegler RC. Simulation games that integrate research, entertainment, and learning around ecosystem services. *Ecosyst. Serv*. 2014; 10: 195–201. doi: 10.1016/j.gloenvcha.2014.04.002
41. Mengist W, Soromessa T, Feyisa GL. Estimating the total ecosystem services value of Eastern Afromontane Biodiversity Hotspots in response to landscape dynamics. *Environmental and Sustainability Indicators*. 2022; 14. <https://doi.org/10.1016/j.indic.2022.100178>
42. Feng B, Zhang Y, Bourke R. Urbanisation impacts on food risks based on urban growth data and coupled food models. *Natural Hazards*. 2021; 106: 613-627. <https://doi.org/10.1007/s11069-020-04480-0>
43. United Republic of Tanzania (URT). Tanzania in figures 2012. National bureau of statistics ministry of finance. 2013.
44. Joseph L. Impacts of traditional extraction of building materials on biodiversity conservation and livelihoods of residing communities in Mwanza City- Tanzania. *African Journal of Social Sciences and Humanities Research*. 2022; 5(1): 55-79. doi: 10.52589/AJSSHRPBCATPPY.