

Spectrum of genetics variations, association, path coefficient analysis in green gram (*Vigna radiata* L. wilzeck)

ABSTRACT

The high protein content and ability of *Vigna radiata* L. Wilczek, sometimes known as the mung bean, to fix nitrogen in the soil make it a highly significant pulse crop. The most challenging feature, seed yield is greatly influenced by the environment and other factors. This interpretation states that the study's goal was to evaluate the relationships between 22 distinct genotypes of mung beans in terms of yield and yield-attributing traits. Research on variations in genetics identified that there was a fair level of diversity among the 22 genotypes. Secondary branches had the highest PCV and GCV, followed by grain yield. The characters, 50% flowering, days to reproductive phase, days to maturity, plant height (cm), pods per plant, length of the pod, seeds per pod, primary branches, secondary branches, clusters per plant, grain yield (g), 100 seed weight (g) had shown high heritability combined with genetic advance as percent mean. 100 seed weight, cluster per plant, primary branches, seeds per pod, and pod length were shown as a highly significant and positive correlation. cluster per plant, plant height, seeds per pod, and reproductive phase explained the highest positive direct effect on seed yield, as shown by the analysis of path coefficients. Therefore, to increase the seed yield of green gram, these traits must be focused on in developing a breeding strategy.

Keywords: Genetic variability; Heritability; GCV; PCV; Correlation; Path analysis.

1. INTRODUCTION

One of the most significant legumes for food in South and Southeast Asia is the Greengram [*Vigna radiata* (L.) Wilczek]. It is India's third-most significant pulse crop [1]. Madhya Pradesh, Maharashtra, Uttar Pradesh, Andhra Pradesh, Karnataka, and Rajasthan are the primary regions where it is grown. The green gram is a semi-erect or upright annual herbaceous plant. The flower grows in axillary racemes with diadelphous stamens and an ovary with a long-bearded style. The leaves are trifoliate with long petioles, stipules with basal appendage, stipules minute, and leaflets entirely oval. [Thanniruet *al.* [2]]. Due to the claimed health benefits and growing popularity of Indian ethnic dishes, domestic consumption of greengram has surged recently [Datta *et al.* [3]]. Protein has a fair amount of lysine, an amino acid deficient in cereal grains. Greengram seeds include a variety of vitamins and minerals, including ascorbic acid, thiamine, riboflavin, niacin, pantothenic acid, and vitamin A. Other minerals include calcium, iron, magnesium, phosphorus, and

Comment [A.S.1]: Don't use numbers at the beginning of a sentence!

Comment [A.S.2]: Which trait? Just mention it in detail here.

Comment [A.S.3]: Please check the correct spelling! Greengram or green gram? Then write consistently from the title to the conclusion.

Comment [A.S.4]: Reference to which statement?

potassium Tang *et al.* [4]. This crop was regarded as a tough pulse crop that thrives in hot, dry climates. After harvest, the crop's cultivation provides 30 to 40 kg N/ha to the soil, increasing its fertility. As a result, roughly 25% less nitrogen needs to be applied to the next crop, which is often a cereal Mbeyagala *et al.* [5].

An effective breeding program must begin with a survey of genetic variability, including heritability, genotypic coefficient of variation (GCV), and phenotypic coefficient of variation (PCV) Kurandale *et al.* [6]. Correlation research can be used to determine the degree of interdependence between significant plant features, which is useful for choosing the right genotype. The majority of plant breeding initiatives focus on increasing yield, a complex trait that depends on numerous other traits that are further related to one another.

Consequently, the correlation study can be deemed inadequate. Finding the appropriate plant varieties for successful plant breeding projects requires determining the correlation and path coefficient between yield and yield criteria. Path analysis was done to determine how much the causal components, directly and indirectly, influenced the effective component. With these considerations in mind, the current study aims to screen greengram germplasm accessions, investigate the high-yield potential and quality traits, and examine genetic variation, heritability, genetic advance, correlation, and yield-related trait path analysis. When choosing a breeding program, heritability knowledge combined with genetic advancement is helpful. Finding optimal procedures for selecting quantitative qualities and comprehending the relationship between yield and contributing characters depend heavily on the correlation coefficient Johnson *et al.* [7].

It should be observed, still that a relationship does not always provide clear insights into the unique effects that each character has on the dependent character Tejaswini *et al.* [8] and Raja Thonta *et al.* [9]. Understandably, identifying the traits that genuinely influence seed yield would need a route analysis. Thus, path analysis is employed to quantify the impacts of attributes, both direct and indirect Desai *et al.* [10]. **This study aim to...**

Comment [A.S.5]: Please make a statement of the purpose of this study in the last paragraph of the introductory chapter

Formatted: Font color: Red

2. MATERIALS AND METHODS

In *Kharif* 2023, twenty-two genotypes of mung beans were used in the experiment, which used a Randomized Block Design with three replications at the Genetics and Plant Breeding (GPB) farm, School of Agriculture, LPU, Phagwara, Punjab. With a gap of 30 cm x 10 cm both between and within rows, each row measured 2 m. The research is located 243 meters above mean sea level at latitude 31° 19'32" N and longitude 75° 34'45" E. During the trial, every other suggested package of procedures was adhered to for healthy crop growth, and the genotypes were harvested as soon as the pods reached maturity. For data collection, ten randomly chosen plants were chosen from each genotype in each replication. Every suggested practice in the package was adhered to. Burton and De Vane's [11] study of variance includes genotypic and phenotypic coefficients of variation and heritability Hanson *et al.* [12] were all performed using standard statistical procedures. The Falconer formula was used to compute the correlations between genotype and phenotype. The path coefficient study was conducted using the Dewey and Lu- [13] recommended methodology. The characters in the present study *viz.*, days to 50% flowering (DF), days to reproductive phase (DP), days to maturity (DM), plant height (PH), number of pods per plant (NPPP), length of the pod (LP), number of seeds per pod (NSPP), number of primary branches (NPB), number of secondary branches (NSB), number of clusters per plant (NCPP), grain yield (GY), 100 seed weight (HSW).

3. RESULTS AND DISCUSSION

ANOVA, or Analysis of variance, was used to examine each of the 12 characters that were being examined. At the 0.01 level of significance, the analysis of variance showed that there was a significant difference across all genotypes, demonstrating the existence of genetic variability (Table 1). All of the features under study have their estimated variance components and broad-sense heritability displayed in (Table 2).

Comment [A.S.6]: This sentence fits better in the materials and methods section.

3.1 PCV AND GCV

The trait secondary branches had the highest PCV (26.941) and GCV (26.771); the number of seeds per pod had the highest PCV (15.942) and GCV (15.764); the grain yield had the highest PCV (15.679) and GCV (14.928). The lowest GCV (3.003) and PCV (3.498) values are indicated by the number of days before 50% blooming. For each of the 12 traits examined in this study, the phenotypic coefficient of variation (PCV) was greater than the genotypic coefficient of variation (GCV). This illustrates how a trait's expression is influenced by its surroundings. Characters with low magnitude GCV and PCV were those with days to 50% flowering, days to maturity, days to reproductive phase, and plant height; characters with moderate magnitude GCV and PCV values were those with pods per plant and pod length. The identical outcomes were observed; Rahim *et al.* [14], and Srivastava and Singh [15].

3.2 Heritability

All of the traits in the current investigation exhibited the highest heritability, which is number of secondary branches (0.987), number of seeds per pod (0.977), number of pods per plant (0.929), number of clusters per plant (0.918), 100 seed weight (0.917), grain yield (0.906), days to reproductive phase (0.886), days to maturity (0.875), number of primary branches (0.863), length of the pod (0.837), days to 50% flowering (0.737), plant height (0.621) shown in below (Table.2). The outcomes lined up with Singh *et al.* [16] for grain yield and number of pods per plant.

Comment [A.S.7]: In this section the author only mentions the results and compares with other people's research. Create a 1-5 sentence discussion about the results obtained in this study.

3.3 Correlation Studies

Studying the relationships between seed yield and its constituents as well as between the constituents is necessary for genetically improving seed yield while seed yield is a polygenic and complex characteristic that depends on several factors that contribute to yield. The directly observed characteristics are the focus of the phenotypic correlation. In this work, 22 genotypes of mung beans were examined to evaluate the genotypic and phenotypic correlations in every pair of character combinations that could exist. Since the environment's influence on a trait's expression is hidden by phenotypic correlation coefficients, genotypic correlation coefficients are thought to have high estimations Dhunde *et al.* [17]. The calculated correlation coefficient values between the various character pairings under consideration are displayed in Table 3. A useful method for elucidating the strength and scope of the relationships between significant plant characteristics is correlation coefficient analysis, which also offers fundamental standards for selection in field trials based on yield and its constituent parts Reshmi *et al.* [18]. To take full advantage of the genetic variability observed in green gram for seed production, it is essential to understand the inheritance patterns of grain yield and its closely related components Khattak *et al.* [19].

Genotypic correlation of the components characters with grain yield: Grain yield showed a positive and highly significant correlation with length of the pod (0.7270), number of seeds per pod (0.6906), number of primary branches (0.7760), number of clusters per plant and 100 seed weight, positive and significant correlation with plant height (0.3778), and number of pods per plant, while the negative and non-significant correlation with days to 50% flowering (-0.3405), days to reproductive phase (-0.0712), and days to maturity (-0.1513). Similar findings were observed; Marak. [20] and Makeen *et al.* [21].

Phenotypic correlation of the components characters with grain yield: Grain yield showed a positive and highly significant correlation with the length of the pod (0.48), number of seeds per pod (0.60), number of primary branches (0.53), number of clusters per plant (0.45), and 100 seed weight (0.63), positive and significant correlation with number of pods per plant (0.35), while negative and non-significant correlation with days to 50% flowering (-0.18), days to reproductive phase (-0.03), days to maturity (-0.12). Similar findings were observed; by Govardhan *et al.* [22], Anand *et al.* [23], and Ramachandra and Lavanya. [24]. The degree of correlation that is shown between characteristics depends on how they relate to one another as well as how genes that cause variation are separated. Some genes may result in a positive association by increasing both attributes, whereas other genes may cause a negative correlation by decreasing one and increasing the other. The competition between features for shared antecedents, which are limited, leads to negative correlations. Positive correlations, on the other hand, result from variations in the genes that supply precursors Madhur and Jinks. [25].

3.4 Path Analysis

Any breeding program's effectiveness in selecting features mostly rests on understanding the relationships between traits. The relationship between any two variables is simply identified by the correlation coefficient; it is not revealed. To determine the relative importance of the causal variables involved, the correlation coefficient can be divided into direct and indirect effects using the Path Coefficient Analysis by Shakeer *et al.* [26]. Consequently, path coefficient analysis takes into account both the direct and indirect impacts by dividing the correlation coefficient between the variables Dawane *et al.* [27]. This helps determine how much a change in one variable will affect a change in another. Eleven characteristics were considered independent characters (causes) in the path coefficient analysis, with seed yield per plant acting as the dependent character (effect). Path coefficient analysis was performed to identify features with major direct and indirect influences on grain yield, and the results were used to define the phenotypic correlation values (Table 4). The factor with the highest positive direct impact on the yield was recorded by the number of clusters per plant (0.546), plant height (0.277), number of seeds per pod (0.257), and reproductive phase (0.250). It's interesting to note that these traits likewise showed a strong positive link with yield; as a result, they ought to be taken into consideration as crucial selection factors for increasing green gram production. While, days to maturity (-0.988), number of pods per plant (-0.045), and 100 seed weight (-0.301) resulting a direct decrease in the number of seeds produced per plant. Similar findings were observed; Garjeet *et al.* [28], and Bhutia *et al.* [29].

Table 1. Analysis of variance for seed yield and its components in green gram

Sr. No	Characters	Mean sum of squares		
		Replication	Genotype	Error
1	Days to 50% flowering	1.44	6.476**	1.705
2	Days to reproductive phase	0.62	14.06**	1.601
3	Days to maturity	2.772	34.66**	4.319
4	Plant height (cm)	0.107	10.933**	4.141
5	No. of pods per plant	2.286	22.617**	1.595
6	Length of the pod (cm)	0.0028	1.341**	0.217
7	No. of seeds per pod	0.035	6.248**	0.138
8	No. of primary branches	0.378	1.410**	0.193
9	No. of secondary branches	0.132	28.109**	0.353
10	No. of clusters per plant	0.0374	3.086**	0.254
11	Grain yield (g)	0.059	6.100**	0.569
12	100 seed weight (g)	0.045	2.397**	0.198

**=significant at 5% and 1% level of probability.

Comment [A.S.8]: Usually ** is significantly different at the 1% level and * is significantly different at the 5% level. Correct this way of writing and apply it to all tables.

UNDER PEER REVIEW

Table 2. Genetic parameters of variation for seed yield and its components in greengram

Trait	DF	RP	DM	PH	NPPP	LP	NSPP	NPB	NSB	NCPP	HSW	GY
-------	----	----	----	----	------	----	------	-----	-----	------	-----	----

Sr. No	characters	Range		Mean	GCV	PCV	h ²	GA%
		Min.	Max.					
1	Days to 50% flowering	39.65	44.65	42.01	3.003	3.498	0.737	5.313
2	Days to reproductive phase	46.37	52.78	48.29	4.221	4.483	0.886	8.186
3	Days to maturity	63.55	75.74	71.02	4.479	4.787	0.875	8.633
4	Plant height (cm)	46	52.91	50.04	3.006	3.814	0.621	4.881
5	No. of pods per plant	32.57	44.08	37.74	7.01	7.272	0.929	13.923
6	Length of the pod (cm)	6.59	9.24	7.84	7.777	8.5	0.837	14.659
7	No. of seeds per pod	5.93	11.09	9.04	15.764	15.942	0.977	32.114
8	No. of primary branches	4.42	6.82	5.44	11.691	12.585	0.863	22.373
9	No. of secondary branches	5.21	15.46	11.36	26.771	26.941	0.987	54.801
10	No. of clusters per plant	6.41	10.03	8.41	11.549	12.057	0.918	22.791
11	Grain yield (g)	7.18	12.14	9.09	14.928	15.679	0.906	29.279
12	100 seed weight (g)	4.5	7.79	6.32	13.55	14.149	0.917	26.73

DF	1.0000	0.3913*	0.4577*	-0.6235**	-0.0424	-0.6225**	-0.3784*	-0.3398	-0.2214	-0.2850	0.0642	-0.3405
RP		1.0000	0.2650	-0.5169**	0.3729*	-0.1545	-0.2534	-0.4942**	0.1333	-0.0230	-0.1826	-0.0712
DM			1.0000	-0.3574*	-0.1301	0.0787	-0.4734*	-0.0303	0.2432	0.2067	-0.0783	-0.1513
PH				1.0000	-0.0726	0.3098	0.4139*	0.5447**	0.3085	0.3093	0.3905*	0.3778*
NPPP					1.0000	0.2758	0.1775	0.0187	0.1839	0.4361*	0.0528	0.3803*
LP						1.0000	0.2603	0.5801**	0.2327	0.4615*	0.3986*	0.7270**
NSPP							1.0000	0.5836**	0.1088	0.3702*	0.4947**	0.6906**
NPB								1.0000	0.2876	0.4896**	0.7947**	0.7760**
NSB									1.0000	0.5590**	0.0331	0.3208
NCCP										1.0000	0.3157	0.6375**
HSW											1.0000	0.9095**
GY												1.0000

Table 3. Genotypic correlation coefficient among characters in greengram

**=significant at 5% and 1% level of probability.




fig. 1: Phenotypic correlation coefficient

Trait	DF	RP	DM	PH	NPPP	LP	NSPP	NPB	NSB	NCPP	HSW	GY
-------	----	----	----	----	------	----	------	-----	-----	------	-----	----

DF	1.4259	0.5580	0.6526	-0.8891	-0.0605	-0.8876	-0.5396	-0.4845	-0.3157	-0.4064	0.0915	-0.3405
RP	0.0981	0.2506	0.0664	-0.1295	0.0934	-0.0387	-0.0635	-0.1238	0.0334	-0.0058	-0.0457	-0.0712
DM	-0.4524	-0.2620	-0.9884	0.3533	0.1285	-0.0778	0.4679	0.0300	-0.2404	-0.2043	0.0774	-0.1513
PH	-0.1728	-0.1433	-0.0991	0.2772	-0.0201	0.0859	0.1147	0.1510	0.0855	0.0857	0.1083	0.3778
NPPP	0.0194	-0.1708	0.0596	0.0333	-0.4579	-0.1263	-0.0813	-0.0086	-0.0842	-0.1997	-0.0242	0.3803
LP	-0.9043	-0.2245	0.1144	0.4501	0.4007	1.4528	0.3781	0.8428	0.3381	0.6705	0.5792	0.7270
NSPP	-0.0976	-0.0654	-0.1221	0.1068	0.0458	0.0671	0.2579	0.1505	0.0281	0.0955	0.1276	0.6906
NPB	-0.0514	-0.0747	-0.0046	0.0824	0.0028	0.0877	0.0882	0.1512	0.0435	0.0740	0.1202	0.7760
NSB	-0.0303	0.0183	0.0333	0.0423	0.0252	0.0319	-0.0149	0.0394	0.1370	0.0766	0.0045	0.3208
NCCP	-0.1557	-0.0126	0.1130	0.1690	0.2383	0.2522	0.2023	0.2676	0.3055	0.5465	0.1725	0.6375
HSW	-0.0194	0.0551	0.0236	-0.1178	-0.0159	-0.1202	-0.1492	-0.2397	-0.0100	-0.0952	-0.3016	0.9095

Table 4. Genotypic path coefficient showing direct and indirect effects of different contributing characters on grain yield per plant ingreengram


R SQUARE = 1.0465 RESIDUAL EFFECT = SQRT (-1 1.0465)

Trait	DF	RP	DM	PH	NPPP	LP	NSPP	NPB	NSB	NCPP	HSW	GY
DF	-0.0860	-0.0267	-0.0331	0.0425	0.0003	0.0474	0.0264	0.0242	0.0162	0.0205	-0.0029	-0.2682
RP	-0.0037	-0.0118	-0.0029	0.0049	-0.0041	0.0014	0.0028	0.0053	-0.0014	0.0005	0.0019	-0.0544
DM	0.0153	0.0097	0.0396	-0.0126	-0.0040	0.0024	-0.0174	-0.0004	0.0087	0.0071	-0.0033	-0.1384
PH	0.0233	0.0196	0.0150	-0.0472	0.0007	-0.0114	-0.0155	-0.0176	-0.0119	-0.0119	-0.0128	0.3232
NPPP	-0.0007	0.0671	-0.0193	-0.0030	0.1914	0.0443	0.0331	0.0019	0.0332	0.0766	0.0064	0.3671
LP	-0.1643	-0.0346	0.0184	0.0717	0.0690	0.2981	0.0644	0.1453	0.0648	0.1220	0.0974	0.6238
NSPP	-0.0984	-0.0767	-0.1411	0.1052	0.0554	0.0691	0.3202	0.1698	0.0340	0.1133	0.1492	0.6545
NPB	0.0597	0.0943	0.0021	-0.0788	-0.0021	-0.1032	-0.1122	-0.2116	-0.0545	-0.0969	-0.1523	0.6775
NSB	-0.0341	0.0220	0.0399	0.0455	0.0315	0.0394	0.0193	0.0467	0.1813	0.0969	0.0057	0.2999
NCPP	-0.0033	-0.0005	0.0025	0.0035	0.0055	0.0056	0.0049	0.0063	0.0073	0.0137	0.0043	0.5650
HSW	0.0239	-0.1167	-0.0594	0.1916	0.0236	0.2306	0.3287	0.5077	0.0222	0.2231	0.7054	0.7991

Table5. Phenotypic path coefficient showing direct and indirect effects of different contributing characters on grain yield per plant in greengram

R SQUARE = 0.9512 RESIDUAL EFFECT = 0.220

UNDER PEER REVIEW



Genotypical Path Diagram for GY




fig.2: Genotypical path diagram for grain yield




fig.3:Phenotypical path diagram for grain yield

4. CONCLUSION

Understanding how different yield characteristics are interconnected helps improve various other traits directly and indirectly, ultimately enhancing the targeted trait. By examining the correlation between yield and its contributing factors, such as the number of clusters per plant, number of seeds per pod, and plant height, we can determine their impact on seed yield per plant. This analysis reveals that these factors have favorable direct effects on seed yield per plant and positive correlations, making them suitable candidates for selection to boost greengram seed yield.

Comment [A.S.9]: Write down in detail what character you are referring to.

REFERENCES

1. Rishi N. Significant plant virus diseases in India and a glimpse of modern disease management technology. *Journal of General Plant Pathology*. 2009; 75: 1-18.
2. Thanniru Bhavyasri, IR Delvadiya, AV Ginoya. Breeding of physiological traits and improving the productivity of green gram (*Vigna radiata*(L.) Wilczek). *The Pharma Innovation Journal*. 2022; 11(9):2591-2601.
3. Datta S, Gangwar S, Shiv Kumar, Gupta S, Rai R, KaashyapM, Singh P, Chaturvedi SK, Singh BB. and Nadarajan N. Genetic diversity in selected Indian greengram [*Vigna radiata* (L.) Wilczek] cultivars using RAPD markers. *American Journal of Plant Sciences*. 2012;3: 1085-1091.
4. TangD, DongY, Guo N, Li L, and Ren H. Metabolomics analysis of the polyphenols in germinating mung bean (*Vigna radiata*) seeds and sprouts. *Agricultural and Food Science*. 2014; 94(8): 1639–1647.
5. Mbeyagala KE, Amayo R, Obuo JP, Abhay KP, and Abdul RW. A manual for mungbean production in Uganda. National Agricultural Research Organization (NARO), 2017, 32.
6. Kurandale R, Padol G, Suresh BG, Lavanya GR. Estimation of genetic variability and interrelationship among yield and yield related characters in Mungbean [*Vigna radiata* (L.) Wilczek]. *International Journal of Chemical Studies*. 2020;8(6):1809-1813.
7. Johnson HW, Robinson HI, and Comstock RE. Estimation of genetic and environmental variability in soybean. *Agronomy Journal*. 1955;47(3):314-318.
8. Tejaswini K, Malarvizhi D, Hemavathy AT, and Senthil N. Character association and path analysis for yield and yield contributing traits in greengram [*Vigna radiata* (L.) Wilczek]. *The Pharma Innovation Journal*. 2022;11(7):2654-2657.
9. Raju Thonta, Manoj Kumar Pandey, Rajneesh Kumar and Santhoshini. Studies on correlation and path coefficient for growth and yield attributes in green gram (*Vigna radiata*(L.) Wilczek). *The Pharma Innovation Journal*. 2023;12(6):1910- 1915.
10. Desai VK, Parmar LD, Chaudhary AR, and Chaudhary NB. Genetic Variability, Correlation, Path Coefficient and Stability Analysis for Yield and its Attributing Traits in Summer Green Gram [*Vigna radiata* (L.) International Wilczek] *Journal Accessions. of Current Microbiology and Applied Sciences*. 2020;9(06):2942- 2955.
11. Burton GW. and De Vane GM. Estimating heritability in tall Fescue (*Festuca arundinaceae*) from replicated clonal material. *Agronomy Journal*. 1953; 45: 478-481.

12. Hanson CH, Robinson HF, and Comstock RE. Biometrical studies on yield in segregating populations of Korean Lespedeza. *Agronomy Journal*. 1956; 48: 286-329.
13. Dewey DR and Lu K. A Correlation and Path coefficient analysis of components of crested wheatgrass seed production. *Agronomy Journal*. 1959; 51(9):515-518.
14. Rahim MA, Mia AA, Mahmud F, Zeba N, Afrin KS. Genetic variability, character association and genetic divergence in mung bean (*Vigna radiata* (L.) Wilczek), *Plant Omics Journal*. 2010;3(1):1-6.
15. Srivastava RL, Singh G. Genetic variability correlation and path analysis in mung bean (*Vigna radiata* (L.) Wilczek), *Journal of the Geological Society of India*. 2012;2(1):61-65.
16. Singh G, Srivastav RL, Prasad BK, Kumar R. Genetic variability and character association in mungbean (*Vigna radiata* (L.) Wilczek). *South Asian Journal of Agricultural Sciences*. 2022;2(1):04-07.
17. Dhunde BB, Devmore JP, Mahadik SG, Palshetkar MG, Vanve PB, Bhawe SG, Thorat BS. Correlation and path analysis studies on yield and its components in green gram [*Vigna radiata*(L.) Wilczek]. *The Pharma Innovation Journal*. 2021;10(1):727-30.
18. Reshmi Raj KR, Baisakh B, Tripathy SK, Lenka D, Salini K, and Mohanty MR. Studies on correlation and path analysis for yield and yield related characters in green gram (*Vigna radiata* (L.) Wilczek). *The Pharma Innovation Journal*. 2022;11(6):2392- 2395.
19. Khattak GSS, Haq MA, Ashraf M, Srinivas P. Combining ability in mungbean [*Vigna radiata* (L.) Wilczek]. Agronomic traits. *Korean Journal Crop Science*. 2001; 46:420-423.
20. Marak JK. Correlation and path analysis study on some morphological and yield parameters of mungbean (*Vigna radiata*(L.) Wilczek). *Journal of Agricultural Science and Food Technology*. 2017; 4(1): 60-65.
21. Makeen K, Abraham G, Jan A, and Singh KA. Genetic variability and correlations studies on yield and its components in mungbean [*Vigna radiata* (L.) Wilczek]. *Journal of Agronomy*. 2007; 6(1): 216-218.
22. Govardhan G, Reddy KHP, Reddy DM, Sudhakar P, Reddy BVB. Genetic variability studies for yield, yield attributes and drought tolerant traits in mung bean (*Vigna radiata* (L.) Wilczek). *Indian Journal of Pure & Applied Biosciences*. 2018;6(5): 674-9.
23. Anand G, Anandhi K. and Paulpandi VK. Genetic variability, correlation and path analysis for yield and yield components in F6 families of Greengram [*Vigna radiata* (L.) Wilczek] under rainfed condition. *Electronic Journal of Plant Breeding*. 2016; 7(2): 434-437.
24. Ramachandra P, and Lavanya GR. Variability, heritability genetic advance and correlation coefficients for yield component characters and seed yield in Greengram [*Vigna radiata* (L.) Wilczek]. *Journal of Pharmacognosy and Phytochemistry*. 2017; 6(4): 1202-1205.
25. Madhur J, Jinks C. Study of different dates of sowing on yield parameters of greengram (*V. mungo*). *Agriculture News* 1994; 12:53-56.

26. Shakeer S, Prameela S, Sharma B, Lavanya GR, Lal GM, Saikumar SS. Direct and Indirect Effects of Yield Contributing Characters on Seed Yield in Greengram (*Vigna radiata* (L.) Wilczek). *International Journal of Environment and Climate Change*. 2022;12(11):3300-10.
27. Dawane BK, Patil DK, Gite VK, Dawane JK, Kale DG. Character association and path analysis studies on seed yield and its yield attributing traits in greengram (*Vigna radiata* (L.) Wilczek). *The Pharma Innovation Journal*. 2022;11(12): 2476-2481.
28. Garje UA, Bhailume MS, Nagawade DR, Parhe SD. Genetic association and path coefficient analysis in Greengram [*Vigna radiata* (L.) Wilczek]. *Journal of Food Legumes*. 2014;27(2):151-4.
29. Bhutia P, Lal GM, Thomas N. Studies on genetic variability, correlation and path analysis in Greengram [*Vigna radiata* (L.) Wilczek] germplasm. *International Journal of Agricultural Sciences*. 2016;8(51):2267-72.

UNDER PEER REVIEW