

PREVALENCE OF MALARIA AND INTESTINAL HELMINTH CO-INFECTIONS AMONG CHILDREN OF AGE UNDER 15 YEARS IN ISUANIOCHA COMMUNITY AWKA NORTH LGA ANAMBRA STATE NIGERIA

ABSTRACT

Malaria parasites and helminth co-infections have long been recognized as major contributors to anaemia in endemic countries. The aim of this study was to assess the prevalence of malaria and helminths co-infection and to determine the associated risk factors among children of age under 15 years in Isuaniocha community Awka North LGA Anambra State Nigeria. This study investigated the prevalence of malaria and intestinal helminth co-infections among children of under fifteen years in 10 villages in Isuaniocha Community Awka North Local Government Anambra State Nigeria. The study was a cross-sectional survey of blood and faecal samples collected from 390 randomly selected under 15 years old children. The study was conducted between June and July 2018 in 10 villages in Isuaniocha. Blood and stool samples were collected from participants in a clean Ethylene Diamine Tetraacetic Acid (EDTA) and dry capped containers respectively and stored in a ice packed for transportation to the laboratory for testing. The blood sample were analyzed using microscopy (thin and thick smears) for the test for the presence of malaria parasite while the stool samples were analyzed using Kato-Katz method for the presence of helminth. The overall prevalence of malaria and helminth co-infection was 9.49%. The prevalence of malaria parasites was 38.23% and the prevalence of helminth infections was (17.44%). The prevalence of malaria parasites and helminth co-infection among children under 15 years showed that there is no significant difference ($P>0.05$) among age under 15. Children below 5 years of age were more susceptible to malaria and anaemia..

Keywords: Malaria parasites; Intestinal Helminths; Co-infection; Ethylene Diamine Tetra Acetic Acid (EDTA); Endemic countries; Isuaniocha

1. INTRODUCTION

"Malaria as been known over the years as one of the most intractable public health problems widely co-endemic with intestinal helminths infections, most especially in children who live in the endemic regions of sub-Saharan Africa resulting in a high rate of co-infection. Malaria epidemics have increased in frequency and intensity in most sub-Saharan African populations due to the emergence of drug resistance over the last two decades" [1]. While the prevalence of symptomatic malaria infection is well documented [2], there seems to be a lack of data on asymptomatic malaria infection [2]. Many of these asymptomatic infections go undetected as well as untreated while causing little or no manifestation in affected individuals. The extent of asymptomatic parasitemia could be said to be inversely related to a population's susceptibility to clinical disease [3,4], as, asymptomatic or sub-clinically symptom people are major reservoirs of infection [4,5].

The co-infection of malaria and intestinal helminths are commonly seen in sub-Saharan Africa because of the environmental factors that supports their coexistence and survival. A prominent factor that favour the transmission of soil-transmitted helminths (Geo-helminths) which makes them widely distributed in rural communities can be said to be the inadequacy or non-existent of sanitary facilities [6,7]. Some major predictors of malaria and helminth infection have been noted in previous studies, including greater susceptibility of females, lack of toilets, non-working status of parents or types of parents occupation, lower socioeconomic status, younger age, household crowding, lower level of education, religion, non-use of footwear, defecation practices, pig ownership, and lack of access to clean water [8,9]. Another possible factor is the subsistence farming, which is widely practiced in rural communities in Africa, which exposes populations to geo-helminths infections. It can be said that through a number of interventions, there has been a global decline in the prevalence of *Ascaris lumbricoides*, *Trichuris trichiura* and the hookworms (*Ancylostoma duodenale* and *Necator americanus*) in the Americas and Asia, the situation in sub-Saharan Africa remains stagnant [10]. "The health outcomes of intestinal parasitic infections include malnutrition, growth stunting, intellectual retardation, and cognitive as well as educational deficiencies" [11]. "However, the interference to nutrient uptake by intestinal helminths can lead to anaemia, malabsorption, impaired physical and mental development which poses a serious threat to children's long-term health, their education, and productivity" [12]. "Developing countries are known to harbour a considerable morbidity and mortality especially among children due to intestinal helminthiasis especially the soil transmitted helminths infections" [12]. Although, there has been an increasing interest to investigate the prevalence of malaria and helminth co-infections, their clinical consequences and the risk factors worldwide, there have been very few longitudinal community based studies that have attempted to investigate the interactions between malaria and helminth co-infections in villages in Isuaniocha community, Awka North Local Government Area Anambra State Nigeria [13][14]. Thus, this study is aimed at providing data on the prevalence of malaria and helminth co-infection in Isuaniocha Community, Awka North Local Government Area Anambra State Nigeria.

2. MATERIAL AND METHODS

2.1. The Study Area

The Study Area is Isuaniocha in Awka North Local Government Area of Anambra State, southeast Nigeria. Isuaniocha community is a rural town located at the outskirts of Awka, the capital of Anambra State. It is located approximately at latitude N6°16'14" and longitude E7°02'37". The population census of the community of study with respect to the creation of Anambra state in 1991, as projected by 3% to 2013 was 7,631 [15]. The study area is within the tropical rain forest zone and has marked wet and dry seasons. The wet season (March-October) is about 8 months while the dry season (October-February) is about 4 months. The relative humidity averages 70%, reaching 80% during rainy season with an annual rainfall of about 2000-3000 mm. The temperature ranges from 26°C-35°C in the dry season and 22°C-30°C in wet season. The community has 10 villages which are; Umuomite, Umudunu, Umueze, Otoko, Adama, Umuneri, Umuelom, Agwu Albert, Ekeagba and Ifite-isu (Fig 1). Isuaniocha is predominated mostly by the tribe called Igbos with few people from other ethnic groups in Nigeria like the Kogi's. Most of the inhabitants are farmers, with some petty traders, civil servants, students and technicians. The community has streams which serve as a source of domestic and drinking water supply to them. Isuaniocha is contingent with Mgbakwu, Awka, Achalla, and Urum.




Figure 1. Map showing villages in Isuaniocha Community, Awka North Local Government, Anambra State Nigeria. Source: Ministry of Lands and Survey

2.2. Study Population

The study population comprised of children under 15 years. The sample size was 390 children which was determined using the formula as cited by [16] as shown below:

$$n = \frac{N}{1 + Ne^2}$$

n =Size of Sample

N =Size of population

e =level of precision (usually 0.05)

2.3. Study Design

The study is a cross-sectional community-based survey and involved collection of blood and stool samples from children below 15 years to determine malaria and helminth co-infection respectively in the 10 villages in Isuaniocha community, Awka North LGA Anambra. The function of the immune system peaks at around puberty and gradually declines thereafter with advance in age [17] which is the reason for the selection of children below 15 years of age. The population sizes of the 10 villages, presence of markets, churches and the topographical nature of the communities inform the selection distribution across villages. For every household, a maximum of 2 children were sampled. Children that showed signs of fever and frequent stooling were selected while a simple random sampling was used to select children who did not show signs of fever or frequent stooling for the study. A total of 8 nurses from the primary health care center in Isuaniocha community were trained in blood and stool sample collections by a medical laboratory scientist and their services were engaged in sample collection from the selected children. Sample collection and waiting time per individual for blood samples took an average time of 30 minutes while participants were given sterile containers to collect their stool sample between 9 pm and midnight or very early in the morning immediately upon arising. Stool sample collection was done with the assistance of their parent.

2.4. Collection of Blood Samples

Blood were collected with a 5ml syringe from the participants and stored in an EDTA container and were used for the preparation of thin and thick blood films on a microscopic slide and in filling of a heparinized capillary tube for the determination of packed cell volume. Thin and thick blood smears was made on a slide and allowed to air-dried [18], and viewed under the microscope to determine the presence of malaria parasite.

2.5. Collection of Faecal Samples

A clean and well labeled stool containers were given to the caregivers with instructions on how to collect the children stool sample the next morning. This stool samples were collected between 8.00 and 10.00am and using the Kato-katz method to determine the presence of intestinal helminths eggs [19, 20].

2.6. Examination of Blood Samples for Malaria Parasites

2.6.1. Preparation of thick blood film

Two drops of the participant's blood was placed on a well labelled grease-free microscopic slide. The blood was homogenously spread in a circular motion using the edge of a spreader slide to make an even smear. The smear was allowed to air dry and placed in a slide box away from flies, dust and heat.

2.6.2. Preparation of thin blood film

A drop of the participant's blood was placed at the center of a well labeled grease-free microscopic slide. A smooth edged spreader slider was placed at an angle of 45° with the horizontal slide in front of the drop of blood. The drop of blood was allowed to run along the edge of the spreader slider. The spreader was pushed gently but firmly along the horizontal slide to make a thin smear of the blood. Care was taken to ensure even contact of the spreader and the surface of the horizontal slide. The film was allowed to air dry on a slide rack and later placed in a slide box away from flies, dust and heat.

2.6.3. Staining of the Blood films

Thick blood films was stained by holding the slide with the dried thick film facing downwards and dipped into Field's stain A for 5 seconds. The excess was drained off by touching a corner of the slide against the side of the container. The slide was washed gently in clean water for about 5 seconds and dipped into Field's Stain B for 3 seconds. The excess was drained off and the slide washed in clean water. The back was wiped clean and placed in an upright position on a draining rack for the film to air dry. The thin blood films was first fixed in absolute methanol for two minutes. Using a pipette, each film was covered with diluted Fields stain B and an equal volume of Field's stain A was added immediately and mixed well using the pipette. The film was left to stain for one minute, washed off with clean water, and placed upright on a draining rack to air dry.

2.6.4. Microscopic Examination of the Blood films

"When the stained thin and thick films were completely dried, a drop of immersion oil was added on each slide and examined under the microscopic using x100 objective lens. The thick film was used to detect the presence of malaria parasites while the thin film was used to identify the species. The results were recorded against the names and number of the participants. The intensity of malaria was recorded using the plus sign thus: Mild infection (+) or 1-10 parasites per 100 high power fields. Moderate infection (++) or 11-100 parasites per 100 high power fields. Heavy infection (+++) or 1-1000 parasites per high power fields" [19].

2.7. Examination of Faecal Samples for Helminths Parasites

Kato-Katz technique was used for the detection of helminths in the stool samples. A well labeled glass slide with the sample number and a template placed on top of it was used. 1 Gram (1.0g) of faecal sample was placed on a newspaper with a piece of nylon screen mesh pressed on top. A spatula was used to scrape the sieved faecal material through the screen mesh so that only the debris remained. Some of the sieved faeces were scrapped up to fill the hole in the template (disc). Air bubbles were avoided and excess faeces removed. Care was taken while lifting off the template (disc). After this, a piece of cellophane which had been soaked overnight in methylene blue glycerol solution was placed over the faecal sample. Then a clean slide was

placed over the top and pressed evenly downwards to spread the faeces in a circle and then the slide was carefully removed and observed under microscope using x10 and x40 objectives [20].

2.8. Determination of the Associated Risk Factors of Malaria and Helminth co-infection

A structured questionnaire were administered to all the 390 study participant with consent from their parents. The information contained include; their demography (gender, age, educational level, name of village and occupation of the parents) and behavioural hygiene such as the use of long lasting insecticide nets, household sanitary facility, licking of fingers, walking barefooted, hand washing after visit to toilet, foot wearing habit and frequency of deworming were the variables determined using the structured questionnaire. For children who could not read or write, the questionnaire were filled with the help of their parents.

2.9. Data Analysis

The data collected was entered into EPI data 3.1 statistical packages and analyzed using Statistical Package for the Social Sciences (SPSS) version 16.0. The outcome of the variables measured were; malaria, parasitaemia, helminthiasis and malaria and intestinal helminths co-infection. Descriptive statistics such as frequencies, proportions, means and confidence intervals were used to summarize the data. While the categorical data were compared using the Chi-square test or the Fisher's exact test, as appropriate, a p-value of < 0.05 of the study was considered for statistical significance.

3. RESULTS

3.1. Prevalence of malaria and intestinal helminth co-infection by gender among children under fifteen years in Isuaniocha Community Anambra

The overall prevalence of malaria and helminth co-infection was (9.49%) while the prevalence of malaria parasites was (38.23%) and the prevalence of helminth infections was (17.44%). Of the 390 participants for the study, (9.14%) of the male had malaria and intestinal helminth co-infection and the (52.49%) of female participants had malaria and intestinal helminth parasites co-infection as shown in (Table 1). The findings showed there is no significant difference ($P=.41, P>.05$) in the prevalence of malaria and helminth parasite co-infection by gender among children under fifteen in Isuaniocha Community.

Table 1. Prevalence of malaria and intestinal helminth co-infection by gender among children under fifteen years in Isuaniocha Community Anambra

Variables:	Number	Number for	Number	Number of	X ²	P-	df
Gender	Examined	malaria	for	positives for		value	
	(%)	Positive (%)	helminth	malaria and			
			Positive	helminths Co-			
			(%)	infection (%)			
Male	186 (47.69)	69(37.10)	32 (17.20)	17 (9.14)	0.679	.410	1
Female	204 (52.31)	84 (41.20)	36 (17.65)	20 (9.80)			
Total	390	153 (39.23)	68 (17.44)	37 (9.49)			

3.2. Prevalence of malaria and intestinal helminth co-infection by age among children under fifteen years in Isuaniocha Community Anambra

The prevalence of malaria and helminth co-infection was highest among the age 11 to 15 years having a prevalence of 11.38% co-infection rate as shown in (Table2). The age group with the least malaria and helminth parasite infection was found in age less or equal to 5 years. The findings showed there is no significant difference ($P=.756, P>.05$) in the prevalence of malaria and helminth co-infection by age among children under fifteen years in Isuaniocha Community.

Table 2. Prevalence of malaria and intestinal helminth co-infection by age among children under fifteen years in Isuaniocha Community Anambra

Variables: Age (in years)	Number Examined (%)	Number for malaria Positive (%)	Number for helminth Positive (%)	Number of positives for malaria and helminths co-infection (%)	X ²	P-value	df
≤5 years	45 (11.54)	18 (40.00)	14 (31.10)	4 (8.89)	0.558	.756	2
6-10 years	178 (45.64)	73 (41.00)	21 (11.80)	14 (7.87)			
11-15 years	167 (42.82)	62 (37.13)	33 (19.80)	19 (11.38)			
Total	390	153 (39.23)	68 (17.44)	37 (9.49)			

3.3. Prevalence of malaria and intestinal helminth co-infection by parents occupation among children under fifteen years in Isuaniocha Community Anambra

The three most prominent occupation of parents of the participant were trading, farming and public servants. The highest prevalence of malaria and helminth co-infections by occupation of parents was found among children whose parents were farmers (12.60%) while the least prevalence was found among children whose parents were public servants (5.80%) (Table 3). The prevalence of Malaria parasites and Helminth co-infections among children below fifteen years by parent's occupation shows that there is a significant difference ($P=.001, P<.05$)

Table 3. Prevalence of malaria and intestinal helminth co-infection by parents occupation among children under fifteen years in Isuaniocha Community Anambra

Parent's occupation	Number Examined (%)	Number of positives for malaria parasite and helminth co-infection (%)	X ²	P-value	df
Trader	139 (35.64)	10 (7.19)	13.080	.001	2
Farmer	182 (46.67)	23 (12.6)			
Public Servant	69 (27.54)	4 (5.80)			
Total	390	37			

3.4. Prevalence of malaria and intestinal helminth co-infection by the villages in Isuaniocha community among children under fifteen years

Among villages in Isuaniocha, Adama village had the highest prevalence malaria and helminth parasite co-infections (16.22%) among children below fifteen in Isuaniocha community while the least co-infection of malaria and helminth parasites infections were recorded in Umueze village (4.00%) (Table 4). The prevalence of malaria parasites and helminth co-infections among children below fifteen years by the Villages in

Isuaniocha shows that there is no significant difference ($P=.685$, $P>.05$) among children below fifteen years in Isuaniocha community.

Table 4. Prevalence of malaria and intestinal helminth co-infection by the villages in Isuaniocha community among children under fifteen years

Villages	Number Examined (%)	Number of Positive for malaria Parasite and Helminth co-infection (%)	X ²	P-value	df
Adama	37 (9.49)	6 (16.22)	6.538	.685	9
Agwu Albert	27 (6.92)	2 (7.41)			
Ekeagba	36 (9.23)	2 (5.56)			
Ifite Isu	44 (11.28)	3 (6.82)			
Otoko	34 (8.72)	4 (11.76)			
Umudunu	43 (11.03)	4 (9.30)			
Umuelom	43 (11.03)	4 (9.30)			
Umueze	50 (12.82)	2 (4.00)			
Umuneri	34 (8.72)	5 (14.71)			
Umuomite	42 (10.77)	5 (11.90)			
Total	390	37			

3.5 Associated Risk factors of Malaria and Helminth co-infections among children under fifteen years in Isuaniocha Community Anambra

The associated risk factors contributing to malaria can be seen by ownership of long-lasting insecticide nets (LLINs) and use of LLINs the previous night. A high percentage of the participant had LLINs 326 (83.59%) (Table 5) while only about 16.92% use LLINs the previous net. Of the 326 (83.59%) who said they had LLINs, 127 (38.97%) had malaria parasite. Of the 66 (16.92%) who used LLINs the previous, 18 (27.27%) had malaria parasite. There is no significant difference between the ownership of long-lasting insecticide nets (LLINs) and malaria parasite infection in children under 15 years in Isuaniocha community ($P=.803$, $P<.05$) while there is a significance difference ($P=.000$, $P<.05$) between the use of LLINs the previous night and malaria parasites infection in Isuaniocha community (Table 5a).

The risk factor for helminth infection transmission by the household sanitary facility showed that Latrine/Pit (50%) is the most used sanitary facility with (21.03%) showing helminth infections. Water Closet (38.97%) was the second most used sanitary facility with (6.58%) having helminth infection while (11.02%) of participants use bush/open defecation with (39.53%) having helminth infections (Table 5b). Children who licks their fingers showed a high risk of helminth infection (25.26%) against children who do not lick their fingers (14.92%) (Table 5b). Children who walk barefooted also showed high risk of helminth infection (26.35%) against the group who do not walk barefooted (11.98%). The risk factor of helminth infection by hand washing after visit to the toilet showed high helminth infection (24.73%) for the group that do not carry out hand washing after visit to toilet against the group that wash their hands after visit to toilet (15.15%) (Table 5b). The source of drinking water can also serve as a means of helminth infection (Table 5b) with the participants drinking from well (10.26%) having the highest helminth infection (65%) against other source of drinking water like borehole and sachet water. There is no significance difference ($P=.000$, $P<.05$) between the sources of drinking water and helminth infection. Children who dewormed in the last three months showed (20.19%) of helminth infection from the 104 (26.6%) participants examined while (16.43%) had helminth infection from the 286 (73.33%) participants examined (Table 5b). There is a significance difference ($P=.387$, $P>.05$) between deworming every three months and helminth infection.

Table 5a: Associated Risk factor of Malaria in children under 15 years in Isuaniocha Community

Risk factors	Number Examined (%)	Number of positive for malaria parasites (%)	X ²	P-value	df
Ownership of LLINs			0.062	.803	1
Yes	326 (83.59%)	127 (38.96%)			
No	64 (16.41%)	26 (41.03%)			
Use of LLINs the previous night			105.451	.000	1
Yes	66 (16.92%)	18 (27.27%)			
No	324	136 (41.98%)			

Table 5b: Associated Risk factor of helminth infections in children under 15 years in Isuaniocha Community

Risk factors	Number Examined (%)	Number of positive for helminth parasites (%)	X ²	P-value	df
Household Sanitary Facility			13.169	.000	2
Water Closet	152 (38.97%)	10 (6.58%)			
Bush/Open defecation	43 (11.02%)	17 (39.53%)			
Latrine/Pit	195 (50%)	41 (21.03%)			
Licking of Finger			4.515	.21	1
Yes	95 (24.35%)	24 (25.26%)			
No	295 (75.64%)	44 (14.92%)			
Walking barefoot			71.931	.000	1
Yes	148 (37.95%)	39 (26.35%)			
No	242 (62.05%)	29 (11.98%)			
Hand washing after visit to toilet			37.001	.034	1
Yes	297 (76.15%)	45 (15.15%)			
No	93 (23.85%)	23 (24.73%)			
Source of drinking water			71.931	.000	2
Well	40 (10.26%)	26 (65.00%)			
Borehole	275 (70.51%)	29 (10.55%)			
Sachet water	75 (19.23%)	13 (17.33%)			
Dewormed last 3 month			0.748	.387	1
Yes	104 (26.67%)	21 (20.19%)			

No

286 (73.33%) 47 (16.43%)

4. DISCUSSION

This study may be considered the first study conducted on the prevalence and malaria and helminth co-infections in Isuaniocha community among age group higher than the usual 0 to 59 months as no published record exists. The knowledge of the prevalence and distribution of malaria and helminth co-infections and associated risk factors in a given community is vital for intensive planning, designing and evaluating integrated health intervention programs [21].

This study reported a prevalence of 38.23% of malaria parasites, which is lower than the high 91.3% and 50.6% prevalence rate reported by Efunshile et al. [22] in Ilero and Kunihya et al [23] in Yola respectively. The study shows a prevalence of 17.44% for helminth infections which was higher than the observed prevalence of 6.02% in Anambra state [24] but lower compare to 52.50% gotten from Ogbaru Local Government Area of Anambra State by Uchenna et al. [25] and the recorded prevalence rates of 89% by Okereke et al. [26], 44.7% by Onyido et al. [27] and 10.18% by Mbanugo et al. [28] all in Anambra State.

The study reported that female under 15 years of age had the highest prevalence of malaria parasites 41.20% than the male under 15 years of age which was 37.10%. This disagrees with Ezeigbo et al. [29] and Kiggundu et al. [30] in Uganda, which showed that confirmed malaria was lower among girls 52.3% than boys 57.3% of that age range and also the findings of Benisheikh et al. [31] as well as Umaru and Uyaiabasi [32] which reported higher prevalence rate in males than their female counterparts but had no bearing on the effect of helminth co-infection. This difference was of significance and had no borderline. This varied finding may be explained by Abdullahi et al. [33] and Okonkwo et al. [34] who stated that “there is no scientific evidence to prove higher prevalence being related to gender as susceptibility to malaria infection is not influence by gender”. “This implies that malaria infection depends on the person’s exposure to infectious bites of mosquito vectors. However, children and pregnant women are biologically more susceptible to malaria; there are compounded gender differences due to sociocultural norms and expectations that influence patterns of exposure, decision making, and economics” [35].

This study shows that a higher prevalence of malaria parasites among age 6 years and above contradicts the usual trends of malaria epidemiological findings [36]. Children age 6 and above often show low prevalence of malaria with high prevalence of malaria found among under 5 years [22]. The study corroborates with the work of Aina et al. [37] in Lagos which shows high prevalence of malaria parasite among age 5-14 years, this may be as a result of a range of factors such as fewer numbers of under 5 years that showed up for the exercise and their clinical stage at the time of conducting the test. The high helminth infections 31.10% from the study among the under 5 years may have conferred immunity against malaria [22]. These findings strengthen the worries that regular deworming of children in endemic areas could potentially be putting the children at increased risk of malaria [38] as it was particularly seen that the protective effect of helminth co-infection was seen in the youngest children under 5 years who are at the highest risk of malaria compare to other ages, as deworming is now a routine biannual public health intervention in Nigeria [39]. It is very unlikely that permission would be granted to conduct a randomized trial of the effect of deworming on malaria risk. Furthermore, historical controls cannot be used to compare malaria risk before and after deworming in a treatment series, because malaria transmission changes with season and annual variation. However, this study raises the question of whether deworming should routinely be combined with intermittent preventive treatment of malaria in children (IPTc) [40] or other interventions aimed at reducing malaria risk.

Knowledge about disease transmission can be linked to educational level and the type of occupation of parents as this will contribute to the behavioural attributes of children. When the occupation of parent’s of the children who participated in the study were determined, based on some indicators like the ownership of long-lasting insecticide nets, use of long-lasting insecticide nets, a high prevalence of malaria parasites was found among children whose parent’s occupation were farmers 43.96% which was expected considering the low educational level of the group with a high significant difference ($P=.000$). Although it was expected that children of parents working as public servants should be more knowledgeable on the cause and prevention of malaria parasites.

Locations with much bushes around its surrounding tend to have high prevalence of malaria parasite and helminth co-infections owing to the conducive breeding environment for the female mosquitoes and the helminth parasites. A survey of the villages in the Isuaniocha community showed that villages that are far located away from the main town had high prevalence of malaria and helminth co-infection. The study showed

that the village Adama in Isuaniocha community, which had a lot of bushes around it had the highest prevalence of malaria and helminth co-infection which agrees with the work of Hotez and Kamath, [41] that rural communities with poor sanitary conditions often experience prevailing high infections.

This study recorded an overall prevalence of 17.44% for helminth infections, which is low and corroborate with the findings from Mirisho et al. [42] in Ghana which also showed a low prevalence of 17.33% compare to other studies done among children in different countries with a prevalence of 41.46% in a district in Northern Ethiopia [43] and 31% from rural Kenya [44]. The result shows that helminth parasites are highly prevalent in rural communities due to poor sanitary conditions prevailing in most of these areas [42]. This low intensity and prevalence of intestinal parasites in Isuaniocha community could be attributed to the bi-annual deworming campaigns that the government implemented in the region.

The behavioural factors which posed as a risk factor to the transmission of malaria and helminth infections was recorded to be, the ownership of LLINs and usage, household sanitary facility, hand washing after defecation, licking of fingers, walking barefooted, source of drinking water and deworming every three months. These behavioural factors may prevent or exacerbate the transmission of malaria and helminths co-infections. This work agrees with the previous studies conducted by Nkuo-Akenji et al. [45] in the Southwest Region of Cameroon and Messina et al. [46] in Congo where they reported that environmental factors like stagnant water, bushes around homes, unhygienic conditions, and high temperatures favor the growth and transmission of both malaria and soil-transmitted helminth. This is not surprising because unwashed hands and fruits contain dust contaminated with eggs of helminth and meanwhile stagnant water pools serve as breeding grounds for Anopheles mosquitos, hence making co-infection inevitable. While engaging in this behaviour as been known to reduce the transmission of malaria and helminth co-infection, studies have not been able to identify the impact of deworming every three months to the reduction of the transmission of malaria and helminth co-infection as often practiced in Nigeria.

Over all, the present study has established the malaria and helminth co-infection in Isuaniocha community in Awka North LGA Anambra State, Nigeria, but there is still scarce information on the pathogenesis of malaria and helminth co-infection and the mechanism at play in coexistence. More research is required to explain the underlying immunological mechanisms behind the high overlapping distributions of malaria and helminth in the study area.

5. CONCLUSION

This study established the prevalence of malaria and helminth co-infection in under 15 years in Isuaniocha community in Awka North LGA Anambra State, Nigeria. Children less than 5 years old were more susceptible to malaria and helminth infections owing to multiple risk factors which probably contributed to the prevalence of co-infection of malaria and helminth. In addition, the occupational status of parent is highly contributing to the transmission of malaria and helminth as children whose parents were farmers experienced high prevalence of malaria and helminth co-infection owing to multiple behavioural factors like poor hygiene status. Studies assessing the effect of helminth infections on the course of malaria are very important in targeted field of vaccine production especially for malaria and its trial in communities. The recent biannual vaccination exercise in the country may probably be a result of previous research findings on their co-infection. Well-structured longitudinal studies investigating the underlying immune mechanisms involved in malaria-helminths co-infection are very important in this regard. Effort should be made to develop a behavioural change model for communication on the use of LLINs and other interventions that will forestall absolute compliance. These results also indicate that employing the proper intervention in a combined malaria-helminth cost effective approach will yield timely results in endemic communities. The government is encouraged to engage scientist and approve studies that will explore the epidemiological impacts of malaria and helminth co-infections as well as their pathogenesis which will help future interventions.

CONSENT

Authors declare that 'written informed consent was obtained from the patient and parents.

ETHICAL APPROVAL

The study was approved by the ethical review committee of Chukwuemeka Odimegwu Ojukwu Teaching Hospital, Awka Anambra, Nigeria. An advocacy visit to the State Ministry of Health with a letter of introduction from the head of Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Anambra State was made to secure their approval. An advocacy visit with a copy of the approval letter from the State Ministry of Health was conducted to the Medical Director of the Health facilities, the community heads and parents to get their support and co-operation.

REFERENCES

1. Breman JG. The ears of the hippopotamus: manifestations, determinants, and estimates of the malaria burden, *Am. J Trop Med Hyg.* 2001;64(1-2):1-11.
2. Ojorongbe O, Adegbayi AM, Bolaji OS, Akindele AA, Adefioye OA, Adeyeba OA. Asymptomatic falciparum malaria and intestinal helminths co-infection among school children in Osogbo, Nigeria. *J Res and Med Sci.* 2011;16:680–86.
3. Bereczky S, Liljander A, Rooth I, Faraja I, Granath F, Montgomery M. Multiclonal asymptomatic *Plasmodium falciparum* infections predict a reduced risk of malaria disease in a Tanzanian population, *Microbes Infect.* 2007;9(1):103-10.
4. Mwangi TW, Fegan G, Williams TN, Kinyanjui SM, Snow RW, Marsh K. Evidence for over-dispersion in the distribution of clinical malaria episodes in children, *PLoS One.* 2008; 3(5): e2196.
5. Bousema JT, Gouagna LC, Drakeley CJ, Meutstege AM, Okech BA, Akim IN. *Plasmodium falciparum* gametocyte carriage in asymptomatic children in Western Kenya, *Malar J.* 2004;3:18-21.
6. De Silva NR, Brooker S, Hotez PZ, Montresor A, Engels D, Savioli L. Soil transmitted helminth infections: updating the global picture, *Trends Parasitol.* 2003;19(12);547–51.
7. Hung L, Vries P, Giao P, Binh T, Nam N, Kager P. Anemia, Malaria and Hookworm Infections in a Vietnamese Ethnic Minority, *Southeast Asian J. Trop. Med. Public Health.* 2005;36(4):816-21.
8. Al-Mekhlafi MH, Surin J, Atiya AS, Ariffin WA, Mahdy AKM, Che HA. Pattern and predictors of soil-transmitted helminth reinfection among aboriginal schoolchildren in rural Peninsular Malaysia, *Acta Trop.* 2008;107(2):200–4.
9. Traub RJ, Robertson ID, Irwin P, Mencke N, Thompson RCA. The prevalence, intensities and risk factors associated with geohelminth infection in tea-growing communities of Assam, India. *Trop. Med. Int. Health.* 2004;9(6):688–701.
10. Pullan R, Smith J, Jasrasaria R, Brooker S. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010, *Parasite Vect.* 2014;7(37); 19-37
11. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm, *Lancet.* 2006;367(9521):1521–32.
12. Aribodor DN, Basse SA, Yoonuan T, Sam-Wobo SO, Aribodor OB, Ugwuanyi IK Analysis of schistosomiasis and soil-transmitted helminths mixed infections among pupils in Enugu State, Nigeria: Implications for control. *Infect Dis Health* 2019;24(2):98–06. <https://doi.org/10.1016/j.idh.2018.12.003>.
13. Egwunyenga OA, Ajayi JA, Nmorsi OPG, Duhlińska-Popova DD. *Plasmodium*/intestinal helminth co-infections among pregnant Nigerian women. *Mem. Inst. Oswaldo Cruz.* 2001;96(8):1055-1059. <http://dx.doi.org/10.1590/S0074-02762001000800005>
14. Mwangi TW, Bethony JM, Brooker S. Malaria and helminth interactions in humans: an epidemiological view point. *Ann. Trop. Med. Parasitol.* 2006;100(7):551-570. . <https://dx.doi.org/10.1179/136485906X118468>
15. National Population Census, Nigerian population statistics. National Bureau of Statistics. 1991.
16. Uzoagulu AE. Practical guide to writing research project reports in tertiary institutions. Cheston Limited, Enugu; 2011;57-58
17. Hirokawa K, Utsuyama M, Kasai M, Kurashima C. Aging and Immunity. *Acta Pathol Jpn.* 1992; 42(8): 537-540. doi: 10.1111/j.1440-1827.1992.tb03103.x.PMID:1449050.

18. Cheesbrough M. Parasitological tests. In: District Laboratory Practice in Tropical Countries. Tropical Health Technologies Cambridge, 2005.
19. Idris .MA, Al-Jabri AM. Usefulness of Kato-Katz and trichrome staining as diagnostic methods for parasitic infections in clinical laboratories, Journal for Scientific Research Medical Sciences of Sultan Qaboos University Hospital, 2001;3:65–8.
20. Katz N, Chaves A, Pellegrino J. A sample device for quantitative stool thick smear technique in *Schistosoma mansoni*. Rev Inst Med Trop Sao Paulo. 1972;14(1):397-400
21. Dada-Adegbola H, Oluwatoba OA, Falade CO. Asymptomatic malaria and intestinal helminth co-infection among children in a rural community in South West Nigeria. Mala. Worl. Journ. 2013;4(18): 1-6.
22. Efunshile M, Olawale T, Stensvold CR, Kurtzhals J and König B. Epidemiological Study of the Association Between Malaria and Helminth Infections in Nigeria. Am j trop med hyg. 2015;92. 10.4269/ajtmh.14-0548.
23. Kunihya IZ, Samaila AB, Nassai I, Sarki A. and Haruna MY. Prevalence of Malaria Infection among Children Attending Specialist Hospital Yola, Adamawa State, Nigeria J Med Biol Sci.Res 2016;2 (8):136-142.
24. Aribodor OB, Ekwunife CA, Sam-Wobo SO. Aribodor DN, Ejiófor OS, Ugwuanyi IK, et al. Status of Intestinal Helminth Infection in Schools Implementing the Home-Grown School Feeding Program and the Impact of the Program on Pupils in Anambra State, Nigeria. Acta Parasitologica 2021;DOI: 10.1007/s11686-021-00429-w
25. Uchenna MO, Ifeoma ME, Ochiabuto OMTB, Obi MC, Emmanuell O, Ekelozie IS. Intestinal Parasites, Malaria and Anemia among School Children in Some Flood Affected Areas of Ogbaru Local Government Anambra State, Nigeria. Arch Clin Microbiol. 2018;9(2):80
26. Okereke JN, Obasi KO, Nwadike PO, Ezeji EU, Udebuani AC Geo-helminthes associated with geophagic pupils in selected primary schools in Oyi, Anambra State. Sci J Public Health 2015;3(5):45. <https://doi.org/10.11648/j.siph.s.2015030501>.
27. Onyido AE, Anumba JU, Ezechukwu GC, Ugha C, Umeanaeto PU, Iwueze MO. Intestinal helminth infections among primary school pupils in Nimo community, Njikoka local government area, Anambra State, Southeastern Nigeria. J Adv Res Biol Pharm Res 2016;1(4):44–8.
28. Umeh C, Mbanugo JI, Nwachukwu E. Prevalence of intestinal helminthes parasite in stools of nursery and primary schools pupils in Uga, Anambra State. Nigeria Sky J Microbiol Res 2015;3(1): 6–10
29. Ezeigbo OR, Osuagwu MC, Ezike MN, Ibegbulem ZO, Kalu S. Malaria Parasitaemia in Children Aged 1-5 Years in Aba, South Eastern Nigeria. US Open Genl. Microbiol. J. 2014; 1(1):1-6.
30. Kiggundu VL, O'Meara WP, Musoke R, Nalugoda FK, Kigozi G, Baghendaghe E, et al. High Prevalence of Malaria Parasitemia and Anemia among Hospitalized Children in Rakai, Uganda. PLoS ONE 2013 8(12): e82455. <https://doi.org/10.1371/journal.pone.0082455>
31. Benisheikh AAG, Biu AA, Awana AU, Shehu BB, Isiak MT. Epidemiology Survey of Malaria Infection among Patients attending General Out-Patient Department of Borno State Specialist Hospital Maiduguri, Borno State. J. Sci. Multidisc. Res., 2014; 6(10): 119- 23.
32. Umaru ML, Uyaiabasi GN. Prevalence of malaria in patients attending the General Hospital Makarfi, Makarfi Kaduna- State, North-Western Nigeria. Am. J. Infect. Dis. Microbiol., 2015 3(1):1-5.
33. Abdullahi K, Abubakar U, Adamu T, Daneji AI, Aliyu RU, Jiya NN, et al. Malaria in Sokoto, North Western Nigeria. Afr. J. Biochem. 2009;8(24): 7101-5.
34. Okonkwo IO, Adejuwon AO, Okerentunba PO, Frank-Peterside N. Plasmodium falciparum and HIV-1/2 coinfection among children presenting at the Out-Patient Clinic of Oni Memorial Children Hospital in Ibadan, Southwestern Nigeria. Natr. Sci. 2012;10(8): 94-100.
35. Ministry of Health Malaria Control Unit. Gender and malaria in Kenya. 2015; pp. 2-11.
36. World Health Organization. World Malaria Report 2012; Geneva: World Health Organization. 2012
37. Aina OO, Agomo CO, Olukosi YA, Okoh HI, Iwalokun BA, Egbuna KN, et al. Malariometric survey of ibeshe community in ikorodu, lagos state: dry season. Malar Res Treat. 2013;2013:487250. doi: 10.1155/2013/487250. Epub 2013 May 23. PMID: 23766924; PMCID: PMC3676961.
38. Specht S, Hoerauf A . Does helminth elimination promote or prevent malaria?. Lancet. 2007;369: 446-7. 10.1016/S0140-6736(07)60210-4.
39. Strunz E. Expanding the Deworming Program in Nigeria. Available at: <http://www.childrenwithoutworms.org/>. Accessed 2011 November 6, 2014.
40. Midzi N, Mtapuri-Zinyowera S, Sangweme D, Paul NH, Makware G, Mapingure MP, et al. Efficacy of integrated school based de-worming and prompt malaria treatment on helminthes-Plasmodium falciparum co-infections: a 33 months follow up study. BMC Int Health Hum Rights. 2011;11: 9.
41. Hotez PJ, Kamath A. Neglected Tropical Diseases in Sub-Saharan Africa: Review of Their Prevalence, Distribution, and Disease Burden. PLoS Negl Trop Dis, 2009;3, e412.<http://dx.doi.org/10.1371/journal.pntd.0000412>

42. Mirisho R, Neizer ML, Sarfo B. Prevalence of Intestinal Helminths Infestation in Children Attending Princess Marie Louise Children's Hospital in Accra, Ghana. *J Parasitol Res.* 2017;2017:8524985. doi: 10.1155/2017/8524985. Epub 2017 Sep 5. PMID: 29057116; PMCID: PMC5605797.
43. Abera A, Nibret E. Prevalence of gastrointestinal helminthic infections and associated risk factors among schoolchildren in Tilili town, northwest Ethiopia. *Asian P J Trop Med.* 2014;7(7):525–30. doi: 10.1016/S1995-7645(14)60088-2.
44. Nguhiu PN, Kariuki HC, Magambo JK, Kimani G, Mwatha JK, Muchiri E et al. Intestinal polyparasitism in a rural Kenyan community. *East Afr Med J.* 2009;86(6):272–8.
45. Nkuo-Akenji TK, Chi PC, Cho JF, Ndamukong KK, Sumbele I. Malaria and helminthes co-infection in children living in a malaria endemic setting of mount Cameroon and predictors of anemia, *J Parasitol.* 2006;92(6):1191-5
46. Messina JP, Taylor SM, Meshnick SR, Linke AM, Tshetu AK, Atua B. et al. "Population, behavioural and environmental drivers of malaria prevalence in the Democratic Republic of Congo," *Malaria Journal*,2011;10(1): 161.

DEFINITIONS, ACRONYMS, ABBREVIATIONS

Asymptomatic (sub-clinical) malaria infection: refers to the presence of malaria parasites in the blood without symptoms, usually provides a reservoir for transmission, and is an antecedent to symptomatic malaria

Parasitemia: Parasitaemia is the quantitative content of parasites in the blood. It is used as a measurement of parasite load in the organism and an indication of the degree of an active parasitic infection

Morbidity and Mortality: Morbidity is when you have a specific illness or condition. Some examples of common morbidities are heart disease, diabetes, and obesity while mortality is the number of deaths due to a specific illness or condition

Ethylene Diamine Tetra Acetic Acid (EDTA): It is an anticoagulant for blood samples for complete blood count (CBC), also known as a full blood count (FBC), where the EDTA chelates the calcium present in the blood specimen, arresting the coagulation process and preserving blood cell morphology.